

# Crocins protect retinal ganglion cells against H<sub>2</sub>O<sub>2</sub>-induced damage through the mitochondrial pathway and activation of NF- $\kappa$ B

BOCHANG LV<sup>1</sup>, TAO CHEN<sup>2</sup>, ZHIGUO XU<sup>1</sup>, FUQUAN HUO<sup>3</sup>, YANYAN WEI<sup>2</sup> and XINGUANG YANG<sup>1</sup>

<sup>1</sup>Shaanxi Ophthalmic Medical Center, Xi'an No. 4 Hospital, Affiliated Guangren Hospital, School of Medicine, Xi'an Jiaotong University, Xi'an, Shaanxi 710004; <sup>2</sup>Department of Anatomy, Histology and Embryology,

K.K. Leung Brain Research Centre, The Fourth Military Medical University, Xi'an, Shaanxi 710032;

<sup>3</sup>Department of Physiology and Pathophysiology, Key Laboratory of Environment and Genes Related to Diseases, Ministry of Education, Xi'an Jiaotong University School of Medicine, Xi'an, Shaanxi 710061, P.R. China

Received February 10, 2015; Accepted October 26, 2015

DOI: 10.3892/ijmm.2015.2418

**Abstract.** Glaucoma is a degenerative nerve disorder that results in irreversible blindness. It has been reported that the apoptosis of retinal ganglion cells (RGCs) is a hallmark of glaucoma. Oxidative stress is one of the major factors that cause apoptosis of RGCs. Crocin has many beneficial effects, including antioxidant and anti-apoptotic actions. However, the mechanism by which crocin protects against oxidative stress-induced damage to RGCs remains unclear. The present study aimed to investigate the mechanism by which crocin protects RGC-5 cells against H<sub>2</sub>O<sub>2</sub>-induced damage. H<sub>2</sub>O<sub>2</sub> was used to establish a model of oxidative stress injury in RGC-5 cells to mimic the development of glaucoma *in vitro*. Different concentrations (0.1 and 1  $\mu$ M) of crocin were added to test whether crocin was capable of protecting RGCs from H<sub>2</sub>O<sub>2</sub>-induced damage. WST-1, lactic dehydrogenase (LDH) release and Annexin V/FITC assays were then performed. Levels of reactive oxygen species (ROS) were detected using a ROS assay kit, mitochondrial membrane potential ( $\Delta\Psi$ m) was analyzed by JC-1 staining, caspase-3 activity was examined using a Caspase-3 assay kit, and the protein levels of Bax, Bcl-1 and cytochrome *c* were measured using western blot analysis. In addition, the protein level of phosphorylated nuclear factor- $\kappa$ B (p-NF- $\kappa$ B)

p65 was also evaluated using western blot analysis. The results showed that crocin protected RGC-5 cells from apoptosis, decreased LDH release and enhanced cell viability. Additional experiments demonstrated that crocin decreased ROS levels, increased  $\Delta\Psi$ m, downregulated the protein expression of Bax and cytochrome *c*, promoted Bcl-2 protein expression and activated NF- $\kappa$ B. Taken together, the findings of this study indicate that crocin prevented H<sub>2</sub>O<sub>2</sub>-induced damage to RGCs through the mitochondrial pathway and activation of NF- $\kappa$ B.

## Introduction

Glaucoma is a degenerative nerve disorder, which is characterized by optic atrophy and visual field defects, and results in irreversible blindness. It has been reported that almost 64.3 million patients suffered from glaucoma worldwide in 2013, and this number is likely to increase to 76 million in 2020 and 111.8 million in 2040 (1). The hallmark of glaucoma is the apoptosis of retinal ganglion cells (RGCs), which are the only efferent neurons that convey visual signals from the retina to the brain. Several risk factors, including elevated intraocular pressure (2), oxidative stress (3), elevated glutamate (4) and aging (5), have been considered to accelerate RGC apoptosis in glaucoma, among which, oxidative stress is considered as the final common pathway in glaucoma (6). Several studies have reported that oxidative stress can lead to apoptosis of RGCs through activation of the mitochondrial pathway (7), and that apoptosis is the main cause of RGC loss (8). Thus, identifying a way of inhibiting oxidative stress-induced apoptosis in RGCs may provide an effective therapy for glaucoma.

Oxidative stress is able to destroy mitochondrial membrane potential ( $\Delta\Psi$ m), induce mitochondrial DNA damage and the release of apoptosis-related factors, and thus, trigger apoptosis (9). Mitochondria play an important role in the functioning and survival of RGCs (10,11), and mitochondrial dysfunction has been observed in glaucoma patients (12). Mitochondrial dysfunction is regarded as an early event in the mitochondrial apoptotic pathway. In the mitochondrial apoptotic pathway, mitochondrial dysfunction and the activation of pro-apoptotic

*Correspondence to:* Dr Xinguang Yang, Shaanxi Ophthalmic Medical Center, Xi'an No. 4 Hospital, Affiliated Guangren Hospital, School of Medicine, Xi'an Jiaotong University, 21 Jiefang Road, Xincheng District, Xi'an, Shaanxi 710004, P.R. China  
E-mail: xinguangyangxa@163.com

**Abbreviations:** RGCs, retinal ganglion cells;  $\Delta\Psi$ m, mitochondrial membrane potential; LDH, lactic dehydrogenase; IR, ischemia/reperfusion; ROS, reactive oxygen species; DMEM, Dulbecco's modified Eagle's medium; FBS, fetal bovine serum; OD, optical density

**Key words:** crocin, retinal ganglion cell, hydrogen peroxide, mitochondrial pathway

Bcl-2 family members has been demonstrated to induce the release of cytochrome *c*, which forms the apoptosome complexes, and contributes to the activation of caspase-9 and the cleavage of caspase-3 (8,13,14). The mitochondrial-dependent apoptosis of RGCs has been previously investigated (15). As an important consequence of mitochondrial dysfunction caused by oxidative stress, excessive reactive oxygen species (ROS) are capable of mediating mitochondrial permeability transition and the release of pro-apoptotic proteins, and thus, stimulate the mitochondrial apoptotic pathway (9).

Saffron is a traditional medicine that is frequently used in clinical therapy. The clinical therapeutic effects of saffron have been demonstrated in cancer (16), hypertension (17), insomnia and anxiety (18), cerebral ischemia (19) and depression (20). The effect of saffron on retinal diseases has also been demonstrated by improving focal macular electroretinogram parameters (21), inhibiting cell death induced by intense light (22), and treating macula lutea and ischemic retinopathy caused by old age (23). Crocetin and crocin are the two major active ingredients of saffron. Crocetin can be capable of preventing the retinal damage induced by oxidative and endoplasmic reticulum stresses through inhibition of the activity of caspase-3 and -9 (24), and of protecting the retina from ischemic damage via the inhibition of oxidative stress (25). Crocin has also been shown to exert a protective effect on retinal ischemia/reperfusion (IR) injury-induced apoptosis of RGCs (26). However, the mechanism by which crocin protects against oxidative stress-induced damage to RGCs remains unclear.

In the present study, we investigated the protective effects of crocin on RGCs under oxidative stress. Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) was used to establish a model of oxidative stress injury in RGCs to mimic RGC injury in glaucoma *in vitro*. The anti-apoptotic effect of crocin was determined, and the mitochondrial-mediated apoptosis pathway was examined to determine the anti-apoptotic mechanism of crocin. In addition, the activity of phosphorylated nuclear factor- $\kappa$ B (p-NF- $\kappa$ B) p65 was also measured using western blot analysis.

## Materials and methods

**Cell culture.** RGC-5 cells, obtained from the American Type Culture Collection (Cat. no. PT6600; ATCC, Manassas, VA, USA), were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) (both from Gibco, Carlsbad, CA, USA), 100 U/ml penicillin and 100  $\mu$ g/ml streptomycin (both from Sigma-Aldrich, St. Louis, MO, USA). Cells were grown in a humidified incubator with 5% CO<sub>2</sub> at 37°C and passaged every three days. Second generation RGCs were used in our experiments.

**Establishment of a model of oxidative stress injury in RGCs and crocin treatment.** The cells were equally divided into five groups, which were treated with different concentrations of H<sub>2</sub>O<sub>2</sub> (0, 200, 400, 800 and 1,000  $\mu$ M) for 16 h. Cell viability and lactic dehydrogenase (LDH) release were tested to investigate the cell injury induced by H<sub>2</sub>O<sub>2</sub> in RGCs, and an appropriate concentration was chosen to establish the model of oxidative stress injury in RGCs.

RGC-5 cells were pre-treated with 0.1 and 1  $\mu$ M of crocin (Sigma-Aldrich) for 24 h, and no drugs were added to the control

group. The cells were then subjected to oxidative insult with H<sub>2</sub>O<sub>2</sub> (800  $\mu$ M) for 16 h and collected for subsequent experiments.

**WST-1 cell proliferation assay.** Cell viability was determined using a WST-1 assay (Roche Diagnostics GmbH, Mannheim, Germany). Briefly, the cells were cultured as described above. WST-1 reagent (10  $\mu$ l) was then added to each well, and incubated for 4 h at 37°C. The optical density (OD) was read at 440 nm using a microplate reader (FLUOstar Omega, BMG Labtech, Ortenberg, Germany).

**LDH release assay.** LDH release was determined using a LDH cytotoxicity detection kit (Takara Bio, Tokyo, Japan). Briefly, the cells were cultured in 96-well plates and 100  $\mu$ l cell suspensions of RGCs were collected to assess the LDH activity. Fresh reaction mixture (100  $\mu$ l) was then added to each well and incubated at room temperature for 30 min. The absorbance was determined at 490 nm using a microplate reader (Bio-Rad, Hercules, CA, USA).

**Annexin V/FITC assay.** Apoptotic cells were quantified using a FITC Annexin V apoptosis detection kit (BD Biosciences, Piscataway, NJ, USA). Approximately 1x10<sup>5</sup> cells were collected and resuspended with 100  $\mu$ l of binding buffer. Then, 5  $\mu$ l of FITC Annexin V and 5  $\mu$ l of propidium iodide (PI) were added to stain cells for 15 min at room temperature in the dark. Subsequently, 400  $\mu$ l of 1X binding buffer was added prior to analysis by flow cytometry (BD Biosciences).

**ROS assay.** The level of intracellular ROS was evaluated using a ROS assay kit (Beyotime Biotech, Jiangsu, China). Briefly, the cells were harvested and washed by 1X buffer, and then stained with 20  $\mu$ M of dichloro-dihydro-fluorescein diacetate (DCFH-DA; 1:1,000) for 20 min at 37°C. The signal was read at excitation/emission (Ex/Em) wavelengths of 488/525 nm filter after the cells were further washed three times using PBS. Cells treated with ROSup (provided with the ROS assay kit) only were used as negative controls.

**Measurement of  $\Delta\Psi_m$ .** JC-1 fluorescent dye 9 (Beyotime Biotech) was used to measure  $\Delta\Psi_m$ . The cells were collected and incubated with JC-1 staining solution at 37°C for 15 min in a 5% CO<sub>2</sub> incubator, and then resuspended with 500  $\mu$ l of preheated incubation buffer. The green fluorescence (JC-1-monomer) was viewed at Ex/Em 490/530 nm, and the red fluorescence (JC-1-aggregate) was viewed at Ex/Em wavelengths of 525/590 nm.

**Caspase-3 activity assay.** The enzymatic activity of caspase-3 was detected by a caspase-3 assay kit (Abcam, Cambridge, MA, USA). Briefly, the cells were suspended in lysis buffer and incubated on ice for 10 min. Reaction buffer and DEVD-AFC substrate were then added prior to being read at Ex/Em wavelengths of 400/505 nm.

**Western blot analysis.** The total protein was extracted from RGC-5 cells using RIPA (Beyotime Biotech) and its concentration was determined using a bicinchoninic acid (BCA) assay. The proteins were electrophoresed on 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred onto nitrocellulose membranes (Amersham;




Figure 1. Effects of H<sub>2</sub>O<sub>2</sub> on cell viability and lactic dehydrogenase (LDH) release in (RGC)-5 cells. RGC-5 cells were pre-treated with 0.1 and 1 μM of crocin for 24 h prior to being exposed to 800 μM of H<sub>2</sub>O<sub>2</sub>. (A) Cell viability was determined by a WST-1 assay. (B) LDH release was analyzed by an LDH cytotoxicity assay kit. The columns indicate the means ± SEM. \*P<0.05 vs. the control group.




Figure 2. Effects of crocin on the release of lactic dehydrogenase (LDH) and cell viability in RGC-5 cells. RGC-5 cells were pre-treated with 0.1 and 1 μM of crocin for 24 h prior to being exposed to 800 μM of H<sub>2</sub>O<sub>2</sub>. (A) LDH release was analyzed by an LDH cytotoxicity assay kit. (B) Cell viability was determined by a WST-1 assay. The column indicates the means ± SEM. \*P<0.05 vs. the control group.

GE Healthcare Europe GmbH, Freiburg, Germany). The membranes were blocked with 5% (v/v) dried milk and probed with anti-Bax, anti-Bcl-2, anti-cytochrome *c* (Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA) and anti p-NF-κB p65 (Cell Signaling Technology, Inc., Danvers, MA, USA) at 4°C overnight. Subsequently, HRP-conjugated goat anti-mouse IgG (Bioworld Technology, Inc., St. Louis Park, MN, USA) was added and incubated with the membranes for 1 h at room temperature. β-actin (Cell Signaling Technology, Inc.) was used as the reference protein.

**Statistical analysis.** Data are presented as the means ± SEM. Statistical comparisons were performed using the Student's t-test. P<0.05 was considered to indicate a statistically significant difference.

## Results

**Effects of different concentrations of H<sub>2</sub>O<sub>2</sub> on cell viability and LDH release in RGC-5 cells.** To establish a model of oxidative stress injury in RGCs, different concentrations of H<sub>2</sub>O<sub>2</sub> were used to evaluate the cytotoxicity to select an appropriate concentration. The results showed that H<sub>2</sub>O<sub>2</sub> is capable of decreasing RGC-5 cell viability in a dose-dependent manner; however, there was no significant difference between concentrations of 800 and 1,000 μM (Fig. 1A). As the concentration of H<sub>2</sub>O<sub>2</sub> increased, there was a corresponding gradual increase

in the release of LDH (Fig. 1B). The LDH release assay and the WST-1 assay showed a very low gradient from 800 to 1,000 μM. Thus, an H<sub>2</sub>O<sub>2</sub> concentration of 800 μM was used to establish the a model of oxidative stress injury in RGCs.

**Effects of crocin on cell viability and LDH release in oxidative stress-injured RGC-5 cells.** LDH is a stable cytoplasmic enzyme that is present in all cells. When the plasma membrane is damaged, LDH is rapidly released into the culture supernatant, thus, LDH release in the culture supernatant is a measure of cytotoxicity. To evaluate the anti-cytotoxic effect of crocin on RGC-5 cells, an LDH assay was performed using an LDH cytotoxicity assay kit. LDH release in RGC-5 cells was significantly decreased in the presence of crocin. There was no significant difference in the LDH release between crocin concentrations of 0.1 and 1 μM (P>0.05) (Fig. 2A). To understand the cytoprotective effects of crocin in H<sub>2</sub>O<sub>2</sub>-insulted RGC-5 cells, a WST-1 assay was performed to determine cell viability. Crocin significantly enhanced RGC-5 cell viability in H<sub>2</sub>O<sub>2</sub>-insulted cells (P<0.05), and this effect was not concentration-dependent (P>0.05) (Fig. 2B). These results indicated that crocin could enhance the cell viability of RGC-5 cells that have been injured by H<sub>2</sub>O<sub>2</sub>.

**Effect of crocin on the apoptosis of oxidative stress-injured RGC-5 cells.** H<sub>2</sub>O<sub>2</sub>-induced apoptosis of RGC-5 cells was analyzed using flow cytometry through Annexin V-FITC/PI




Figure 3. Crocin inhibits H<sub>2</sub>O<sub>2</sub>-induced apoptosis of RGC-5 cells. RGC-5 cells were pre-treated with 0.1 and 1 μM of crocin for 24 h prior to being exposed to 800 μM H<sub>2</sub>O<sub>2</sub>. Apoptotic cells were detected by a FITC Annexin V apoptosis detection kit. The columns indicate the means ± SEM. \*P<0.05 vs. the control group.

staining. The percentages of apoptotic cells decreased from 45.39% without crocin, to 32.15% in the presence of 0.1 μM and 28.16% in the presence of 1 μM crocin (Fig. 3). Crocin significantly inhibited H<sub>2</sub>O<sub>2</sub>-induced apoptosis in RGC-5 cells and there was no significant difference in the percentages of apoptotic cells between crocin concentrations of 0.1 and 1 μM.

*Effect of crocin on the production of ROS in oxidative stress-injured RGC-5 cells.* Intracellular ROS is an oxidative stress indicator in cells that plays an important role in apoptosis induction under physiological and pathological conditions (27). ROS are released from the mitochondria, and excessive ROS are able to disrupt the ΔΨ<sub>m</sub> in return (27). In the present study, excessive ROS was generated in H<sub>2</sub>O<sub>2</sub>-injured cells (Fig. 4). With the addition of crocin, the intracellular ROS content was markedly reduced compared with the control group, and the difference between crocin concentrations of 0.1 and 1 μM was not significant. These results indicated that crocin exerted an antioxidant effect on the oxidative stress-injured RGC-5 cells.

*Effect of crocin on ΔΨ<sub>m</sub> in oxidative stress-injured RGC-5 cells.* Mitochondria are closely associated with cell apoptosis, and a decrease in ΔΨ<sub>m</sub> is considered as one of the earliest hallmark events in the cascade reaction process of apoptosis (28). To examine the effect of crocin on H<sub>2</sub>O<sub>2</sub>-induced ΔΨ<sub>m</sub> disruption, the lipophilic cation JC-1 was used to evaluate ΔΨ<sub>m</sub>. Crocin significantly increased ΔΨ<sub>m</sub> in oxidative stress-injured RGC-5 cells (P<0.05). A significant difference between crocin concentrations of 0.1 and 1 μM was identified, which suggested that the mitochondria-dependent pathway may be involved in the protective effect of crocin on H<sub>2</sub>O<sub>2</sub>-injured cells (Fig. 5).

*Effects of crocin on the activity of caspase-3 and the expression of Bcl-2, Bax and cytochrome c in oxidative stress-injured RGC-5 cells.* Caspase-3 is the final effector in the mitochondria-mediated apoptotic pathway (9). Cytochrome *c*, the anti-apoptotic factor Bcl-2 and pro-apoptotic factor Bax are the key regulating factors in the mitochondrial pathway (14). To investigate the effect of crocin on the apoptosis of RGC-5 cells, we detected the expression of Bcl-2, Bax and cytochrome *c* using western blot analysis. As shown in Fig. 6A, the expression level of Bcl-2 was markedly higher in the crocin groups than the control group (P<0.05), and the difference between 0.1 and 1 μM crocin was not significant (Fig. 6B). There was also a significant difference in the expression of Bax between the crocin groups and the control group (P<0.05) (Fig. 6C), while no significant difference was observed between 0.1 and 1 μM crocin. Cytochrome *c* release in oxidative-stress-injured RGC-5 cells was significantly suppressed at crocin concentrations of 0.1 and 1 μM (Fig. 6D), and the inhibitory effect of 1 μM of crocin was significantly stronger than that of 0.1 μM of crocin.

Caspases are aspartic acid proteases containing cysteine, which selectively cleave the target protein of aspartate residue, and thus, induce cell apoptosis. In caspase-dependent signaling, caspase-3 is one of the most important effector caspases, and its activation is the final step of apoptosis (29). To investigate the effect of crocin on H<sub>2</sub>O<sub>2</sub>-induced activation of caspase-3, we used a caspase-3 assay kit to detect caspase-3 activity. The results showed that crocin treatment significantly inhibited the activation of caspase-3 activity, while there was no significant difference between 0.1 and 1 μM crocin (Fig. 6E).




Figure 4. Crocin reduces intracellular reactive oxygen species (ROS) production. RGC-5 cells were pre-treated with 0.1 and 1 μM crocin for 24 h prior to being exposed to 800 μM H<sub>2</sub>O<sub>2</sub>. The level of intracellular ROS was evaluated using flow cytometry. The columns indicate the means ± SEM. \*P<0.05 vs. the control group.




Figure 5. Crocin increases mitochondrial membrane potential (ΔΨm) in oxidative-stress-injured retinal ganglion cells (RGCs). RGC-5 cells were pre-treated with 0.1 and 1 μM crocin for 24 h before being exposed to 800 μM H<sub>2</sub>O<sub>2</sub>. A mitochondrial membrane potential assay kit was used to measure ΔΨm. The columns indicate means ± SEM. \*P<0.05 vs. control group; #P<0.05 vs. 0.1 μM crocin.

Effect of crocin on p-NF-κB 65 in oxidative stress-injured RGC-5 cells. NF-κB is a family of nuclear transcription

factors that include the subunits Rel, p65, RelB, p50 and p52, which influence cell apoptosis by regulating the expression of




Figure 6. Effects of crocin on the activity of caspase-3 and the expression of Bcl-2, Bax and cytochrome *c* in oxidative stress-injured RGC-5 cells. RGC-5 cells were pre-treated with 0.1 and 1  $\mu$ M crocin for 24 h prior to being exposed to 800  $\mu$ M H<sub>2</sub>O<sub>2</sub>. (A) Representative western blot image shows the expression of Bcl-2, Bax and cytochrome *c* in all the groups, and the columns show the quantification of (B) Bcl-2, (C) Bax and (D) cytochrome *c*. (E) The enzymatic activity of caspase-3 was detected by a caspase-3 assay kit. The columns indicate the means  $\pm$  SEM. \* $P$ <0.05 vs. control group; # $P$ <0.05 vs. 0.1  $\mu$ M crocin.




Figure 7. Crocin increases the level of p-nuclear factor- $\kappa$ B (NF- $\kappa$ B) p65 protein in oxidative stress-injured retinal ganglion cells (RGCs). RGC-5 cells were pre-treated with 0.1 and 1  $\mu$ M crocin for 24 h prior to being exposed to 800  $\mu$ M H<sub>2</sub>O<sub>2</sub>. Western blot analysis was used to measure the level of p-NF- $\kappa$ B p65 protein. The columns indicate the means  $\pm$  SEM. \* $P$ <0.05 vs. control group.

cell survival genes (30). The phosphorylation of the transactivation p65 subunit is essential for efficient transcriptional activation by NF- $\kappa$ B (31). Thus, the level of p-NF- $\kappa$ B p65 was measured using western blot analysis in the present experiment. The results showed that the relative p-NF- $\kappa$ B p65 level was markedly increased in the presence of crocin (Fig. 7), which indicates that crocin upregulated the activity of NF- $\kappa$ B in oxidative stress-injured RGC-5 cells.

## Discussion

In glaucoma, elevated intraocular pressure is the most significant risk factor for accelerated RGC death. It is widely accepted that oxidative damage in response to pressure elevation is an important underlying mechanism of elevated

intraocular pressure-induced cell damage and neuronal death (32,33). Thus, H<sub>2</sub>O<sub>2</sub> was used to establish a model of oxidative stress injury in RGCs to mimic RGC injury in glaucoma *in vitro*. The LDH and WST-1 assay results showed that H<sub>2</sub>O<sub>2</sub> decreased cell viability and increased LDH release. Efficiency was highest at a concentration of 800  $\mu$ M, therefore a concentration of 800  $\mu$ M H<sub>2</sub>O<sub>2</sub> was used to establish the model of oxidative stress injured in RGCs for subsequent experiments.

Crocin is one of the active ingredients of saffron, which is frequently used as a traditional medicine for its antitoxic properties (34). The anti-apoptotic and antioxidant effects of crocin have been stated in several studies (35,36). Qi *et al.* (26) have reported that crocin injections prevented apoptosis of RGCs subsequent to retinal IR injury. In the present study, we

detected changes in the cell viability and apoptosis of H<sub>2</sub>O<sub>2</sub>-insulted RGC-5 cells by WST-1 and Annexin V/PI staining *in vitro*, respectively. Our results were consistent with those of Mehri et al (37), who reported that crocin enhanced cell viability and reduced apoptosis. In addition, we also detected LDH release in H<sub>2</sub>O<sub>2</sub>-insulted RGCs using an LDH cytotoxicity assay kit. LDH release was significantly decreased by crocin concentrations of 0.1 and 1 μM. Taken together, these results suggest that crocin prevented H<sub>2</sub>O<sub>2</sub>-induced damage to RGCs.

One of the important mechanisms by which crocin exerts its biological effects is its ability to modulate the redox status of organisms. Evidence has suggested that overproduction of ROS plays an important role in the protective effects of crocin in serum-deprived and hypoxic PC12 cells (38). Mousavi et al (39) have confirmed that crocin decreased the production of ROS induced by glucose in PC12 cells. To determine the effects of crocin on the production of ROS in H<sub>2</sub>O<sub>2</sub>-injured RGC-5 cells, we determined the production of ROS by performing a cellular ROS assay. The results showed that H<sub>2</sub>O<sub>2</sub>-induced production of ROS was significantly suppressed by crocin, suggesting that crocin is capable of reducing the ROS level and suppressing H<sub>2</sub>O<sub>2</sub>-induced oxidative stress in RGC-5 cells.

There are two main pathways of oxidative stress-induced apoptosis: mitochondrial- and death receptor-mediated pathways (40). In the mitochondrial pathway, Bcl-2 and Bax are the key regulators. Bcl-2 inhibits apoptosis by suppressing cytochrome *c* release and caspase activation, while Bax promotes apoptosis by inducing the release of cytochrome *c*, which then triggers the downstream apoptosis event (29). On the other hand, the release of apoptosis-related factor cytochrome *c* may also be inhibited by the rise of the ΔΨ<sub>m</sub>, which decreased intimal permeability (41). Our results show that crocin effectively prevented H<sub>2</sub>O<sub>2</sub>-induced apoptosis by increasing ΔΨ<sub>m</sub>, downregulating Bax and cytochrome *c* and caspase-3, and upregulating Bcl-2. This finding indicates that crocin stabilized the mitochondria and inhibited apoptosis mediated by the mitochondrial pathway, thereby protecting RGCs from apoptosis.

NF-κB activity helps cells to avoid the sustained phase of JNK activation which has been demonstrated to activate the mitochondrial apoptotic pathway (42), and thus, promotes cell survival (43,44). NF-κB plays an important role in the apoptosis of RGCs mediated by H<sub>2</sub>O<sub>2</sub> (45,46). The present study revealed that the level of p-NF-κB p65 was significantly higher in the crocin groups than in the control group. This result suggests that crocin initiated the activation of NF-κB in the presence of H<sub>2</sub>O<sub>2</sub>, thereby reducing H<sub>2</sub>O<sub>2</sub>-induced apoptosis.

Taken together, our results demonstrate that crocin is capable of protecting H<sub>2</sub>O<sub>2</sub>-injured RGC-5 cells from apoptosis through the mitochondrial pathway, and by upregulating the activity of NF-κB.

## Acknowledgements

The present study was supported by the National Natural Science Foundation of China (no. 81273902).

## References

1. Tham YC, Li X, Wong TY, Quigley HA, Aung T and Cheng CY: Global prevalence of glaucoma and projections of glaucoma burden through 2040: a systematic review and meta-analysis. *Ophthalmology* 121: 2081-2090, 2014.
2. Khan AK, Tse DY, van der Heijden ME, Shah P, Nusbaum DM, Yang Z, Wu SM and Frankfort BJ: Prolonged elevation of intraocular pressure results in retinal ganglion cell loss and abnormal retinal function in mice. *Exp Eye Res* 130: 29-37, 2015.
3. Wang Z, Pan X, Wang D, Sun H, Han F, Lv C and Zhang X: Protective effects of protocatechuic acid on retinal ganglion cells from oxidative damage induced by H<sub>2</sub>O<sub>2</sub>. *Neuro Res* 37: 159-166, 2015.
4. Harada T, Harada C, Nakamura K, Quah HM, Okumura A, Namekata K, Saeki T, Aihara M, Yoshida H, Mitani A and Tanaka K: The potential role of glutamate transporters in the pathogenesis of normal tension glaucoma. *J Clin Invest* 117: 1763-1770, 2007.
5. Levkovitch-Verbin H, Vander S, Makarovskiy D and Lavinsky F: Increase in retinal ganglion cells' susceptibility to elevated intraocular pressure and impairment of their endogenous neuroprotective mechanism by age. *Mol Vis* 19: 2011-2022, 2013.
6. Chrysostomou V, Rezanian F, Trounce IA and Crowston JG: Oxidative stress and mitochondrial dysfunction in glaucoma. *Curr Opin Pharmacol* 13: 12-15, 2013.
7. Sancho P, Fernández C, Yuste VJ, Amrán D, Ramos AM, de Blas E, Susin SA and Aller P: Regulation of apoptosis/necrosis execution in cadmium-treated human promonocytic cells under different forms of oxidative stress. *Apoptosis* 11: 673-686, 2006.
8. Almasieh M, Wilson AM, Morquette B, Cueva Vargas JL and Di Polo A: The molecular basis of retinal ganglion cell death in glaucoma. *Prog Retin Eye Res* 31: 152-181, 2012.
9. Circu ML and Aw TY: Reactive oxygen species, cellular redox systems, and apoptosis. *Free Radic Biol Med* 48: 749-762, 2010.
10. Yang XJ, Ge J and Zhuo YH: Role of mitochondria in the pathogenesis and treatment of glaucoma. *Chin Med J (Engl)* 126: 4358-4365, 2013.
11. Lascaratos G, Garway-Heath DF, Willoughby CE, Chau KY and Schapira AH: Mitochondrial dysfunction in glaucoma: Understanding genetic influences. *Mitochondrion* 12: 202-212, 2012.
12. Abu-Amero KK, Morales J and Bosley TM: Mitochondrial abnormalities in patients with primary open-angle glaucoma. *Invest Ophthalmol Vis Sci* 47: 2533-2541, 2006.
13. Brentnall M, Rodriguez-Menocal L, De Guevara RL, Cepero E and Boise LH: Caspase-9, caspase-3 and caspase-7 have distinct roles during intrinsic apoptosis. *BMC Cell Biol* 14: 32, 2013.
14. Estaquier J, Vallette F, Vayssiere JL and Mignotte B: The mitochondrial pathways of apoptosis. *Adv Exp Med Biol* 942: 157-183, 2012.
15. Meng QF, Lv J, Ge H, Zhang L, Xue F, Zhu Y and Liu P: Overexpressed mutant optineurin (E50K) induces retinal ganglion cells apoptosis via mitochondrial pathway. *Mol Biol Rep* 39: 5867-5873, 2012.
16. Abdullaev FI and Espinosa-Aguirre JJ: Biomedical properties of saffron and its potential use in cancer therapy and chemoprevention trials. *Cancer Detect Prev* 28: 426-432, 2004.
17. Imenshahidi M, Hosseinzadeh H and Javadpour Y: Hypotensive effect of aqueous saffron extract (*Crocus sativus* L.) and its constituents, safranal and crocin, in normotensive and hypertensive rats. *Phytother Res* 24: 990-994, 2010.
18. Hosseinzadeh H and Noraei NB: Anxiolytic and hypnotic effect of *Crocus sativus* aqueous extract and its constituents, crocin and safranal, in mice. *Phytother Res* 23: 768-774, 2009.
19. Hosseinzadeh H, Sadeghnia HR, Ghaeni FA, Motamedshariaty VS and Mohajeri SA: Effects of saffron (*Crocus sativus* L.) and its active constituent, crocin, on recognition and spatial memory after chronic cerebral hypoperfusion in rats. *Phytother Res* 26: 381-386, 2012.
20. Akhondzadeh Basti A, Moshiri E, Noorbala AA, Jamshidi AH, Abbasi SH and Akhondzadeh S: Comparison of petal of *Crocus sativus* L. and fluoxetine in the treatment of depressed outpatients: A pilot double-blind randomized trial. *Prog Neuropsychopharmacol Biol Psychiatry* 31: 439-442, 2007.
21. Falsini B, Piccardi M, Minnella A, Savastano C, Capoluongo E, Fadda A, Balestrazzi E, Maccarone R and Bisti S: Influence of saffron supplementation on retinal flicker sensitivity in early age-related macular degeneration. *Invest Ophthalmol Vis Sci* 51: 6118-6124, 2010.
22. Maccarone R, Di Marco S and Bisti S: Saffron supplement maintains morphology and function after exposure to damaging light in mammalian retina. *Invest Ophthalmol Vis Sci* 49: 1254-1261, 2008.
23. Moghaddasi MS: Saffron chemicals and medicine usage. *J Med Plants Res* 4: 427-430, 2010.

24. Yamauchi M, Tsuruma K, Imai S, Nakanishi T, Umigai N, Shimazawa M and Hara H: Crocetin prevents retinal degeneration induced by oxidative and endoplasmic reticulum stresses via inhibition of caspase activity. *Eur J Pharmacol* 650: 110-119, 2011.
25. Ishizuka F, Shimazawa M, Umigai N, Ogishima H, Nakamura S, Tsuruma K and Hara H: Crocetin, a carotenoid derivative, inhibits retinal ischemic damage in mice. *Eur J Pharmacol* 703: 1-10, 2013.
26. Qi Y, Chen L, Zhang L, Liu WB, Chen XY and Yang XG: Crocetin prevents retinal ischaemia/reperfusion injury-induced apoptosis in retinal ganglion cells through the PI3K/AKT signalling pathway. *Exp Eye Res* 107: 44-51, 2013.
27. Simon HU, Haj-Yehia A and Levi-Schaffer F: Role of reactive oxygen species (ROS) in apoptosis induction. *Apoptosis* 5: 415-418, 2000.
28. Mignotte B and Vayssiere JL: Mitochondria and apoptosis. *Eur J Biochem* 252: 1-15, 1998.
29. Granville DJ and Gottlieb RA: Mitochondria: Regulators of cell death and survival. *ScientificWorldJournal* 2: 1569-1578, 2002.
30. Tang G, Minemoto Y, Dibling B, Purcell NH, Li Z, Karin M and Lin A: Inhibition of JNK activation through NF-kappaB target genes. *Nature* 414: 313-317, 2001.
31. Zhong H, Voll RE and Ghosh S: Phosphorylation of NF-kB p65 by PKA stimulates transcriptional activity by promoting a novel bivalent interaction with the coactivator CBP/p300. *Mol Cell* 1: 661-671, 1998.
32. Ju WK, Liu Q, Kim KY, Crowston JG, Lindsey JD, Agarwal N, Ellisman MH, Perkins GA and Weinreb RN: Elevated hydrostatic pressure triggers mitochondrial fission and decreases cellular ATP in differentiated RGC-5 cells. *Invest Ophthalmol Vis Sci* 48: 2145-2151, 2007.
33. Liu Q, Ju WK, Crowston JG, Xie F, Perry G, Smith MA, Lindsey JD and Weinreb RN: Oxidative stress is an early event in hydrostatic pressure induced retinal ganglion cell damage. *Invest Ophthalmol Vis Sci* 48: 4580-4589, 2007.
34. Huynh TP, Mann SN and Mandal NA: Botanical compounds: effects on major eye diseases. *Evid Based Complement Alternat Med* 2013: 549174, 2013.
35. Soeda S, Ochiai T, Paopong L, Tanaka H, Shoyama Y and Shimeno H: Crocetin suppresses tumor necrosis factor-alpha-induced cell death of neuronally differentiated PC-12 cells. *Life Sci* 69: 2887-2898, 2001.
36. Ochiai T, Soeda S, Ohno S, Tanaka H, Shoyama Y and Shimeno H: Crocetin prevents the death of PC-12 cells through sphingomyelinase-ceramide signaling by increasing glutathione synthesis. *Neurochem Int* 44: 321-330, 2004.
37. Mehri S, Abnous K, Mousavi SH, Shariaty VM and Hosseinzadeh H: Neuroprotective effect of crocin on acrylamide-induced cytotoxicity in PC12 cells. *Cell Mol Neurobiol* 32: 227-235, 2012.
38. Ochiai T, Shimeno H, Mishima K, Iwasaki K, Fujiwara M, Tanaka H, Shoyama Y, Toda A, Eyanagi R and Soeda S: Protective effects of carotenoids from saffron on neuronal injury in vitro and in vivo. *Biochim Biophys Acta* 1770: 578-584, 2007.
39. Mousavi SH, Tayarani NZ and Parsaee H: Protective effect of saffron extract and crocin on reactive oxygen species-mediated high glucose-induced toxicity in PC12 cells. *Cell Mol Neurobiol* 30: 185-191, 2010.
40. Hengartner MO: The biochemistry of apoptosis. *Nature* 407: 770-776, 2000.
41. Hao M, Li Y, Lin W, Xu Q, Shao N, Zhang Y and Kuang H: Estrogen prevents high-glucose-induced damage of retinal ganglion cells via mitochondrial pathway. *Graefes Arch Clin Exp Ophthalmol* 253: 83-90, 2015.
42. Weston CR and Davis RJ: The JNK signal transduction pathway. *Curr Opin Cell Biol* 19: 142-149, 2007.
43. Kamata H, Honda S, Maeda S, Chang L, Hirata H and Karin M: Reactive oxygen species promote TNFalpha-induced death and sustained JNK activation by inhibiting MAP kinase phosphatases. *Cell* 120: 649-661, 2005.
44. Nakano H, Nakajima A, Sakon-Komazawa S, Piao JH, Xue X and Okumura K: Reactive oxygen species mediate crosstalk between NF-kappaB and JNK. *Cell Death Differ* 13: 730-737, 2006.
45. Gupta VK, You Y, Li JC, Klistorner A and Graham SL: Protective effects of 7,8-dihydroxyflavone on retinal ganglion and RGC-5 cells against excitotoxic and oxidative stress. *J Mol Neurosci* 49: 96-104, 2013.
46. Ozawa Y, Yuki K, Yamagishi R, Tsubota K and Aihara M: Renin-angiotensin system involvement in the oxidative stress-induced neurodegeneration of cultured retinal ganglion cells. *Jpn J Ophthalmol* 57: 126-132, 2013.