

Spontaneous cervical hematoma caused by giant parathyroid adenoma in an elderly patient: A case report

BELÉN MATÍAS-GARCÍA, FERNANDO MENDOZA-MORENO, MANUEL DÍEZ-ALONSO, ENRIQUE OVEJERO-MERINO, CRISTINA VERA-MANSILLA, ALMA BLÁZQUEZ-MARTÍN, ANA QUIROGA-VALCÁRCEL, RUBÉN JIMÉNEZ-MARTÍN, REBECA D'AMICO, INMACULADA LASA-UNZÚE and ALBERTO GUTIÉRREZ-CALVO

Department of General and Digestive Surgery, Príncipe de Asturias Teaching Hospital, 28805 Madrid, Spain

Received October 5, 2023; Accepted November 21, 2023

DOI: 10.3892/mi.2023.125

Abstract. Spontaneous cervical hematoma usually occurs as a consequence of extracapsular bleeding from a parathyroid gland, generally due to the presence of an adenoma (giant adenoma), glandular hyperplasia, cystic component, or, less frequently, due to the existence of a carcinoma. The hematoma can be confined to the cervical compartment or extend to the mediastinum, potentially causing airway compression. Despite this, the recommended management in hemodynamically stable patients consists of surveillance and hospital monitoring with delayed surgery after a few weeks. On the other hand, in those patients with airway compromise and instability, emergency surgery, consisting of cervicotomy and drainage, is mandatory. The present study describes the case of a 78-year-old patient with a medical history of high blood pressure, non-insulin-dependent diabetes mellitus, dyslipidemia, moderate aortic stenosis, chronic kidney disease and sarcoidosis under pharmacological treatment who attended the emergency department due to symptoms of neck pain, an increase in soft tissue, and dyspnea on moderate exertion with an evolution leading to respiratory failure. This was secondary to a diagnosis of spontaneous cervical hematoma that required urgent surgical intervention. The results of histopathological analysis revealed that a giant parathyroid adenoma was responsible for the bleeding. The patient had a complicated post-operative period with a prolonged admission to the intensive care unit.

Introduction

Primary hyperparathyroidism (PHPT) is the most common pathological condition affecting the parathyroid glands and represents the third most common cause of endocrine disorders worldwide (1,2). It is estimated that the incidence of PHPT is ~2-4 cases per 10,000 inhabitants, with a higher prevalence among post-menopausal women (1). In 80-90% of cases, PHPT is attributed to the presence of a parathyroid adenoma, which autonomously produces parathyroid hormone (PTH), affecting calcium and phosphorus metabolism (3,4).

Spontaneous parathyroid hemorrhage is an exceptionally rare, yet potentially life-threatening condition due to airway compromise, often necessitating urgent surgical intervention. Clinically, it manifests with symptoms, such as cervical pain, dysphagia, dysphonia, or dyspnea, and physical examination may reveal ecchymosis in the cervical and/or thoracic regions (5).

The present study describes the case of a patient with spontaneous hemorrhage from a giant parathyroid adenoma resulting in acute airway compromise requiring surgical drainage.

Case report

A 78-year-old female patient with a medical history of high blood pressure, non-insulin-dependent diabetes mellitus, dyslipidemia, moderate aortic stenosis, chronic kidney disease and sarcoidosis, receiving pharmacological treatment, presented to the Emergency Department of Príncipe de Asturias Teaching Hospital (Madrid, Spain) with complaints of increased cervical swelling, dyspnea during moderate exertion and odynophagia.

Her vital signs upon admission were as follows: Blood pressure, 185/88 mmHg; heart rate, 86 bpm; and baseline oxygen saturation, 96%. A physical examination revealed a cervical soft tissue hematoma without crepitation or fluctuation upon palpation. A cervical ultrasound was performed as part of the imaging assessment, which revealed a markedly enlarged thyroid with multiple nodules and intrathoracic extension, as well as increased subcutaneous tissue thickness with altered echogenicity, without distinct collections. A posteroanterior

Correspondence to: Dr Fernando Mendoza-Moreno, Department of General and Digestive Surgery, Príncipe de Asturias Teaching Hospital, Carretera Alcalá Meco 1, Alcalá de Henares, 28805 Madrid, Spain E-mail: f.mendoza@uah.es

Abbreviations: PHPT, primary hyperparathyroidism; PTH, parathyroid hormone; CT scan, computed tomography scan

Key words: parathyroid adenoma, thyroidectomy, cervical hematoma, primary hyperparathyroidism

chest X-ray displayed a reduction in tracheal caliber in the cervical region related to a mixed multinodular goiter.

Due to the clinical suspicion of spontaneous cervical hematoma and the diagnosis of a hypertensive crisis, the patient remained under hospital observation. With stable clinical condition, blood pressure was managed through intravenous urapidil infusion and arterial phase control. However, during her stay in the emergency room, the patient experienced a desaturation episode with sudden respiratory distress (oxygen saturation decreased to 90%) and dysphonia. In response to this, a cervico-thoracic axial computed tomography (CT) scan with intravenous contrast was performed, revealing a lesion of heterogeneous density ~2x5 cm (anteroposterior x transverse) in the left thyroid lobe, extending to the second thoracic vertebra, with hyperdensity in different phases, suggestive of a hemorrhagic complication. Additionally, an increase in soft tissue density in the parapharyngeal and retropharyngeal space was observed, without signs of active bleeding (Fig. 1).

Given the worsening clinical condition of the patient and the potential for airway compromise, urgent surgery was deemed necessary. Under general anesthesia, following orotracheal intubation with a video laryngoscope, a median cervicotomy was performed. This revealed a large cervical hematoma extending into the mediastinum and parapharyngeal spaces, associated with the thyroid gland, leading to destruction and bleeding in the left thyroid lobe. The hematoma within the thyroid gland was evacuated, and a total thyroidectomy and tracheostomy were performed due to persistent bleeding (Fig. 2).

Following the surgery, the patient was admitted to the intensive care unit, where she experienced a complicated post-operative course marked by the development of pneumonia associated with mechanical ventilation and disruptions in phosphorus-calcium metabolism, including hypocalcemia refractory to replacement therapy and hypothyroidism. Following a 97-day hospital stay, the patient was discharged.

The histopathological examination of the surgical specimen revealed a thyroidectomy specimen weighing 15.8 g with dimensions of 4x4x12.5 cm, including a papillary microcarcinoma of the thyroid (1.5 mm) with clear surgical margins. Furthermore, the specimen contained parathyroid tissue measuring 6x4x3 cm and weighing 20.51 g, consistent with an adenoma and showing the presence of a hemorrhagic focus.

Discussion

The parathyroid glands are small endocrine glands nestled in the neck, typically located behind the larger thyroid gland. While the majority of individuals have four parathyroid glands, the exact number can vary among individuals.

These glands play a pivotal role in regulating calcium levels in both the bloodstream and cells. Their primary function is to produce and secrete PTH, which plays a crucial role in maintaining the calcium and phosphorus levels in the body. They typically measure between 3-5 mm in size and have a weight of ~70 mg (6).

A parathyroid adenoma is a benign (non-cancerous) tumor that can affect one of the parathyroid glands. By contrast, a 'giant' parathyroid adenoma refers to a significantly enlarged parathyroid adenoma. While the majority of parathyroid adenomas are relatively small, with an average weight of <1 g, giant adenomas can be substantially larger, often weighing >3.5 g and measuring >2 cm in length. The largest reported giant parathyroid adenoma, as described in the study by Mahmodlou *et al* (1), measured 9x6x4 cm and weighed 122 g.

Giant parathyroid adenomas are a less common subtype of parathyroid adenomas, although they are still benign (non-cancerous) tumors. These adenomas overproduce PTH, leading to primary hyperparathyroidism, and they represent 5% of all parathyroid adenomas (7).

Spontaneous cervical hematoma is a rare complication of PHPT. The first documented case of spontaneous cervical hematoma due to a parathyroid adenoma dates back to 1934 when Capps (8) described it in the literature. Since then, <60 cases have been reported worldwide (3).

In the absence of cervical trauma, it is hypothesized that an imbalance between the gland's tissue growth and its blood flow may lead to phenomena, such as hemorrhage, infarction and subsequent necrosis, similar to the apoplexy observed in other endocrine neoplasms (9). Unlike the thyroid gland, the capsule surrounding the parathyroid glands is exceptionally thin, which can result in a rapid extension into the cervical and/or mediastinal regions in case of rupture (10,11).

Imaging tests, including CT scans, ultrasonography, or magnetic resonance imaging, are indispensable for diagnosing and pinpointing the location of the hematoma. The differential diagnosis should consider other conditions, such as cervical abscess, superior vena cava syndrome, cervical dissection, or subacute thyroiditis (5,12).

As regards the symptoms, patients with spontaneous cervical hematoma may experience neck swelling, tracheal and esophageal compression, and develop dysphagia, dyspnea and hoarseness. Simcic *et al* (13) proposed a more specific triad of symptoms, namely neck swelling, neck ecchymosis and hypercalcemia. However, elevated calcium levels have been observed in <70% of spontaneous hematoma cases. In the patient described herein, despite her chronic renal failure, neither hypercalcemia nor elevated PTH levels were found.

According to the study by Alegre *et al* (14), the management of patients with spontaneous cervical hematoma should be conservative if there is no airway compromise and hemodynamic instability persists. In such cases, intervention is typically delayed, involving procedures, such as thyroidectomy or parathyroidectomy, typically performed within ~ 6 weeks. This conservative approach has a success rate of 51.4% and is particularly suitable for elderly patients with high morbidity (3). In the case described in the present study, given the absence of hemodynamic instability, respiratory compromise and the notable morbidity of the patient, a non-operative approach involving monitoring and blood pressure control was initially favored. However, as the respiratory condition of the patient deteriorated, surgical intervention became necessary.

In conclusion, in the face of adversity, spontaneous cervical hematoma emerges as a rare yet formidable challenge, often originating from a distressed parathyroid gland (be it an adenoma, hyperplasia, cyst, or parathyroid carcinoma). It serves as a poignant reminder that within life's unpredictable narrative, the unexpected can occur.

The occurrence of a non-traumatic cervical hemorrhage, poses as a reminder of the resilience that resides within





Figure 1. (A) Chest X-ray illustrating the tapering of the trachea (red arrow). (B-D) Cervical hematoma with progressive airway compression.



Figure 2. (A) Cervical spontaneous hematoma. (B) Cervicotomy surgical field following thyroidectomy and drainage of the cervical hematoma.

humans. The path to healing may take various forms, with options ranging from a prudent, watchful eye to timely surgical intervention. In this journey, the path chosen is a reflection of the patient's clinical stability, underscoring the importance of individualized care.

Emphasizing the importance of reminding clinical staff to be vigilant in recognizing cervical hematoma induced by parathyroid adenoma as a life-threatening condition cannot be overstated. While it is a rare occurrence, the consequences of overlooking this condition can be severe. Cervical hematoma in the context of parathyroid adenoma can lead to airway obstruction and respiratory compromise, rendering it an urgent medical emergency. This life-threatening situation underscores the critical need for early detection and immediate intervention. By raising awareness among clinical staff, healthcare professionals can ensure that timely assessments and treatments are administered, potentially saving lives in situations where every moment counts. Moreover, such reminders can help prevent misdiagnosis or delayed intervention, reducing the risk of adverse outcomes for patients with this uncommon, yet 'high-stakes' medical condition.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

All authors (FMM, BMG, MDA, EOM, CVM, ABM, AQV, RJM, RDA, ILU and AGC) contributed to the diagnosis and treatment of the patient, and in the design of the study. FMM was a major contributor to the writing of the manuscript. BMG and FMM confirm the authenticity of all the raw data. All authors have read and approved the final manuscript.

Ethics approval and consent to participate

The present study followed international and national regulations and was in agreement with the Declaration of Helsinki, and ethical principles. The patient signed an informed consent form prior to the performance of the surgery.

Patient consent for publication

The patient provided written informed consent for the publication of any data and/or accompanying images, before the surgery was performed. Patients have a right to anonymity and privacy, and authors have a legal and ethical responsibility to respect this right.

Competing interests

The authors declare that they have no competing interests.

References

- Mahmodlou R, Sedokani A, Pezeshk A and Najafinejad B: Giant parathyroid adenoma: A case report. J Med Case Rep 16: 150, 2022.
- Spanheimer PM, Stoltze AJ, Howe JR, Sugg SL, Lal G and Weigel RJ: Do giant parathyroid adenomas represent a distinct clinical entity? Surgery 154: 714-719, 2013.
- Tessler I, Adi M, Diment J, Lahav Y, Halperin D and Cohen O: Spontaneous neck hematoma secondary to parathyroid adenoma: A case series. Eur Arch Otorhinolaryngol 277: 2551-2558, 2020.
- 4. Khan S, Choe CC, Shabaik A and Bouvet M: Parathyroid adenoma presenting with spontaneous cervical and anterior mediastinal hemorrhage: A case report. Medicine (Baltimore) 98: e14347, 2019.
- Familiari M, Di Santo D, Galli A, Danè G, Giordano L, Mellone R and Bussi M: Spontaneous extracapsular parathyroid adenoma hemorrhage: When surgery is required? Endocrine 75: 575-582, 2022.
- 6. Jayant M, Kaushik R, Kochar S and Attri AK: Giant parathyroid adenoma. Kathmandu Univ Med J (KUMJ) 9: 77-79, 2011.
- Rutledge S, Harrison M, O'Connell M, O'Dwyer T and Byrne MM: Acute presentation of a giant intrathyroidal parathyroid adenoma: A case report. J Med Case Rep 10: 286, 2016.
- Capps RB: Multiple parathyroid tumors with massive mediastinal and subcutaneous hemorrhage. Am J Med Sci 188: 801-804, 1934.
- 9. Howard J, Follis RH, Yendt ER and Connor TB: Hyperparathyroidism; case report illustrating spontaneous remission due to necrosis of adenoma, and a study of the incidence of necrosis in parathyroid adenomas. J Clin Endocrinol 13: 997-1008, 1953.
- Osorio Silla I, Lorente L, Sancho JJ and Sitges-Serra A: Spontaneous neck hematoma due to rupture of parathyroid adenoma: Report of 3 cases and literature review. Endocrinol Nutr 61: e5-e6, 2014.
- Yoshimura N, Mukaida H, Mimura T, Iwata K, Amioka A, Hirabayashi N and Takiyama W: A case of an acute cervicomediastinal hematoma secondary to the spontaneous rupture of a parathyroid adenoma. Ann Thorac Cardiovasc Surg 20: 816-820, 2014.
- Aggarwal V, Mishra A, Bhargav PR and Ramakant P: Giant parathyroid adenoma. ANZ J Surg 79: 91, 2009.
- Simcic KJ, McDermott MT, Crawford GJ, Marx WH, Ownbey JL and Kidd GS: Massive extracapsular hemorrhage from a parathyroid cyst. Arch Surg 124: 1347-1350, 1989.
- Alegre B, Rodríguez-VanStrahlen C and Vilaseca I: Spontaneous haemorrhage from a parathyroid adenoma with airway involvement. Med Clin (Barc) 161: 273-274, 2023.



Copyright © 2023 Matías-García et al. This work is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) License.