

# Beneficial effects of protease preparations derived from *Aspergillus* on the colonic luminal environment in rats consuming a high-fat diet

YONGSHOU YANG<sup>1</sup>, NOVITA VIVI SITANGGANG<sup>1</sup>, NORIHISA KATO<sup>1</sup>, JUNJI INOUE<sup>2</sup>,  
TAKAYUKI MURAKAMI<sup>2</sup>, TOSHIRO WATANABE<sup>3</sup>, TAKAFUMI IGUCHI<sup>3</sup> and YUKAKO OKAZAKI<sup>4</sup>

<sup>1</sup>Graduate School of Biosphere Science, Hiroshima University, Higashi-Hiroshima 739-8528; <sup>2</sup>Ahjikan Co., Ltd., Hiroshima 733-8677; <sup>3</sup>Technology Development Laboratory, Yaegaki Bio-Industry Incorporated, Himeji 679-4298;

<sup>4</sup>Faculty of Human Life Sciences, Fuji Women's University, Ishikari, Hokkaido 061-3204, Japan

Received April 16, 2015; Accepted July 3, 2015

DOI: 10.3892/br.2015.490

**Abstract.** The present study investigated the effects of the dietary addition of the protease preparations derived from *Aspergillus* on the colonic luminal environment. Rats were fed a 30% beef tallow diet with or without the protease preparations, including Amano protease (protease A 'Amano SD', neutral proteases from *Aspergillus* spp.) or orientase (orientase AY, acid proteases from *Aspergillus niger*) at the dose of 0.2% for 3 weeks. Cecal *Bifidobacterium* was significantly elevated in the dietary Amano protease group (194-fold,  $P < 0.05$ ), but not in the orientase group. *Lactobacillus* was elevated in the two groups ( $P < 0.05$ ). Cecal *n*-butyrate, propionate and lactate were higher in the Amano protease and orientase groups compared with the controls ( $P < 0.05$ ). Fecal immunoglobulin A and mucins were elevated in the Amano protease group ( $P < 0.05$ ). These results suggest the potential effect of the consumption of *Aspergillus*-derived protease preparations that are favorable for the colonic luminal environment in rats fed a high-fat diet.

## Introduction

Prebiotics, such as dietary fiber and oligosaccharides, contribute to the improvement of human health. Numerous attempts to promote health by prebiotics aim to selectively increase colonic bacterial groups, mainly *Bifidobacterium*. Increasing evidence indicates that *Bifidobacterium* improve the colonic luminal environment. Elevated levels of colonic *Bifidobacterium* are associated with increased colonic production of organic acids,

such as butyrate and propionate (1); butyrate is an important energy source in the colonic epithelium that regulates cell growth and differentiation in healthy colon cells, and increases mucin production, which contributes to intestinal barrier function and induces apoptosis in tumor cells (2,3). Furthermore, the administration of certain *Bifidobacterium* species, such as *Bifidobacterium bifidum* (*B. bifidum*), can enhance intestinal immunoglobulin A (IgA), which is an indicator of intestinal immune function (4).

Our previous study developed a method for producing *Aspergillus awamori*-fermented burdock. Burdock is traditionally consumed as a root vegetable or herbal medicine in Asia, Europe and North America. It is rich in dietary fibers and has been studied extensively due to its prebiotic effects (5). A high-fat (HF) diet decreases the levels of colonic probiotics and organic acids, and it is believed to elevate the risk of colon cancer (6). Notably, our previous study reported that the consumption of dietary *Aspergillus awamori*-fermented burdock markedly increased the levels of cecal *Bifidobacterium* and organic acids, including lactate, propionate, acetate and butyrate, and fecal IgA and mucins (index of colonic barrier function) compared to burdock powder in rats fed an HF diet (7). Our previous study indicated that consumption of the water-soluble fraction of *Aspergillus awamori*-fermented burdock markedly increases cecal *Bifidobacterium* levels (unpublished data); this fraction of *Aspergillus awamori*-fermented burdock contains extracellular neutral and acid proteases of *Aspergillus awamori*. Accordingly, we hypothesize that the beneficial effect of *Aspergillus awamori*-fermented burdock on colonic health is derived from the proteases of *Aspergillus per se*. This hypothesis will aid in the clarification of the possible mechanism by which *Aspergillus awamori*-fermented burdock increased the cecal *Bifidobacterium* and organic acids, and fecal IgA and mucins. Our preliminary study investigated the effect of the consumption of several food-processing proteases derived from *Aspergillus* on colonic luminal microflora. Among them, two enzyme preparations, protease A 'Amano SD' and orientase AY, were notably identified to cause a marked increase in the colonic levels of *Bifidobacterium* and

---

Correspondence to: Dr Yukako Okazaki, Faculty of Human Life Sciences, Fuji Women's University, 4-5 Hanakawa Minami, Ishikari, Hokkaido 061-3204, Japan  
E-mail: yokazaki@fujijoshi.ac.jp

**Abbreviations:** FIAF, fasting-induced adipose factor; HF, high-fat

**Key words:** *Aspergillus*-derived proteases, intestinal microflora, organic acids, immunoglobulin A, mucins

*Lactobacillus* in rats fed an HF diet. Accordingly, the present study analyzed the effects of these enzyme preparations on colonic variables, including microflora, fermentation, IgA, mucins and gene expression in rats fed an HF diet.

## Materials and methods

**Animals and diets.** A total of 27 male Sprague-Dawley rats (4-week-old) were purchased from the Hiroshima Laboratory Animal Center (Hiroshima, Japan) and were maintained according to the 'Guide for the Care and Use of Laboratory Animals' established by Hiroshima University. The study was approved by the Ethics Committee of Hiroshima University. The rats were housed individually in an air-conditioned room at 23–24°C under a 12-h light/dark cycle (lights on from 08:00 a.m. to 20:00 p.m.). Following acclimatization with a non-purified commercial rodent diet (moderate fat diet; Oriental Yeast Co., Tokyo, Japan) for 7 days, the rats (mean body weight, 108 g) were divided into 3 groups with 9 rats in each.

The composition of the experimental diets (Table I) was based on an HF diet (30% beef tallow). The group of rats were randomly assigned to 1 of 3 diets: A control diet and experimental diets containing protease A 'Amano SD' [neutral proteases and peptidase from *Aspergillus* spp., containing 70% (w/w) dextrin, protease activity at pH 6.0; 50,000 U/g; Amano Enzyme Co., Ltd., Nagoya, Japan], or orientase AY [acid proteases from *Aspergillus niger*, containing 20% (w/w) dextrin, protease activity at pH 4.0; 200,000 U/g; HBI Enzymes Inc., Shisou, Japan]. Amano protease (protease A 'Amano SD') or orientase (orientase AY) was added to the experimental diet at 0.2% (w/w) (Table I). Equal amounts of the experimental diets were incorporated daily into food cups at 19:00 p.m. (9, 10, 12, 14 and 15 g on days 1, 2–4, 5–7, 8–13 and 14–21, respectively) to prevent differences in food intake. All the food was consumed each day until the food was served on the following day. The weight of any spilled food was recorded daily and was accounted for in the calculation of food intake. Feces were collected during the last 3 days, stored at -30°C, freeze-dried and milled. At the end of the feeding period, the rats were sacrificed by decapitation under diethyl ether anesthesia. Blood was collected and serum was separated by centrifugation at 2,000 x g for 20 min and stored at -80°C. The cecum was excised and its contents were immediately collected, weighed and stored at -80°C until subsequent analysis. The colonic mucosa was scraped with a sterilized glass slide and used for RNA extraction.

**Measurements.** Bacterial genomic DNA was isolated from the cecal digesta using the UltraClean™ Fecal DNA extraction kit (MO BIO Laboratories, Inc., Carlsbad, CA, USA) according to the manufacturer's instructions. Bacterial groups were quantified by quantitative polymerase chain reaction (qPCR) using a LightCycler 480 System II (Roche Applied Science, Indianapolis, IN, USA). The group-specific primers for qPCR are shown in Table II. qPCR was performed in a reaction volume of 20 µl containing 10 µl SYBR qPCR mix (Toyobo Co., Ltd., Osaka, Japan), 200 nM each of the forward and reverse primers (8–13), and 2 µl cecal DNA samples. The reaction conditions were 95°C for 30 sec followed by 40 cycles

Table I. Composition of experimental diets.

Characteristics	Control (%, w/w)	Amano protease (%, w/w)	Orientase (%, w/w)
Beef tallow	30.00	30.00	30.00
Casein <sup>a</sup>	20.00	20.00	20.00
L-cystine	0.30	0.30	0.30
Vitamin mixture <sup>b</sup>	1.00	1.00	1.00
Mineral mixture <sup>b</sup>	3.50	3.50	3.50
Cellulose	5.00	5.00	5.00
Sucrose	20.00	20.00	20.00
Corn starch	20.20	19.53	19.95
Amano protease <sup>c</sup>		0.67	
Orientase <sup>d</sup>			0.25

<sup>a</sup>Casein: Net protein content, 87% (w/w). <sup>b</sup>American Institute for Nutrition (AIN-93). <sup>c</sup>Protease A 'Amano SD': This powder contains 70% (w/w) dextrin. <sup>d</sup>Orientase AY: This powder contains 20% (w/w) dextrin.

at 95°C for 5 sec, 55°C for 15 sec and 72°C for 30 sec. The fluorescent products were detected at the last step of each cycle. Melting curve analysis was performed following amplification to distinguish the targeted PCR product from the non-targeted PCR product. Data were analyzed by the second derivative maximum method of the LightCycler 480 Basic Software. The relative abundances of the microbial populations are expressed as the proportions of total bacterial 16S rDNA gene.

The pH of cecal digesta was measured directly by a compact pH meter (B-212; Horiba, Ltd., Kyoto, Japan). Cecal organic acids were measured according to the internal standard method using high-performance liquid chromatography (HPLC) (L-2130; Hitachi, Tokyo, Japan) equipped with an Aminex HPX-87H ion exclusion column (7.8 mm i.d. x 30 cm; Bio-Rad, Richmond, CA, USA) (14). Briefly, 500 mg of cecal digesta was homogenized in 5 ml 50 mmol/l H<sub>2</sub>SO<sub>4</sub> containing 10 mmol/l 2,2-dimethyl butyric acid as an internal standard and subsequently centrifuged at 17,000 x g at 2°C for 20 min. The supernatant was ultrafiltered and the filtrate was applied to the HPLC. Cecal ammonia levels were measured according to the method of Lin and Visek (15).

Total IgA concentrations in feces were measured by using an ELISA quantitation kit (Bethyl Laboratories, Montgomery, TX, USA). Mucins were extracted as described by Bovee-Oudenhoven *et al* (16) and quantitated by a fluorometric assay (17).

Total RNA was extracted from colonic mucus using QIAzol lysis reagent (Qiagen, Hilden, Germany) according to the manufacturer's instructions. Isolated RNA was purified using the RNeasy® Lipid Tissue Mini kit (Qiagen). cDNA was synthesized from total RNA using the Revertrace reverse transcription-PCR kit (Toyobo Co., Ltd.). qPCR was performed on an Opticon 2 system (Bio-Rad) using the SYBR qPCR mix (Toyobo Co., Ltd.) and the following primers: Fasting-induced adipose factor (FIAF) forward, GACTGCCAGGAAGCTTTCA and reverse, TGTGATGCTGTGCATCTTCT;

Table II. Target bacteria group, primers sequence and product size for quantitative PCR.

Target bacteria group	Sequence (5'-3')	Product size, bp	Reference
Total bacteria	F: ACTCCTACGGGAGGCAG R: GTATTACCGCGGCTGCTG	200	(8)
<i>Bifidobacterium</i> spp.	F: CGCGTCYGGTGTGAAAG R: CCCACATCCAGCATCCA	244	(9)
<i>Lactobacillus</i> spp.	F: GAGGCAGCAGTAGGGAATCTTC R: GGCCAGTTACTACCTCTATCCTTCTTC	126	(9)
<i>Clostridium coccooides</i> group	F: AAATGACGGTACCTGACTAA R: CTTTGAGTTTCATTCTTGCGAA	440	(10)
<i>Clostridium leptum</i> subgroup	F: GCACAAGCAGTGGAGT R: CTCCTCCGTTTTGTCAA	239	(10)
<i>Akkermansia muciniphila</i>	F: CAGCACGTGAAGGTGGGGAC R: CCTTGCGGTTGGCTTCAGAT	327	(11)
<i>Enterobacteriaceae</i>	F: CATTGACGTTACCCGCAGAAGAAGC R: CTCTACGAGACTCAAGCTTGC	195	(12)
<i>Bacteroides</i>	F: GTCAGTTGTGAAAGTTTGC R: CAATCGGAGTTCTTCGTG	127	(13)

PCR, polymerase chain reaction; bp, base pairs; F, forward; R, reverse.

Table III. Effects of Amano protease and orientase on the profile of microflora in the cecal digesta of rats fed an HF diet<sup>a</sup>.

Bacteria	Control (% of total bacteria)	Amano protease (% of total bacteria)	Orientase (% of total bacteria)
<i>Bifidobacterium</i> spp.	0.016±0.002 <sup>b</sup>	3.105±0.758 <sup>c</sup>	0.882±0.203 <sup>b,c</sup>
<i>Lactobacillus</i> spp.	3.5±0.5 <sup>b</sup>	26.7±3.0 <sup>d</sup>	13.7±2.1 <sup>c</sup>
<i>Clostridium</i>			
<i>Clostridium coccooides</i> group	6.60±0.07 <sup>b,c</sup>	3.88±0.85 <sup>b</sup>	9.48±1.32 <sup>c</sup>
<i>Clostridium leptum</i> group	0.263±0.071 <sup>b</sup>	0.004±0.001 <sup>c</sup>	0.034±0.006 <sup>c</sup>
<i>Akkermansia muciniphila</i>	1.332±0.512 <sup>b</sup>	0.0009±0.0004 <sup>c</sup>	0.601±0.278 <sup>b</sup>
<i>Enterobacteriaceae</i>	0.21±0.09 <sup>b</sup>	8.24±0.84 <sup>c</sup>	6.57±1.70 <sup>c</sup>
<i>Bacteroides</i>	14.0±1.79 <sup>b</sup>	1.10±0.69 <sup>c</sup>	3.57±1.65 <sup>c</sup>

<sup>a</sup>Mean ± standard error (n=9). <sup>b-d</sup>Significantly different by Tukey's multiple-range test (P<0.05). HF, high-fat.

proglucagon forward, GGTTGATGAACACCAAGAGG and reverse, GAAGGATCCATCAGCATGTC; *GAPDH* forward, TGACAACCTCCCTCAAGATTGTCA and reverse, GGCATGGACTGTGGTCATGA. The expression of the target genes, FIAF and proglucagon, was normalized to that of *GAPDH*, the endogenous control gene (18).

**Statistical analysis.** Data are expressed as mean ± standard error. Statistical analysis was performed by one-way analysis of variance and Tukey's post hoc test (Excel Statistics 2010 for Windows; Social Survey Research Information, Tokyo, Japan). P<0.05 was considered to indicate a statistically significant difference.

## Results

**Characteristics of the groups.** The final body weights of the control, Amano protease and orientase groups were 254±2, 246±2 and 252±2 g, respectively (P>0.05). In addition, food intake, epididymal and perirenal adipose tissue weight did not differ significantly among the groups (data not shown).

**Proportions of bacteria.** The proportions of *Bifidobacterium*, *Lactobacillus*, *Clostridium*, *Akkermansia muciniphila*, *Enterobacteriaceae*, and *Bacteroides* in cecal digesta are shown in Table III. The proportions of cecal microflora were markedly different in the Amano protease and orientase

Table IV. Effects of Amano protease and orientase on the cecal organic acids in rats fed an HF diet<sup>a</sup>.

Variables	Control	Amano protease	Orientase
Cecal digesta, g	2.21±0.10 <sup>b</sup>	7.02±0.29 <sup>d</sup>	4.10±0.33 <sup>c</sup>
pH of cecal digesta	7.16±0.10 <sup>b</sup>	5.32±0.12 <sup>c</sup>	5.46±0.11 <sup>c</sup>
Organic acids, $\mu$ mol/g wet cecal digesta			
<i>n</i> -Butyrate	18.6±1.8 <sup>b</sup>	78.9±7.9 <sup>c</sup>	58.9±7.2 <sup>c</sup>
Propionate	10.9±0.9 <sup>b</sup>	35.5±4.2 <sup>c</sup>	28.7±3.6 <sup>c</sup>
Lactate	4.7±0.6 <sup>b</sup>	38.7±3.5 <sup>c</sup>	31.9±6.2 <sup>c</sup>
Acetate	31.2±3.8 <sup>b</sup>	19.4±1.7 <sup>c</sup>	24.2±3.9 <sup>b,c</sup>
Succinate	18.3±3.5 <sup>b</sup>	13.6±2.6 <sup>b,c</sup>	7.9±1.5 <sup>c</sup>
Total organic acids	84±7 <sup>b</sup>	186±12 <sup>d</sup>	152±14 <sup>c</sup>
Ammonia, $\mu$ mol/g wet cecal digesta	3.84±0.28	3.38±0.21	4.25±0.36

<sup>a</sup>Mean  $\pm$  standard error (n=9). <sup>b-d</sup>Significantly different by Tukey's multiple-range test (P<0.05). HF, high-fat.

Table V. Effects of Amano protease and orientase on the fecal parameters of rats fed an HF diet<sup>a</sup>.

Variables	Control	Amano protease	Orientase
Dry wt, g/3 days <sup>b</sup>	3.39±0.07 <sup>c</sup>	4.13±0.26 <sup>d</sup>	2.91±0.17 <sup>c</sup>
IgA, mg/g dry wt	0.267±0.025 <sup>c</sup>	0.634±0.057 <sup>d</sup>	0.500±0.092 <sup>c</sup>
IgA, mg/3 days <sup>b</sup>	0.90±0.09 <sup>c</sup>	2.37±0.15 <sup>d</sup>	1.21±0.19 <sup>c</sup>
Mucins, mg/g dry wt	0.44±0.03 <sup>c</sup>	4.55±0.45 <sup>d</sup>	1.11±0.23 <sup>c</sup>
Mucins, mg/3 days <sup>b</sup>	1.48±0.09 <sup>c</sup>	18.85±2.26 <sup>d</sup>	3.35±0.85 <sup>c</sup>

<sup>a</sup>Mean  $\pm$  standard error (n=9). <sup>b</sup>Fecal collection for final 3 days. <sup>c,d</sup>Significantly different by Tukey's multiple-range test (P<0.05). HF, high-fat; IgA, immunoglobulin A.

Table VI. Effects of Amano protease and orientase on the colonic gene expression in rats fed an HF diet<sup>a</sup>.

Genes	Relative expression		
	Control	Amano protease	Orientase
FIAF	1.00±0.22	0.96±0.12	0.73±0.14
Proglucagon	1.00±0.18 <sup>b</sup>	3.80±0.43 <sup>c</sup>	2.17±0.45 <sup>d</sup>

<sup>a</sup>Mean  $\pm$  standard error (n=9). <sup>b-d</sup>Significantly different by Tukey's multiple-range test (P<0.05). HF, high-fat; FIAF, fasting-induced adipose factor.

groups compared to those in the control group. The proportion of *Bifidobacterium* spp. was significantly higher in the Amano protease group compared to the control group (194-fold greater, P<0.05). The proportions of *Lactobacillus* spp. were also markedly higher in the Amano protease and orientase groups compared to the control group (7.6- and 3.9-fold, respectively, P<0.05). There were no significant changes in the proportion of *Clostridium cocoides* with protease supplementation. However, the proportion of *Clostridium cocoides* was significantly higher in the orientase group compared to the Amano protease group (P<0.05). The proportions of *Clostridium leptum* were significantly lower in the protease groups compared to the control group. The proportion of *Akkermansia muciniphila* was markedly lower in the Amano protease group compared to the control group (P<0.05). The proportions of *Enterobacteriaceae* were significantly higher in the two protease groups compared to the control group (P<0.05). The proportions of *Bacteroides* were significantly lower in the two protease groups compared to the control group (P<0.05).

**Cecal organic acids.** Compared to the control group, the cecal digesta weights were significantly greater in the Amano protease and orientase groups (3.2- and 1.9-fold greater, respectively, P<0.05) (Table IV). The pH of the cecal digesta was significantly lower in the two protease-treated groups compared to the control group (P<0.05). Total organic acid concentration was significantly higher in the Amano protease group compared to the control group; compared to the control group, *n*-butyrate, propionate and lactate concentrations were 4.2-, 3.3- and 8.2-fold greater (P<0.05), respectively, whereas acetate concentration was significantly lower (P<0.05). Similarly, total organic acid concentrations were also significantly higher in the orientase group, and compared to the control group, *n*-butyrate, propionate and lactate concentrations were 3.2-, 2.6- and 6.8-fold greater (P<0.05), respectively, whereas succinate concentration was significantly lower (P<0.05). Cecal ammonia content did not differ significantly among the 3 groups.

**Fecal IgA and mucins.** Fecal dry weight was significantly greater in the Amano protease-treated group compared to the other groups (P<0.05; Table V). IgA and mucins were measured in the fecal samples as indices of colonic immune and barrier functions. As a result, fecal IgA and mucin levels were also significantly higher in the Amano protease-treated group compared to the other groups (P<0.05). The proportion of *Akkermansia muciniphila*, a mucin-degrading commensal bacterium, was inversely correlated with mucin release ( $r=-0.466$ , P<0.05).

**Colonic gene expression.** Compared to the control group, colonic proglucagon gene expression was 3.8- and 2.2-fold higher in the Amano protease and orientase groups, respectively (P<0.05; Table VI). FIAF expression did not differ among the 3 groups.

## Discussion

The present results indicate that the consumption of Amano protease, which consists of neutral proteases derived from

*Aspergillus* spp., improves the colonic luminal parameters favorable to colonic health. In general, shifts in colonic microflora composition are one of the numerous factors involved in the development of colonic diseases. In particular, *Bifidobacterium* and *Lactobacillus* are considered to have important roles in colonic health (19). In the present study, Amano protease supplementation with an HF diet markedly elevated the proportions of cecal *Bifidobacterium* and *Lactobacillus*, as well as concentrations of organic acids such as *n*-butyrate, propionate and lactate. As lactate is absorbed more slowly in the gut compared to other organic acids (20), the pH of the cecal digesta was considerably lower; such an acidic environment favors acid-resistant bacteria including *Bifidobacterium* and *Lactobacillus*. An HF diet causes lower colonic organic acids, and is considered to increase the risk of colon cancer (21). Propionate and *n*-butyrate are also considered to have important roles in colonic health (22). In particular, butyrate is shown to modulate cell proliferation, apoptosis and activity of immune cells in the gut epithelial layer (23,24). Thus, the present results suggest dietary Amano protease preparation has a favorable effect on cecal fermentation when rats were fed an HF diet.

Some of the end products of protein and amino acids metabolism are considered toxic to the host, such as ammonia and phenolic compounds. However, the present results showed that the cecal ammonia content did not differ significantly among the 3 groups. It has been suggested that elevation in cecal organic acids and a lower pH are associated with suppression of ammonia-producing bacteria (25). Thus, supplementation of 0.2% Amano protease may not alter the ammonia level by modulating the amount of ammonia-producing bacteria in the colon.

Kondo *et al* (26) recently reported that intake of *Bifidobacterium breve* B-3 induces proglucagon expression in the mouse colon. Proglucagon is suggested to be a regulatory signal that controls energy homeostasis. Therefore, whether an increased proportion of *Bifidobacterium* caused by protease treatment increases proglucagon expression was determined in the present study. As expected, the results indicated that Amano protease and orientase treatment significantly increased proglucagon gene expression in the colon.

The high production of intestinal IgA and mucins is associated with a lower risk of colon cancer (27,28). The present study further shows that the consumption of Amano protease increases fecal IgA and mucin levels. The present results suggest that Amano protease may have a favorable effect on colonic immune and barrier functions. The administration of certain *Bifidobacterium* species, such as *B. bifidum*, has been reported to enhance intestinal IgA production (4). Elevated IgA levels may be associated with elevated *Bifidobacterium* levels. The decrease in colonic pH due to increased organic acid production may be responsible for increased mucin production (29). A study using a rat colon model shows that acetate and butyrate stimulate mucin release, although the underlying mechanism remains unknown (30). In the present study, the two proteases decreased pH and increased concentrations of certain organic acids in the cecal digesta. However, only Amano protease significantly increased mucin release; the reason why Orientase did not affect fecal mucin levels is

unknown. Notably, Amano protease supplementation markedly decreased the cecal proportion of *Akkermansia muciniphila*, a mucin-degrading commensal bacterium. Furthermore, the proportion of *Akkermansia muciniphila*, a mucin-degrading commensal bacterium, was inversely correlated with mucin release. Therefore, the higher mucin level in the Amano protease group may be, at least in part, associated with the reduction in the proportion of *Akkermansia muciniphila*.

Mitsuoka (31) reported that bacterial extracellular compounds and the metabolites, which are defined as biogenics, may exert beneficial effects on the balance of colonic microflora by elevating the bifidobacteria and/or reducing the harmful bacteria, resulting in improving colon health. The present results suggest that the extracellular products, including several proteases released by *Aspergillus*, may have an important potential as biogenics. However, the study did not indicate any information regarding the underlying mechanism of improved colonic microflora, fermentation, immune and barrier functions by supplemental *Aspergillus*-derived protease preparations. Further study is required to investigate the mechanisms.

In conclusion, the present study suggests that the *Aspergillus*-derived protease preparations may beneficially modify the composition of cecal microflora, organic acids, IgA and mucins in rats fed an HF diet. The powdered protease preparations used contain several proteases and other factors derived from *Aspergillus*. Therefore, further study is in progress to investigate the effects of purified *Aspergillus* proteases on the colonic luminal environment.

## References

- de Vrese M and Schrezenmeir J: Probiotics, prebiotics and synbiotics. *Adv Biochem Eng Biotechnol* 111: 1-66, 2008.
- Ziegler TR, Evans ME, Fernández-Estívariz C and Jones DP: Trophic and cytoprotective nutrition for intestinal adaptation, mucosal repair, and barrier function. *Annu Rev Nutr* 23: 229-261, 2003.
- Shimotodome A, Meguro S, Hase T, Tokimitsu I and Sakata T: Short chain fatty acids but not lactate or succinate stimulate mucus release in the rat colon. *Comp Biochem Physiol A Mol Integr Physiol* 125: 525-531, 2000.
- Park JH, Um JI, Lee BJ, Goh JS, Park SY, Kim WS and Kim PH: Encapsulated *Bifidobacterium bifidum* potentiates intestinal IgA production. *Cell Immunol* 219: 22-27, 2002.
- Li D, Kim JM, Jin Z and Zhou J: Prebiotic effectiveness of inulin extracted from edible burdock. *Anaerobe* 14: 29-34, 2008.
- Reddy BS: Dietary fat and its relationship to large bowel cancer. *Cancer Res* 41: 3700-3705, 1981.
- Okazaki Y, Sitanggang NV, Sato S, Ohnishi N, Inoue J, Iguchi T, Watanabe T, Tomotake H, Harada K and Kato N: Burdock fermented by *Aspergillus awamori* elevates cecal *Bifidobacterium*, and reduces fecal deoxycholic acid and adipose tissue weight in rats fed a high-fat diet. *Biosci Biotechnol Biochem* 77: 53-57, 2013.
- Parnell JA and Reimer RA: Prebiotic fibres dose-dependently increase satiety hormones and alter Bacteroidetes and Firmicutes in lean and obese JCR:LA-cp rats. *Br J Nutr* 107: 601-613, 2012.
- Delroisse JM, Boulvin AL, Parmentier I, Dauphin RD, Vandenbol M and Portetelle D: Quantification of *Bifidobacterium* spp. and *Lactobacillus* spp. in rat fecal samples by real-time PCR. *Microbiol Res* 163: 663-670, 2008.
- Matsuki T, Watanabe K, Fujimoto J, Takada T and Tanaka R: Use of 16S rRNA gene-targeted group-specific primers for real-time PCR analysis of predominant bacterial human feces. *Appl Environ Microbiol* 70: 7220-7228, 2004.
- Sonoyama K, Ogasawara T, Goto H, Yoshida T, Takemura N, Fujiwara R, Watanabe J, Ito H, Morita T, Tokunaga Y, *et al*: Comparison of gut microbiota and allergic reactions in BALB/c mice fed different cultivars of rice. *Br J Nutr* 103: 218-226, 2010.

12. Bartosch S, Fite A, Macfarlane GT and McMurdo MET: Characterization of bacterial communities in feces from healthy elderly volunteers and hospitalized elderly patients by using real-time PCR and effects of antibiotic treatment on the fecal microbiota. *Appl Environ Microbiol* 70: 3575-3581, 2004.
13. Ahmed S, Macfarlane GT, Fite A, McBain AJ, Gilbert P and Macfarlane S: Mucosa-associated bacterial diversity in relation to human terminal ileum and colonic biopsy samples. *Appl Environ Microbiol* 73: 7435-7442, 2007.
14. Okazaki Y, Tomotake H, Tsujimoto K, Sasaki M and Kato N: Consumption of a resistant protein, sericin, elevates fecal immunoglobulin A, mucins and cecal organic acids in rats fed a high-fat diet. *J Nutr* 141: 1975-1981, 2011.
15. Lin HC and Visek WJ: Large intestinal pH and ammonia in rats: Dietary fat and protein interactions. *J Nutr* 121: 832-843, 1991.
16. Bovee-Oudenhoven IM, Termont DS, Heidt PJ and Van der Meer R: Increasing the intestinal resistance of rats to the invasive pathogen *Salmonella enteritidis*: Additive effects of dietary lactulose and calcium. *Gut* 40: 497-504, 1997.
17. Crowther RS and Wetmore RF: Fluorometric assay of O-linked glycoproteins by reaction with 2-cyanoacetamide. *Anal Biochem* 163: 170-174, 1987.
18. Dharmani P, Strauss J, Ambrose C, Allen-Vercoe E and Chadee K: *Fusobacterium nucleatum* infection of colonic cells stimulates MUC2 mucin and tumor necrosis factor alpha. *Infect Immun* 79: 2597-2607, 2011.
19. Brownawell AM, Caers W, Gibson GR, Kendall CWC, Lewis KD, Ringel Y and Slavin JL: Prebiotics and the health benefits of fiber: Current regulatory status, future research and goals. *J Nutr* 142: 962-974, 2012.
20. Hoshi S, Sakata T, Mikuni K, Hashimoto H and Kimura S: Galactosylsucrose and xylosylfructoside alter digestive tract size and concentrations of cecal organic acids in rats fed diets containing cholesterol and cholic acid. *J Nutr* 124: 52-60, 1994.
21. Wu WT and Chen HL: Effects of konjac glucomannan on putative risk factors for colon carcinogenesis in rats fed a high-fat diet. *J Agric Food Chem* 59: 989-994, 2011.
22. Topping DL and Clifton PM: Short-chain fatty acids and human colonic function: Roles of resistant starch and nonstarch polysaccharides. *Physiol Rev* 81: 1031-1064, 2001.
23. Scheppach W: Effects of short chain fatty acids on gut morphology and function. *Gut* 35 (Suppl 1): S35-S38, 1994.
24. Tang Y, Chen Y, Jiang H, Robbins GT and Nie D: G-protein-coupled receptor for short-chain fatty acids suppresses colon cancer. *Int J Cancer* 128: 847-856, 2011.
25. Han KH, Azuma S and Fukushima M: In vitro fermentation of spent turmeric powder with a mixed culture of pig faecal bacteria. *Food Funct* 5: 2446-2452, 2014.
26. Kondo S, Xiao JZ, Satoh T, Odamaki T, Takahashi S, Sugahara H, Yaeshima T, Iwatsuki K, Kamei A and Abe K: Antiobesity effects of *Bifidobacterium breve* strain B-3 supplementation in a mouse model with high-fat diet-induced obesity. *Biosci Biotechnol Biochem* 74: 1656-1661, 2010.
27. Velcich A, Yang W, Heyer J, Fragale A, Nicholas C, Viani S, Kucherlapati R, Lipkin M, Yang K and Augenlicht L: Colorectal cancer in mice genetically deficient in the mucin Muc2. *Science* 295: 1726-1729, 2002.
28. Ito H, Tanabe H, Kawagishi H, Tadashi W, Yasuhiko T, Sugiyama K, Kiriyama S and Morita T: Short-chain inulin-like fructans reduce endotoxin and bacterial translocations and attenuate development of TNBS-induced colitis in rats. *Dig Dis Sci* 54: 2100-2108, 2009.
29. Van den Abbeele P, Gérard P, Rabot S, Bruneau A, El Aidy S, Derrien M, Kleerebezem M, Zoetendal EG, Smidt H, Verstraete W, *et al*: Arabinoxylans and inulin differentially modulate the mucosal and luminal gut microbiota and mucin-degradation in humanized rats. *Environ Microbiol* 13: 2667-2680, 2011.
30. Barcelo A, Claustre J, Moro F, Chayvialle JA, Cuber JC and Plaisancié P: Mucin secretion is modulated by luminal factors in the isolated vascularly perfused rat colon. *Gut* 46: 218-224, 2000.
31. Mitsuoka T: Development of functional foods. *Biosci Microbiota Food Health* 33: 117-128, 2014.