

Low hemoglobin levels are associated with upper gastrointestinal bleeding

MINORU TOMIZAWA¹, FUMINOBU SHINOZAKI², RUMIKO HASEGAWA³, YOSHINORI SHIRAI³, YASUFUMI MOTOYOSHI⁴, TAKAO SUGIYAMA⁵, SHIGENORI YAMAMOTO⁶ and NAOKI ISHIGE⁷

Departments of ¹Gastroenterology, ²Radiology, ³Surgery, ⁴Neurology, ⁵Rheumatology, ⁶Pediatrics and ⁷Neurosurgery, National Hospital Organization Shimoshizu Hospital, Yotsukaido, Chiba 284-0003, Japan

Received May 20, 2016; Accepted July 17, 2016

DOI: 10.3892/br.2016.727

Abstract. Upper gastrointestinal (GI) bleeding can be fatal. Blood test variables were reviewed in search of threshold values to detect the presence of occult upper GI bleeding. The records of 1,023 patients who underwent endoscopy at the National Hospital Organization Shimoshizu Hospital from October 2014, to September 2015, were retrospectively reviewed. Of those, 95 had upper GI bleeding. One-way analysis of variance was applied to blood test variables comparing patients with and without upper GI bleeding. Logistic regression analysis was applied to detect the association of blood test parameters with upper GI bleeding, and receiver-operator characteristics were applied to establish threshold values. White blood cell count (WBC), platelet (Plt) count, and blood urea nitrogen (BUN) levels were higher, and hemoglobin (Hb) and albumin (Alb) levels were lower in patients with upper GI bleeding. Logistic regression analysis showed that low Hb was significantly associated with upper GI bleeding and a Hb value of 10.8 g/dl was established as the threshold for the diagnosis. In patients with upper GI bleeding, WBC, Plt count, and BUN levels were higher and Hb and Alb levels were reduced. Hb at 10.8 g/dl was established as a threshold value to detect upper GI bleeding.

Introduction

Upper gastrointestinal (GI) bleeding is defined as mucosal bleeding from the proximal portion of the esophagus to the ligament of Treitz. Upper GI bleeding is caused by gastric ulcers, duodenal ulcers, and gastric cancer (1) and can be fatal,

with a mortality rate of 3.5-7.4% (2,3). Currently, patients with upper GI bleeding are diagnosed and treated with endoscopy (4), and embolization with angiography is performed on refractory patients (5,6). When patients develop hemodynamic instability, the mortality rate rises to 40% (7). It is pivotal that patients with upper GI bleeding are identified and treated promptly to avoid hemodynamic crisis.

Identification of upper GI bleeding is essential to its management. Glasgow Batchford Scores, Modified Early Warning Score, and pre-endoscopic Rockall scores contribute to the stratification of patients at risk for hemodynamic crisis, the need for transfusion, and hospitalization (8). However, scoring systems can fail to detect patients who are hemodynamically stable despite upper GI bleeding. Not only is it important to identify acute bleeding, but also to identify bleeding caused by gastric cancer and other less acute conditions that require intervention. Blood tests are non-invasive, rapid and inexpensive and offer a group of parameters in which thresholds might suggest the presence of occult but significant upper GI bleeding.

Thus, the aim of the present study was to establish threshold values of screening blood tests to detect upper GI bleeding. A retrospective investigation was conducted to compare blood test variables between patients with or without upper GI bleeding.

Materials and methods

Patients. Patients were selected from all of those who had endoscopy at the National Hospital Organization Shimoshizu Hospital, Yotsukaido, Chiba, Japan from October 2014, to September 2015. Endoscopy was performed in 1,023 patients, 431 men (mean age 68.1±12.9 years), and 592 women (66.4±12.3 years). Indications for endoscopy included anemia, abdominal pain, and melena but also included some screening procedures. The present study was approved by the National Hospital Organization Shimoshizu Hospital Ethics Committee. Since the procedures were performed as part of our routine clinical care, written informed consent for inclusion in the study was waived. Patient records were depersonalized and retrospectively analyzed. Written informed consent was obtained for the endoscopy from all the patients.

Correspondence to: Dr Minoru Tomizawa, Department of Gastroenterology, National Hospital Organization Shimoshizu Hospital, 934-5 Shikawatashi, Yotsukaido, Chiba 284-0003, Japan
E-mail: nihminor-cib@umin.ac.jp

Key words: hemoglobin, receiver-operator characteristics, blood urea nitrogen, albumin, upper gastrointestinal bleeding

Table I. Number of patients with upper gastrointestinal bleeding.

Type	Patients (no.)
Gastric ulcer	58
Gastric cancer	9
Hemorrhagic gastritis	9
Reflux esophagitis	7
Duodenal ulcer	8
Acute gastric mucosal lesion	2
Esophageal varix	2
Total	95

Table II. Comparison of blood test variables between patients with or without upper gastrointestinal bleeding.

Variables	Normal range	Upper gastrointestinal bleeding (mean \pm SD)		P-value
		Absent	Present	
Age, years		69.3 \pm 13.9	69.3 \pm 12.9	0.983
Gender, m/f		459/812	53/42	
WBC, $10^3/\mu\text{l}$	3.5-8.5	6.6 \pm 2.2	8.4 \pm 6.2	0.020
Hb, g/dl	13.5-17.0	12.8 \pm 2.0	9.7 \pm 3.1	<0.001
Plt, $10^4/\mu\text{l}$	15.0-35.0	23.9 \pm 6.7	30.0 \pm 15.9	0.005
CRP, mg/dl	0.0-0.3	1.0 \pm 2.5	2.4 \pm 3.8	0.074
TP, g/dl	6.7-8.3	6.8 \pm 0.6	8.0 \pm 9.6	0.298
Alb, g/dl	3.9-4.9	3.9 \pm 0.8	3.3 \pm 1.0	0.005
ALP, IU/l	115-359	267 \pm 263	275 \pm 191	0.899
AST, IU/l	13-33	27 \pm 32	140 \pm 585	0.083
ALT, IU/l	8-42	23 \pm 25	122 \pm 516	0.089
γ -GTP, IU/l	10-47	51.4 \pm 89.8	108 \pm 261	0.213
LDH, IU/l	119-229	209 \pm 83	243 \pm 172	0.242
UA, mg/dl	2.3-7.0	5.17 \pm 1.52	4.46 \pm 1.59	0.090
BUN, mg/dl	8.0-22.0	14.7 \pm 5.0	18.8 \pm 12.9	0.043
Cre, mg/dl	0.40-0.70	1.78 \pm 2.02	0.77 \pm 0.26	0.490
TG, mg/dl	30-150	115 \pm 62	91 \pm 27	0.163
HDL, mg/dl	49-75	62.5 \pm 18.9	56.7 \pm 26.7	0.392
LDL (mg/dl	0-139	112 \pm 29	99 \pm 48	0.214
BG, mg/dl	69-104	111 \pm 52	118 \pm 35	0.669
HbA1c, %	4.7-6.2	5.8 \pm 0.7	7.1 \pm 3.5	0.107

SD, standard deviation; m, male; f, female; WBC, white blood cell count; Hb, hemoglobin; Plt, platelet; CRP, C-reactive protein; TP, total protein; Alb, albumin; ALP, alkaline phosphatase; AST, aspartate aminotransferase; ALT, alanine aminotransferase; γ -GTP, γ -glutamyl transpeptidase; LDH, lactate dehydrogenase; UA, uric acid; BUN, blood urea nitrogen; Cre, creatinine; TG, triglyceride; HDL, high-density lipoprotein cholesterol; LDL, low-density lipoprotein cholesterol; BG, blood glucose; HbA1c, hemoglobin A1c.

Endoscopy. The endoscopes used included Olympus models GIF-N260H, GIF-XP260NS, GIF-PG260, GIF-XQ260, and GIF-Q260 (Olympus Corp., Tokyo, Japan). Bleeding from a

Table III. Results of logistic regression analysis.

Variables	Odds ratio	95% CI	P-value
WBC	2.67	0.000-3350	0.801
Hb	49.5	1.007-6229	0.049
Plt	0.212	0.0032-9.592	0.432
Alb	2.26	0.535-122.4	0.670
BUN	0.118	0.000-16.12	0.420

CI, confidence interval; WBC, white blood cell count; Hb, hemoglobin; Plt, platelet; Alb, albumin; BUN, blood urea nitrogen.

Table IV. Discriminatory threshold value for Hb.

Variable	Area under curve	Threshold value (g/dl)	Sensitivity (%)	Specificity (%)
Hb	0.7867	10.8	62.5	91.6

Hb, hemoglobin.




Figure 1. Receiver-operator characteristics analysis. Receiver-operator characteristics analysis was performed in search of a threshold value for the identification of upper gastrointestinal bleeding. Solid straight line, a line with a slope of 45° to calculate threshold using JMP 10.0.2 software; broken line, reference.

gastric or duodenal ulcer was classified as a spurting vessel (1a), an oozing vessel (1b), a visible vessel (2a), or a clot (2b), according to the Forrest classification system (9). Table I shows the diagnosis of patients.

Blood test variables. Screening blood tests included white blood cell count (WBC), hemoglobin (Hb), platelet (Plt), C-reactive protein (CRP), total protein, albumin (Alb), alkaline phosphatase, aspartate aminotransferase, alanine aminotransferase (ALT), γ -glutamyl transpeptidase, lactate dehydrogenase, uric acid, blood urea nitrogen (BUN), creatinine (Cre), triglycerides, high-density lipoprotein cholesterol,

low-density lipoprotein cholesterol, blood glucose, and HbA1c.

Statistical analysis. One-way analysis of variance was applied to identify an association between blood test variables and the presence of upper GI bleeding. Logistic regression analysis was then used to correlate values linked to upper GI bleeding. Receiver-operator characteristics analysis was finally applied to establish threshold values. JMP software, version 10.0.2 (SAS Institute, Cary, NC, USA) was used for statistical analysis.

Results

To identify blood test variables associated with upper GI bleeding, one-way analysis of variance was performed. WBC ($P=0.02$), Plt ($P=0.005$), and BUN ($P=0.04$) were elevated and Hb ($P<0.001$) and Alb ($P=0.005$) were reduced in patients with upper GI bleeding (Table II).

Logistic regression analysis was performed to reveal blood test variables associated with upper GI bleeding (Table III). Hb had a significant association with upper GI bleeding ($P=0.049$) with an odds ratio of 45.9 (95% confidence interval, 0.007-6.229).

Since these results indicated that Hb might be a strong discriminatory parameter, receiver-operator characteristics were utilized to establish a threshold value for the existence of upper GI bleeding (Fig. 1). The area under the curve was 0.7867 and the threshold value calculated was 10.8 g/dl, with a sensitivity of 62.5% and specificity of 91.6% (Table IV).

Discussion

Blood test variables are easy to evaluate. Hoffmann *et al* (10) investigated a number of blood test variables and demonstrated that changes in WBC, Hb, Plt, CRP, ALT, and Cre levels are associated with upper GI bleeding. In the present study, WBC, Plt, and BUN were elevated, and Hb and Alb were reduced in patients with upper GI bleeding. Elevated BUN and low Alb have not been reported previously. BUN is known to be produced through protein metabolism (11,12). Blood in the GI tract is digested to protein, and is eventually metabolized to urea (13,14). It is therefore reasonable to conclude that BUN was elevated in patients with upper GI bleeding. Higher BUN levels indicate more severe upper GI bleeding (15), and elevated BUN can also discriminate upper from lower GI bleeding (16). Furthermore, Alb is a marker of nutritional state (17) and can reflect a severe state of upper GI bleeding (18). The results of the present study were consistent with these observations.

Upper GI bleeding is sometimes fatal, and the European Society of Gastrointestinal Endoscopy recommends maintaining Hb levels between 7.0-9.0 g/dl using blood transfusion in order to reduce mortality (19). In the present study, the threshold value calculated to select patients at risk for ongoing upper GI bleeding was 10.8 g/dl. Previously we calculated the threshold value of Hb to be 11.7 g/dl. Our threshold values were higher than the recommendation (19). It is possible the present study included patients with mild upper GI bleeding. Dahlerup *et al* (20) reported that mild upper GI bleeding includes patients with malignancy. Patients should

be subjected to endoscopy before they develop hemodynamic instability, and those who are hemodynamically stable but with a Hb <10.8 g/dl may be recommended for endoscopy, since diagnosis would improve management and treatment would reduce the risk of further or delayed bleeding. One limitation of the present study was that it was based on a small number of patients with upper GI bleeding. Future studies with a greater number of recruited patients should be undertaken.

In conclusion, WBC, Plt, and BUN were elevated and Hb and Alb were lower in patients with upper GI bleeding. We established that a threshold value of 10.8 g/dl of Hb suggests the presence of upper GI bleeding and identifies patients at risk for hemodynamic crisis.

References

1. Theocharis GJ, Thomopoulos KC, Sakellaropoulos G, Katsakoulis E and Nikolopoulou V: Changing trends in the epidemiology and clinical outcome of acute upper gastrointestinal bleeding in a defined geographical area in Greece. *J Clin Gastroenterol* 42: 128-133, 2008.
2. Hearnshaw SA, Logan RF, Lowe D, Travis SP, Murphy MF and Palmer KR: Acute upper gastrointestinal bleeding in the UK: Patient characteristics, diagnoses and outcomes in the 2007 UK audit. *Gut* 60: 1327-1335, 2011.
3. Leontiadis GI, Molloy-Bland M, Moayyedi P and Howden CW: Effect of comorbidity on mortality in patients with peptic ulcer bleeding: Systematic review and meta-analysis. *Am J Gastroenterol* 108: 331-345, quiz 346, 2013.
4. Kim SY, Hyun JJ, Jung SW and Lee SW: Management of non-variceal upper gastrointestinal bleeding. *Clin Endosc* 45: 220-223, 2012.
5. Wilkins T, Khan N, Nabh A and Schade RR: Diagnosis and management of upper gastrointestinal bleeding. *Am Fam Physician* 85: 469-476, 2012.
6. Katano T, Mizoshita T, Senoo K, Sobue S, Takada H, Sakamoto T, Mochiduki H, Ozeki T, Kato A, Matsunami K, *et al*: The efficacy of transcatheter arterial embolization as the first-choice treatment after failure of endoscopic hemostasis and endoscopic treatment resistance factors. *Dig Endosc* 24: 364-369, 2012.
7. Walsh RM, Anain P, Geisinger M, Vogt D, Mayes J, Grundfest-Broniatowski S and Henderson JM: Role of angiography and embolization for massive gastroduodenal hemorrhage. *J Gastrointest Surg* 3: 61-65, discussion 66, 1999.
8. Bozkurt S, Köse A, Arslan ED, Erdoğan S, Üçbilek E, Çevik İ, Ayrik C and Sezgin O: Validity of modified early warning, Glasgow Blatchford, and pre-endoscopic Rockall scores in predicting prognosis of patients presenting to emergency department with upper gastrointestinal bleeding. *Scand J Trauma Resusc Emerg Med* 23: 109, 2015.
9. Forrest JA, Finlayson ND and Shearman DJ: Endoscopy in gastrointestinal bleeding. *Lancet* 2: 394-397, 1974.
10. Hoffmann V, Neubauer H, Heinzler J, Smarczyk A, Hellmich M, Bowe A, Kuetting F, Demir M, Pelc A, Schulte S, *et al*: A novel easy-to-use prediction scheme for upper gastrointestinal bleeding: Cologne-WATCH (C-WATCH) risk score. *Medicine (Baltimore)* 94: e1614, 2015.
11. Schutz Y: Protein turnover, ureagenesis and gluconeogenesis. *Int J Vitam Nutr Res* 81: 101-107, 2011.
12. Adeva MM, Souto G, Blanco N and Donapetry C: Ammonium metabolism in humans. *Metabolism* 61: 1495-1511, 2012.
13. Mardini H and Record C: Pathogenesis of hepatic encephalopathy: Lessons from nitrogen challenges in man. *Metab Brain Dis* 28: 201-207, 2013.
14. Häberle J: Clinical and biochemical aspects of primary and secondary hyperammonemic disorders. *Arch Biochem Biophys* 536: 101-108, 2013.
15. Tomizawa M, Shinozaki F, Hasegawa R, Shirai Y, Motoyoshi Y, Sugiyama T, Yamamoto S and Ishige N: Patient characteristics with high or low blood urea nitrogen in upper gastrointestinal bleeding. *World J Gastroenterol* 21: 7500-7505, 2015.
16. Tomizawa M, Shinozaki F, Hasegawa R, Shirai Y, Motoyoshi Y, Sugiyama T, Yamamoto S and Ishige N: Laboratory test variables useful for distinguishing upper from lower gastrointestinal bleeding. *World J Gastroenterol* 21: 6246-6251, 2015.

17. Lee JL, Oh ES, Lee RW and Finucane TE: Serum albumin and prealbumin in calorically restricted, nondiseased individuals: A systematic review. *Am J Med* 128: 1023.e1-22, 2015.
18. Tung CF, Chow WK, Chang CS, Peng YC and Hu WH: The prevalence and significance of hypoalbuminemia in non-variceal upper gastrointestinal bleeding. *Hepatogastroenterology* 54: 1153-1156, 2007.
19. Gralnek IM, Dumonceau JM, Kuipers EJ, Lanas A, Sanders DS, Kurien M, Rotondano G, Hucl T, Dinis-Ribeiro M, Marmo R, *et al*: Diagnosis and management of nonvariceal upper gastrointestinal hemorrhage: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. *Endoscopy* 47: a1-a46, 2015.
20. Dahlerup JF, Eivindson M, Jacobsen BA, Jensen NM, Jørgensen SP, Laursen SB, Rasmussen M and Nathan T: Diagnosis and treatment of unexplained anemia with iron deficiency without overt bleeding. *Dan Med J* 62: C5072, 2015.