

Changes in FABP1 and gastrin receptor expression in the testes of rats that have undergone electrical injury

XUE-FANG LI¹, QUAN-YONG HUANG² and SHUI-PING LIU³

¹Library of Dali University, Dali, Yunnan 671003; ²Department of Pathology, School of Basic Medical Sciences, Dali University, Dali, Yunnan 671000; ³Department of Forensic Pathology, Zhongshan School of Medicine, Sun Yat-Sen University, Guangzhou, Guangdong 510080, P.R. China

Received July 3, 2014; Accepted March 16, 2015

DOI: 10.3892/etm.2015.2382

Abstract. Testicular trauma may occur due to accidental electrical injury. The aim of this study was to investigate alterations in the levels of fatty acid-binding protein 1 (FABP1) and gastrin receptor (gastrin R) in the testes following electrical injury. Sprague-Dawley rats were divided into control, fatal electrocution (220 V, 50 Hz, 60 sec) and electrical injury (220 V, 50 Hz, 60 sec) groups (n=8 per group). The animals in the fatal electrocution and electrical injury groups were deeply anesthetized with sodium pentobarbital prior to each treatment, in which the current was delivered via an anode connected to the left foreleg and a cathode to the right hindleg. The rats that survived were subsequently sacrificed by cervical dislocation. Control animals received cervical dislocation alone. Immunohistochemical analysis was performed to evaluate the protein expression of FABP1 and gastrin R in the testes. Sections were evaluated by digital image analysis. The expression levels of FABP1 and gastrin R were significantly increased following electrical injury, supported by an increase in the integrated optical density (IOD) when compared with that in the control group ($P<0.05$). However, no significant difference was found in FABP1 and gastrin R expression levels between the fatal electrocution and control groups. In summary, the protein expression levels of FABP1 and gastrin R were found to be significantly altered by electrical injury, suggesting that these two proteins may be important in underlying mechanisms of testicular injury during electrical injury. The findings indicate that such alterations would be reflected in abnormal testicular function.

Introduction

Electrical stimulation plays an important role in the cure of disease and relief of pain. It has been reported that patients with spinal cord injury (SCI) have many factors that are associated with pressure ulcer formation, including paralysis, loss of sensation, poor nutrition, anemia, and skin maceration associated with incontinence (1). Chronic pain in SCI is disabling and resistant to common pharmacologic approaches. It has been demonstrated that electrical and magnetic neural stimulation techniques are potential tools for use in the management of patients with this condition (2,3). The stimulation of leg and gluteal muscles during functional electrical stimulation cycling increases ankle excursion in individuals with spinal cord injury (4).

However, it has been shown that electrical injuries induce progressive tissue loss (5). Electrical injuries are devastating and are challenging to treat due to the complexity of the tissue damage and physiological impact (6). When the injury occurs at a high voltage, significant damage occurs to blood vessels via a number of mechanisms; the rupture of a major vessel is a rare but life-threatening sequela of electrical injury (7). Following high-voltage electrical injury, the endothelial and smooth muscle functions of the brachial artery are significantly decreased (8). In addition, electrical injury is reportedly a major cause of morbidity (9).

It has previously been shown that the effects of electrical injury include changes in the levels of type III collagen expressed in cardiac tissue (10); however, little is known about the changes in fatty acid-binding protein 1 (FABP1) and gastrin receptor (gastrin R) expression in testes that have undergone electrical injury.

The current study used immunohistochemistry to evaluate the expression of FABP1 and gastrin R in the testes of electrically injured rats, in order to determine whether their expression at the protein level is altered by electrical injury.

Materials and methods

Animal care. A total of 24 Sprague-Dawley rats, weighing 180-200 g, were provided by Sun Yat-Sen University (Guangzhou, China). All animals were given free access to standard pellet chow and water prior to the experiments. All

Correspondence to: Dr Quan-Yong Huang, Department of Pathology, School of Basic Medical Sciences, Dali University, Building 1F, Block D, Xiaguan Wanhua Road, Dali, Yunnan 671000, P.R. China
E-mail: hqy0726@126.com

Key words: fatty acid-binding protein 1, gastrin receptor, testis, rats, electrical injury

procedures described in this study were approved by the Ethics Committee of Sun Yat-Sen University (Guangzhou, China).

Animal treatment and study design. The 24 Sprague-Dawley rats were divided into three groups, namely a control group, a fatal electrocution group and an electrical injury group ($n=8$ per group). Sixteen of the rats were deeply anesthetized with sodium pentobarbital. A TMB 1000VA control transformer was used to provide electrical current via an anode and cathode (Zhejiang 001 Group Co., Ltd., Quzhou, Zhejiang, China). With the anode connected to the left foreleg and the cathode to the right hindleg, the rats were electrocuted (220 V, 50 Hz) for 60 sec. The 8 rats that died were defined as the fatal electrocution group, and the others were defined as the electrical injury group. The rats in the electrical injury group were sacrificed by cervical dislocation. In the control group, 8 rats were sacrificed by cervical dislocation without any prior electrical stimulus. Following sacrifice, the rat testes were rapidly excised and perfused with 10% neutral-buffered formalin solution (10).

Histopathological examinations. Specimens of the testes from each group were removed for histopathological examination. The testicular tissues were fixed in phosphate-buffered 4% formalin (pH 7.4) for 24 h and then embedded in paraffin. Sections were cut into 4- μ m slices, and stained with hematoxylin and eosin. The slides were coded and semiquantitative analysis of the sections was performed by a pathologist who was blinded to the treatment that the rats had received. The pathological changes in these testicular tissues were then evaluated (11,12).

Tissue sections and immunohistochemical staining. All rat testes were immersed in 10% neutral-buffered formalin solution for 24 h, prior to being embedded in paraffin and sectioned with a microtome into 4- μ m sections. All rat testes were investigated by immunohistochemistry to determine the difference between the electrical injury and control groups or the fatal electrocution and control groups with respect to FABP1 and gastrin R expression. The immunohistochemical analyses were performed as described previously (13-15). The primary antibodies were mouse monoclonal anti-FABP1 [sc-271591; L-FABP (F-9); Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA] and anti-gastrin R (BA2104; Wuhan Boster Biological Technology, Ltd., Wuhan, China), which were used at a dilution of 1:400.

The total integrated optical density (IOD), a parameter representing the expression levels of FABP1 and gastrin R in the testicular tissue, was determined using a Bx41 cast-grid microscope with DP10 camera (Olympus, Tokyo, Japan) together with an image-analysis program (MetaMorph offline, version 4.65; Molecular Devices, Sunnyvale, CA, USA). Under a magnification of $\times 200$, five images were examined in each immunostained section and the average IOD was calculated (15-17).

Statistical analysis. Results are expressed as mean \pm standard error of the mean. The significance of differences in total IOD values was tested by Kruskal-Wallis analysis. $P<0.05$ was considered to indicate a statistically significant difference. All

Table I. IOD of FABP1 and gastrin receptor in immunohistochemically stained rat testes.

Groups	FABP1	Gastrin receptor
Control	0.0069 \pm 0.00035	0.0066 \pm 0.00048
Fatal electrocution	0.0077 \pm 0.00041	0.0075 \pm 0.00052
Electrical injury	0.0134 \pm 0.00056 ^a	0.0126 \pm 0.00033 ^a

Results presented are the IOD per field, which is proportional to the total amount of staining. ^a $P<0.05$ vs. control group. The total IOD values for FABP1 and gastrin R in the rats subjected to electrical injury were significantly higher compared with those in the control rats. IOD, integrated optical density; FABP1, fatty acid-binding protein 1.

analyses were performed using SPSS software, version 12.0 (SPSS Inc., Chicago, IL, USA).

Results

Histological examination. Routine histological examination revealed little morphological change in the rat testes from each group (not shown).

Expression of FABP1 protein. The distribution of FABP1 in the rat testes of the control (Fig. 1A), fatal electrocution (Fig. 1B) and electrical injury (Fig. 1C) groups is displayed in Fig. 1. The total IOD for FABP1 in the testes of animals subjected to electrical injury (0.0134 \pm 0.00056) was significantly higher than that of the testes in the control group (0.0069 \pm 0.00035; $P<0.05$; Table I). The expression level of FABP1 in the control group was the lowest among the three groups; however, no significant difference in FABP1 expression was found between the fatal electrocution (0.0077 \pm 0.00041) and control groups.

Expression of gastrin R protein. The distribution of gastrin R protein in the rat testes of the control (Fig. 2A), fatal electrocution (Fig. 2B) and electrical injury (Fig. 2C) groups is displayed in Fig. 2. The total IOD of gastrin R in the testes of animals subjected to electrical injury (0.0126 \pm 0.00033) was significantly higher than that of control testes (0.0066 \pm 0.00048; $P<0.05$; Table I). The expression level of gastrin R in the control group was the lowest among the three groups; however, no significant difference in gastrin R expression was found between the fatal electrocution (0.0075 \pm 0.00052) and control groups.

Discussion

FABP1, also known as liver fatty acid-binding protein (L-FABP) has been demonstrated to be strongly associated with several diseases. FABP1 is a highly conserved key factor in lipid metabolism (18). The structures, functions and expression of a whole family of FABPs have been investigated extensively (19). Plasma and especially urinary levels of intestinal (I)-FABP and L-FABP and urinary levels of ileal bile acid binding protein (I-BABP) can improve the early diagnosis of intestinal ischemia, and plasma I-BABP levels can assist in




Figure 1. Effect of electrical injury on FABP1 expression. Photomicrographs display a representative distribution of positive FABP1 expression in the testes of (A) control, (B) fatal electrocution and (C) electrical injury group rats. Magnification, x200. Positive immunostaining appears as brown staining. Total FABP1 expression in rat testes subjected to electrical injury was significantly higher compared with that in control testes. FABP1, fatty acid-binding protein 1.




Figure 2. Effect of electrical injury on gastrin receptor (gastrin R) expression. Photomicrographs showing a representative distribution of positive gastrin R expression in the testes of (A) control, (B) fatal electrocution and (C) electrical injury group rats. Magnification, x200. Positive immunostaining appears as brown staining. Total gastrin R expression in rat testes subjected to electrical injury was significantly higher compared with that in control testes.

the localization of ileal ischemia (20). It has been suggested that L-FABP may have an important role in the prevention of age- or diet-induced obesity (21). A previous study has indicated that alterations in FABP1 may be associated with the development of aspirin-exacerbated respiratory disease (22). The FABP1 gene is highly transcribed in liver-derived cells, and regulated predominantly by liver-enriched transcription factors HNF3 β and C/EBP α (23). A role of L-FABP in the attenuation of hepatic steatosis has also been demonstrated (24). SCP-2 expression has also been indicated to play a significant role in high density lipoprotein-mediated cholesterol efflux by regulating the size of rapid versus slow cholesterol efflux pools and/or eliciting the concomitant upregulation of L-FABP in cultured primary hepatocytes (25).

FABP1 serves as a key regulator of hepatic lipid metabolism, and FABP1 gene polymorphisms have been associated with several metabolic traits (26). It has also been shown that phosphorylation of Sar1b disrupts the FABP1-containing four-membered 75-kDa protein complex in the cytosol, enabling it to bind to the endoplasmic reticulum and generate pre-chylomicron transport vesicle (27). FABP1 plays an important role in the male reproductive system (28). In the present study, FABP1 expression was found to be significantly increased in electrically injured rat testes, indicating that elevated FABP1 expression is associated with testicular injury.

Gastrin is a peptide hormone, which regulates gastric acid secretion and exerts physiological actions such as the regulation of sodium balance (29). Gastrin has been suggested to be involved in blood pressure regulation, possibly via the regulation of sodium and water metabolism and/or the renin-angiotensin-aldosterone system (29). Measurements

of gastrin levels are taken primarily for the diagnosis of gastrin-producing tumors (gastrinomas), which cause Zollinger-Ellison syndrome. Gastrin circulates as several bioactive peptides, and the peptide pattern in gastrinoma patients often deviates from that in normal individuals (30). Gastrin and its precursors have been shown to promote mitogenesis and angiogenesis in gastrointestinal tumors (31).

Gastrin-releasing peptide (GRP), which is an unfolded protein response regulator that functions as a Ca²⁺-binding molecular chaperone in the endoplasmic reticulum, is a regulatory human peptide that induces the release of gastrin and regulates gastric acid secretion and enteric motor function (32). GRP is a member of the bombesin-like peptide family. It has been reported that GRP stimulates the proliferation and invasiveness of androgen-independent prostate carcinoma (33). GRP mediates its action through the membrane-bound receptor, GRP receptor (GRPR), which is characterized by high-affinity binding for GRP and bombesin (33). GRPR is an attractive target for therapeutic and diagnostic applications, and is overexpressed in prostate cancer (34). The expression of GRPR is elevated in mucosa adjacent to head and neck squamous cell carcinoma (HNSCC) compared with mucosa from cancer-free controls, suggesting that elevated GRPR expression may indicate presence of HNSCC (35). GRP acts as a neuropeptide through G protein-coupled receptors involved in signal transmission in the central and peripheral nervous systems (36). It has also been proposed that GRPR is an alternative chemotactic receptor that may be involved in the pathogenesis of inflammatory disorders (36). In addition, gastrin plays an important role in the male reproductive system (37,38). In the current

study, it was revealed that the expression level of gastrin R following electrical injury was higher than that in the control testes, indicating that elevated gastrin R levels are associated with testicular injury.

In the present study, it was demonstrated that the levels of expression of FABP1 and gastrin R in the testes were altered in electrically injured rats. Following an electrical injury, the expression levels of FABP1 and gastrin R were significantly elevated in the rat testes, indicating an association with testicular trauma in rats undergoing electrical injury.

In conclusion, the current findings indicate that such alterations would be reflected in abnormal testis function.

Acknowledgements

This study was supported by a grant from the National Natural Science Foundation of China (no. 81260466) and Dali University (no. KYBS201104).

References

- Recio AC, Felter CE, Schneider AC and McDonald JW: High-voltage electrical stimulation for the management of stage III and IV pressure ulcers among adults with spinal cord injury: Demonstration of its utility for recalcitrant wounds below the level of injury. *J Spinal Cord Med* 35: 58-63, 2012.
- Moreno-Duarte I, Morse LR, Alam M, *et al*: Targeted therapies using electrical and magnetic neural stimulation for the treatment of chronic pain in spinal cord injury. *Neuroimage* 85: 1003-1013, 2014.
- Gorgey AS, Harnish CR, Daniels JA, *et al*: A report of anticipated benefits of functional electrical stimulation after spinal cord injury. *J Spinal Cord Med* 35: 107-112, 2012.
- Fornusek C, Davis GM and Baek I: Stimulation of shank muscles during functional electrical stimulation cycling increases ankle excursion in individuals with spinal cord injury. *Arch Phys Med Rehabil* 93: 1930-1936, 2012.
- Benlier E, Eskiocak S, Puyan FO, *et al*: Effect of lidocaine on reducing injury in a rat electrical burn model. *Ann Plast Surg* 69: 152-156, 2012.
- Shupp JW, Moffatt LT, Nguyen T, *et al*: Examination of local and systemic in vivo responses to electrical injury using an electrical burn delivery system. *J Burn Care Res* 33: 118-129, 2012.
- Toy J, Ball BJ and Tredget EE: Carotid rupture following electrical injury: a report of two cases. *J Burn Care Res* 33: e160-e165, 2012.
- Park KH, Park WJ, Kim MK, *et al*: Alterations in arterial function after high-voltage electrical injury. *Crit Care* 16: R25, 2012.
- Mashreky SR, Hossain MJ, Rahman A, *et al*: Epidemiology of electrical injury: Findings from a community based national survey in Bangladesh. *Injury* 43: 113-116, 2012.
- Huang QY, Chen YC and Liu SP: Connexin 43, angiotensin II, endothelin 1, and type III collagen alterations in heart of rats having undergone fatal electrocution. *Am J Forensic Med Pathol* 33: 215-221, 2012.
- Helin HO, Lundin ME, Laakso M, *et al*: Virtual microscopy in prostate histopathology: Simultaneous viewing of biopsies stained sequentially with hematoxylin and eosin and alpha-methylacyl-coenzyme A racemase/p63 immunohistochemistry. *J Urol* 175: 495-499, 2006.
- De Rossi A, Rocha LB and Rossi MA: Application of fluorescence microscopy on hematoxylin and eosin-stained sections of healthy and diseased teeth and supporting structures. *J Oral Pathol Med* 36: 377-381, 2007.
- Huang QY, Li XF and Liu SP: E-cadherin and 5-HT alterations in the heart of rats having undergone atropine-induced toxicity. *Mol Med Rep* 5: 700-704, 2012.
- Huang QY, Li XF and Liu SP: Elevated enolase and caveolin-1 in the heart of rats following dexamethasone-induced toxicity. *Mol Med Rep* 5: 1232-1236, 2012.
- Huang QY, Li XF and Liu SP: Connexin43 and angiotensin II alterations in hearts of rats having undergone an acute exposure to alcohol. *Am J Forensic Med Pathol* 34: 68-71, 2013.
- Dong J, Yin H, Liu W, Wang P, Jiang Y and Chen J: Congenital iodine deficiency and hypothyroidism impair LTP and decrease C-fos and C-jun expression in rat hippocampus. *Neurotoxicology* 26: 417-426, 2005.
- van Kuijk AW, Gerlag DM, Vos K, *et al*: A prospective, randomised, placebo-controlled study to identify biomarkers associated with active treatment in psoriatic arthritis: Effects of adalimumab treatment on synovial tissue. *Ann Rheum Dis* 68: 1303-1309, 2009.
- Gao N, Qu X, Yan J, Huang Q, Yuan HY and Ouyang DS: L-FABP T94A decreased fatty acid uptake and altered hepatic triglyceride and cholesterol accumulation in Chang liver cells stably transfected with L-FABP. *Mol Cell Biochem* 345: 207-214, 2010.
- Ono T and Odani S: Initial studies of the cytoplasmic FABP superfamily. *Proc Jpn Acad Ser B Phys Biol Sci* 86: 220-228, 2010.
- Thuijls G, van Wijck K, Grootjans J, *et al*: Early diagnosis of intestinal ischemia using urinary and plasma fatty acid binding proteins. *Ann Surg* 253: 303-308, 2011.
- Atshaves BP, Martin GG, Hostetler HA, McIntosh AL, Kier AB and Schroeder F: Liver fatty acid-binding protein and obesity. *J Nutr Biochem* 21: 1015-1032, 2010.
- Kim TH, Lee JY, Park JS, *et al*: Fatty acid binding protein 1 is related with development of aspirin-exacerbated respiratory disease. *PLoS One* 6: e22711, 2011.
- Wu YL, Peng XE, Wang D, Chen WN and Lin X: Human liver fatty acid binding protein (hFABP1) gene is regulated by liver-enriched transcription factors HNF3 β and C/EBP α . *Biochimie* 94: 384-392, 2012.
- Newberry EP, Kennedy SM, Xie Y, *et al*: Decreased body weight and hepatic steatosis with altered fatty acid ethanolamide metabolism in aged L-Fabp^{-/-} mice. *J Lipid Res* 53: 744-754, 2012.
- Storey SM, Atshaves BP, McIntosh AL, *et al*: Effect of sterol carrier protein-2 gene ablation on HDL-mediated cholesterol efflux from cultured primary mouse hepatocytes. *Am J Physiol Gastrointest Liver Physiol* 299: G244-G254, 2010.
- Peng XE, Wu YL, Lu QQ, Hu ZJ and Lin X: Two genetic variants in FABP1 and susceptibility to non-alcohol fatty liver disease in a Chinese population. *Gene* 500: 54-58, 2012.
- Siddiqi S and Mansbach CM II: Phosphorylation of Sar1b protein releases liver fatty acid-binding protein from multiprotein complex in intestinal cytosol enabling it to bind to endoplasmic reticulum (ER) and bud the pre-chylomicron transport vesicle. *J Biol Chem* 287: 10178-10188, 2012.
- Liu RZ, Li X and Godbout R: A novel fatty acid-binding protein (FABP) gene resulting from tandem gene duplication in mammals: Transcription in rat retina and testis. *Genomics* 92: 436-445, 2008.
- Jiang X, Wang W, Ning B, *et al*: Basal and postprandial serum levels of gastrin in normotensive and hypertensive adults. *Clin Exp Hypertens* 35: 74-78, 2013.
- Rehfeld JF, Bardram L, Hilsted L, Poitras P and Goetze JP: Pitfalls in diagnostic gastrin measurements. *Clin Chem* 58: 831-836, 2012.
- Xiao L, Kovac S, Chang M, Shulkes A, Baldwin GS and Patel O: Induction of gastrin expression in gastrointestinal cells by hypoxia or cobalt is independent of hypoxia-inducible factor (HIF). *Endocrinology* 153: 3006-3016, 2012.
- Ni C, Zhao X, Sun T, Liu Y, Gu Q and Sun B: Role of gastrin-releasing peptides in breast cancer metastasis. *Hum Pathol* 43: 2342-2347, 2012.
- Nagasaki S, Nakamura Y, Maekawa T, *et al*: Immunohistochemical analysis of gastrin-releasing peptide receptor (GRPR) and possible regulation by estrogen receptor β in human prostate carcinoma. *Neoplasia* 59: 224-232, 2012.
- Beer M, Montani M, Gerhardt J, *et al*: Profiling gastrin-releasing peptide receptor in prostate tissues: Clinical implications and molecular correlates. *Prostate* 72: 318-325, 2012.
- Egloff AM, Liu X, Davis AL, *et al*: Elevated gastrin-releasing peptide receptor mRNA expression in buccal mucosa: Association with head and neck squamous cell carcinoma. *Head Neck* 35: 270-279, 2012.
- Czepielewski RS, Porto BN, Rizzo LB, *et al*: Gastrin-releasing peptide receptor (GRPR) mediates chemotaxis in neutrophils. *Proc Natl Acad Sci USA* 109: 547-552, 2012.
- Levy R, Eustache F, Pilikian S, *et al*: Effect of gastrin-releasing peptide on sperm functions. *Mol Hum Reprod* 2: 867-872, 1996.
- Sakamoto H, Takamami K, Zuloaga DG, *et al*: Androgen regulates the sexually dimorphic gastrin-releasing peptide system in the lumbar spinal cord that mediates male sexual function. *Endocrinology* 150: 3672-3679, 2009.