

# FK506 induces biphasic $\text{Ca}^{2+}$ release from microsomal vesicles of rat pancreatic acinar cells

TERUTAKA OZAWA

Department of Physiology, Tohoku University Graduate School of Medicine,  
Seiryō-machi 2-1, Aoba-ku, Sendai 980-8575, Japan

Received January 31, 2006; Accepted March 21, 2006

**Abstract.** The effect of the immunosuppressant drug FK506 on microsomal  $\text{Ca}^{2+}$  release was investigated in rat pancreatic acinar cells. When FK506 (0.1–200  $\mu\text{M}$ ) was added to the microsomal vesicles at a steady state of ATP-dependent  $^{45}\text{Ca}^{2+}$  uptake, FK506 caused a dose-dependent and a biphasic release of  $^{45}\text{Ca}^{2+}$ . Almost 10% of total  $^{45}\text{Ca}^{2+}$  uptake was released at FK506 concentrations up to 10  $\mu\text{M}$  ( $K_m=0.47 \mu\text{M}$ ), and 60% of total  $^{45}\text{Ca}^{2+}$  uptake was released at FK506 concentrations over 10  $\mu\text{M}$  ( $K_m=55 \mu\text{M}$ ). Preincubation of the vesicles with cyclic ADP-ribose (cADPR, 0.5  $\mu\text{M}$ ) increased the FK506 ( $\leq 10 \mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release (Ozawa T, *Biochim Biophys Acta* 1693: 159–166, 2004). Preincubation with heparin (200  $\mu\text{g}/\text{ml}$ ) resulted in significant inhibition of the FK506 (30  $\mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release. Subsequent addition of inositol 1,4,5-trisphosphate ( $\text{IP}_3$ , 5  $\mu\text{M}$ ) after FK506 (100  $\mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release did not cause any release of  $^{45}\text{Ca}^{2+}$ . These results indicate that two types of FK506-induced  $\text{Ca}^{2+}$  release mechanism operate in the endoplasmic reticulum of rat pancreatic acinar cells: a high-affinity mechanism of  $\text{Ca}^{2+}$  release, which involves activation of the ryanodine receptor, and a low-affinity mechanism of  $\text{Ca}^{2+}$  release, which involves activation of the  $\text{IP}_3$  receptor.

## Introduction

Mobilization of  $\text{Ca}^{2+}$  from intracellular  $\text{Ca}^{2+}$  stores plays a crucial role in triggering enzyme secretion and muscle contraction. Inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ), which is

produced after receptor stimulation by hormones or neurotransmitters, induces  $\text{Ca}^{2+}$  release from the  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  channel ( $\text{IP}_3$  receptor:  $\text{IP}_3\text{R}$ ) on the endoplasmic reticulum (ER). The features of the  $\text{IP}_3\text{R}$  in many tissues have been characterized (1,2). The channel protein has been purified (3,4) and cloned (5,6). In addition to the  $\text{IP}_3\text{R}$ , the ryanodine-sensitive  $\text{Ca}^{2+}$  channel (ryanodine receptor:  $\text{RyR}$ ), which is activated by caffeine, ryanodine or  $\text{Ca}^{2+}$ , was first identified in excitable tissue. The channel protein has been purified (7,8) and cloned (9,10) in skeletal and cardiac muscles. The mechanism of ryanodine-sensitive  $\text{Ca}^{2+}$  release in nonexcitable tissue such as the liver and exocrine glands has recently been elucidated (11–15).

The immunosuppressant drug FK506 has been shown to increase (modulate)  $\text{Ca}^{2+}$  release through the  $\text{RyR}$  (16,17) or the  $\text{IP}_3\text{R}$  (18). In skeletal muscle sarcoplasmic reticulum (SR), the  $\text{RyR}$  is known to be tightly associated with a 12-kDa FK506-binding protein (FKBP12) (19,20). In association with the  $\text{RyR}$ , FKBP12 has been shown to stabilize the closed conformation of the  $\text{Ca}^{2+}$  release channel (16), and FK506 has been shown to promote dissociation of FKBP12 from the  $\text{RyR}$  complex (16). In cardiac muscle, FKBP12.6 has been found to be associated with  $\text{RyR}$  (21,22).

The  $\text{IP}_3\text{R}$ , a structurally related tetramer that has up to 40% sequence identity with  $\text{RyR}$  (5,6), is also associated with FKBP12 (18). The  $\text{RyR}$ -FKBP12 complex or the  $\text{IP}_3\text{R}$ -FKBP12 complex can associate with  $\text{Ca}^{2+}$ -activated phosphatase calcineurin (CN) (23,24). CN has been shown to dephosphorylate the protein kinase C (PKC) phosphorylation site on the  $\text{IP}_3\text{R}$  (23). When FKBP12 is dissociated from the  $\text{IP}_3\text{R}$  complex by FK506, binding of CN to the FK506-FKBP12 complex is stimulated (23). Thereby, the  $\text{IP}_3\text{R}$  is phosphorylated by PKC, and  $\text{Ca}^{2+}$  flux through the  $\text{IP}_3\text{R}$  is increased (23).

In rat pancreatic acinar cells, the presence of an  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  release mechanism (25,26) and the presence of a ryanodine (caffeine)-sensitive  $\text{Ca}^{2+}$  release mechanism (14,27) have been shown. Recently, it has also been shown that the presence of FK506 modulates  $\text{Ca}^{2+}$  release through a ryanodine-sensitive mechanism (28). However, details of the FK506-dependent mechanism, including the presence of FKBP, are unclear.

In the present study, the properties of microsomal  $\text{Ca}^{2+}$  release induced by the addition of FK506 were investigated in

---

*Correspondence to:* Dr Teruoka Ozawa, Department of Physiology, Tohoku University Graduate School of Medicine, Seiryō-machi 2-1, Aoba-ku, Sendai 980-8575, Japan  
E-mail: teozawa@mail.tains.tohoku.ac.jp

**Abbreviations:** cADPR, cyclic ADP-ribose; CN, calcineurin; ER, endoplasmic reticulum; FKBP, FK506-binding protein;  $\text{IP}_3$ , inositol 1,4,5-trisphosphate;  $\text{IP}_3\text{R}$ ,  $\text{IP}_3$  receptor; PKC, protein kinase C;  $\text{RyR}$ , ryanodine receptor; SR, sarcoplasmic reticulum

**Key words:** FK506, FK506-binding protein,  $\text{IP}_3$  receptor, ryanodine receptor,  $\text{Ca}^{2+}$  release, pancreatic acinar cells




Figure 1. Effect of FK506 on  $^{45}\text{Ca}^{2+}$  release from microsomal vesicles of rat pancreatic acinar cells. (A) Typical curves for FK506 (10  $\mu\text{M}$ ,  $\circ$ ; 30  $\mu\text{M}$ ,  $\blacktriangle$ ; 100  $\mu\text{M}$ ,  $\triangle$ )-induced  $^{45}\text{Ca}^{2+}$  release from the vesicles. The vesicles (1 mg/ml) were preincubated in a KCl buffer containing  $^{45}\text{Ca}^{2+}$  (1  $\mu\text{Ci}/\text{ml}$ ) for 15 min.  $^{45}\text{Ca}^{2+}$  uptake was initiated by the addition of 2 mM  $\text{K}_2\text{ATP}$  (at 0 min). FK506 was added to the vesicles as indicated. (B) Dose-response curve of FK506 (0.1–200  $\mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release from the vesicles. The vertical line represents the ratio of released  $^{45}\text{Ca}^{2+}$  over a 6-min period after the addition of FK506 to the  $^{45}\text{Ca}^{2+}$  that had been taken up by ATP. Each point is the mean  $\pm$  S.E. from 3 to 11 experiments of the type shown in (A). For values at concentrations up to 10  $\mu\text{M}$ , the percentages reported in a previous study (28) were plotted. Note that the effect of FK506 on  $^{45}\text{Ca}^{2+}$  release from the vesicles is biphasic. (C) Dose-response curve of FK506 (>10  $\mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release from the vesicles. The  $V_{\text{max}}$  value ( $V_{1\text{max}}$ : 8.1%) of the first phase was subtracted from the values at concentrations over 10  $\mu\text{M}$  shown in (B), and the corrected values were plotted. Hill plot analysis (inset) yielded a Hill coefficient ( $n$ ) of 3.0, an apparent  $K_m$  value of 55  $\mu\text{M}$  and  $V_{2\text{max}}$  value of 60%.

rat pancreatic acinar cells. It was found that there are two different types of FK506-induced  $\text{Ca}^{2+}$  release mechanism in the ER of rat pancreatic acinar cells: one mechanism, which

has a high sensitivity to FK506 ( $K_m=0.5$   $\mu\text{M}$ ), involves activation of the ryanodine-sensitive  $\text{Ca}^{2+}$  channel and the other, which has a lower sensitivity to FK506 ( $K_m=55$   $\mu\text{M}$ ), involves activation of the  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  channel.

## Materials and methods

**Materials.** Creatine kinase and trypsin inhibitor were obtained from Boehringer Mannheim (Mannheim, Germany). Adenosine triphosphate dipotassium salt ( $\text{K}_2\text{ATP}$ ), creatine phosphate disodium salt, benzamidine, D-myo-inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ) and heparin were purchased from Sigma Chemical (St. Louis, MO, USA). FK506 was kindly provided by Fujisawa Pharmaceutical Co., Ltd. (Osaka, Japan). Cyclic ADP-ribose (cADPR) was obtained from Wako Pure Chemical (Osaka, Japan). Ryanodine was obtained from Calbiochem (La Jolla, CA, USA) and  $^{45}\text{CaCl}_2$  (14–19 C/g) was purchased from New England Nuclear (Boston, MA, USA).

**Preparation of microsomal vesicles.** Pancreatic microsomal vesicles were prepared as described previously (26,27,29). Briefly, isolated acinar cells from male Wistar rats (180–200 g) were homogenized in an ice-cold 'mannitol buffer' containing 290 mM mannitol, 10 mM KCl, 5 mM HEPES, 1 mM  $\text{MgCl}_2$ , 1 mM benzamidine, 0.2 mM phenylmethylsulfonyl fluoride, and 20  $\mu\text{g}/\text{ml}$  of trypsin inhibitor, adjusted with Tris to pH 7.0. After centrifugation of the cell homogenate at 11,000  $\times$  g for 15 min, the 'fluffy layer' on top of the pellet, which is enriched by approximately twofold in the ER (26), was collected. The microsomal vesicles were kept frozen in liquid nitrogen until use. The protein concentration was measured by the method of Bradford (30) using bovine serum albumin as a standard.

**Measurement of  $^{45}\text{Ca}^{2+}$  uptake.** Isolated membrane vesicles were preincubated for 15 min at a protein concentration of 1.0 mg/ml in 0.6 ml of an incubation buffer containing 155 mM KCl, 5 mM HEPES, 0.15 mM  $\text{CaCl}_2$  (corresponding to 2  $\mu\text{M}$  free  $\text{Ca}^{2+}$  concentration), 1.0 mM EDTA, 3.57 mM  $\text{MgCl}_2$  (corresponding to 1.0 mM free  $\text{Mg}^{2+}$  concentration), 10 mM  $\text{NaN}_3$ , 5  $\mu\text{M}$  oligomycin, 5  $\mu\text{g}/\text{ml}$  antimycin A, 10 mM creatine phosphate, 8 U/ml creatine kinase, and 1  $\mu\text{Ci}/\text{ml}$  of  $^{45}\text{CaCl}_2$ , adjusted with Tris/HCl to pH 7.0 at 25°C.  $^{45}\text{Ca}^{2+}$  uptake into the vesicles was initiated by the addition of  $\text{K}_2\text{ATP}$  at a final concentration of 2 mM. After a steady state of  $^{45}\text{Ca}^{2+}$  uptake had been reached, FK506 was added to the medium. FK506 was dissolved in DMSO and was added from the stock solution in a volume not exceeding 0.5% (vol/vol). At indicated times,  $^{45}\text{Ca}^{2+}$  content of membrane vesicles was determined by a rapid filtration technique, as described previously (13). Radioactivity was determined by a liquid scintillation counter (LS6500, Beckman).

Values are presented as means  $\pm$  S.E. Statistical analysis was performed using Student's t-test.

## Results

A steady state of  $^{45}\text{Ca}^{2+}$  uptake into pancreatic microsomal vesicles was reached  $\sim 20$  min after the addition of ATP (Fig. 1A). The amount of  $^{45}\text{Ca}^{2+}$  taken up into the microsomal




Figure 2. Effect of heparin on FK506-induced  $^{45}\text{Ca}^{2+}$  release from pancreatic microsomal vesicles. The experimental procedures were the same as those described in the legend to Fig. 1A. Heparin was present from the beginning of the incubation. FK506 (3 or 30  $\mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release in the absence (Control) and that in the presence of heparin (200  $\mu\text{g/ml}$ ) were compared. The vertical line represents the ratio of released  $^{45}\text{Ca}^{2+}$  as described in the legend to Fig. 1B. Each value is the mean  $\pm$  S.E. from 5 to 9 experiments. Note that heparin significantly inhibited the  $^{45}\text{Ca}^{2+}$  release induced by 30  $\mu\text{M}$  of FK506 but not that induced by 3  $\mu\text{M}$  of FK506. \*\* indicate the level of significant difference compared to the control without heparin ( $P < 0.01$ ).

vesicles over a period of 20 min after the addition of ATP was  $5.3 \pm 0.2$  nmol/mg protein ( $n=94$ ). The addition of FK506 (0.1–200  $\mu\text{M}$ ) to the vesicles at a steady state of ATP-dependent  $^{45}\text{Ca}^{2+}$  uptake caused a dose-dependent release of  $^{45}\text{Ca}^{2+}$  (Fig. 1A). The ratio of  $^{45}\text{Ca}^{2+}$  released after the addition of FK506 to the  $^{45}\text{Ca}^{2+}$  that had been taken up by an ATP-dependent mechanism showed a biphasic change (Fig. 1B). Approximately 10% of the  $^{45}\text{Ca}^{2+}$  that had been taken up by ATP was released in the first phase at FK506 concentrations of less than 10  $\mu\text{M}$  (Fig. 1B). Lineweaver-Burk plot evaluation of the first phase showed a  $K_m$  value of 0.47  $\mu\text{M}$  and  $V_{\max}$  value of 8.1% (28). Approximately 60% of the  $^{45}\text{Ca}^{2+}$  that had been taken up by ATP was further released in the second phase at FK506 concentrations of more than 10  $\mu\text{M}$  (Fig. 1B). To estimate the real percentage of the second phase, the maximal value (8.1%) of the first phase was subtracted from the percentage at concentrations over 10  $\mu\text{M}$  shown in Fig. 1B. The corrected values are plotted in Fig. 1C. Hill plot evaluation of the second phase (Fig. 1C, inset) showed a Hill coefficient ( $n$ ) of 3.0, a  $K_m$  value of 55  $\mu\text{M}$  and  $V_{\max}$  value of 60%.

To determine the involvement of the RyR in the FK506-induced  $^{45}\text{Ca}^{2+}$  release from the vesicles, the effect of cADPR, which is known to modulate the ryanodine-sensitive  $\text{Ca}^{2+}$  release mechanism (13,28,31), on the FK506-induced  $^{45}\text{Ca}^{2+}$  release was investigated. Preincubation of the vesicles with cADPR (0.5  $\mu\text{M}$ ) shifted the dose-response curve of the FK506 ( $\leq 10 \mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release upward (28). It was shown from Lineweaver-Burk plot evaluation of the release that the presence of cADPR (0.5  $\mu\text{M}$ ) increased the  $V_{\max}$  value from 8.1% to 14.4% without affecting the  $K_m$  value (28). The extent of stimulation of FK506 (30  $\mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release by cADPR (0.5  $\mu\text{M}$ ) was nearly the same as that of the  $V_{\max}$  value of the first phase by cADPR (0.5  $\mu\text{M}$ ) [ratio of released  $^{45}\text{Ca}^{2+}$  over a 6-min period after the addition of FK506 (30  $\mu\text{M}$ ) to the  $^{45}\text{Ca}^{2+}$  that had been taken up:  $18.3 \pm 1.6\%$  ( $n=5$ ) without




Figure 3. Effect of subsequent addition of  $\text{IP}_3$  (5  $\mu\text{M}$ ) after FK506 (100  $\mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release from pancreatic microsomes. The experimental procedures were the same as those described in the legend to Fig. 1A. FK506 (100  $\mu\text{M}$ ) was added to the medium 21 min after the addition of ATP ( $\bullet$ ).  $\text{IP}_3$  (5  $\mu\text{M}$ ) was added where indicated in the absence ( $\circ$ ) and presence of FK506 ( $\bullet$ ). Note that subsequent addition of  $\text{IP}_3$  after the FK506 (100  $\mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release did not cause any release of  $^{45}\text{Ca}^{2+}$ . Curves are representative of similar experiments.

cADPR vs.  $23.2 \pm 2.1\%$  ( $n=6$ ) with cADPR (0.5  $\mu\text{M}$ )]. These data indicate that a high-affinity mechanism of the FK506-induced  $\text{Ca}^{2+}$  release activates the RyR but that a low-affinity mechanism does not.

To determine the involvement of the  $\text{IP}_3\text{R}$  in the FK506-induced  $^{45}\text{Ca}^{2+}$  release from the vesicles, the effect of heparin, an inhibitor of the  $\text{IP}_3\text{R}$  (2,3,32), on the release was investigated. Preincubation of the vesicles with heparin (200  $\mu\text{g/ml}$ ) resulted in significant ( $P < 0.01$ ) inhibition of  $^{45}\text{Ca}^{2+}$  release induced by 30  $\mu\text{M}$  of FK506 but not that induced by 3  $\mu\text{M}$  of FK506 (Fig. 2). This result indicates that a low-affinity mechanism of the FK506-induced  $\text{Ca}^{2+}$  release activates the  $\text{IP}_3\text{R}$  but that a high-affinity mechanism of that does not.

To determine whether a high concentration of FK506 induces  $\text{Ca}^{2+}$  release from the  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  pool,  $\text{IP}_3$  (5  $\mu\text{M}$ ) was added to the vesicles after 100  $\mu\text{M}$  of FK506 released  $^{45}\text{Ca}^{2+}$  from the vesicles. Fig. 3 shows that the addition of  $\text{IP}_3$  after FK506 (100  $\mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release did not cause any release of  $^{45}\text{Ca}^{2+}$ . This result indicates that a high concentration of FK506 emptied the  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  pool.

## Discussion

In the present study, according to the different sensitivities for FK506, it was demonstrated that two different types of FK506-induced  $\text{Ca}^{2+}$  release mechanism operate in the ER of rat pancreatic acinar cells; i.e., FK506 can activate the RyR at concentrations up to 10  $\mu\text{M}$  ( $K_m=0.5 \mu\text{M}$ ) and can activate the  $\text{IP}_3\text{R}$  at concentrations over 10  $\mu\text{M}$  ( $K_m=55 \mu\text{M}$ ). It is unlikely that a low concentration ( $\leq 10 \mu\text{M}$ ) of FK506 acts on the  $\text{IP}_3\text{R}$  since the component of the first phase of FK506-induced  $^{45}\text{Ca}^{2+}$  release was not inhibited by heparin (Fig. 2). It is also unlikely that a high concentration ( $> 10 \mu\text{M}$ ) of FK506 acts on the RyR since the extent of stimulation of 30  $\mu\text{M}$  FK506-induced  $^{45}\text{Ca}^{2+}$  release by cADPR (0.5  $\mu\text{M}$ ) was nearly the same as that of



$^{45}\text{Ca}^{2+}$  release induced by FK506 at concentrations of less than  $10\ \mu\text{M}$  (see Results).

FK506 is known to modulate the RyR by dissociating FKBP from the receptor (16,17). It has been shown that the presence of FK506 decreases the threshold of caffeine or ryanodine required for  $\text{Ca}^{2+}$  release in skeletal muscle cells (16) and pancreatic acinar cells (28). cADPR dissociates FKBP12.6 from the RyR in pancreatic islets (33) and coronary arterial smooth muscle cells (34). In a previous study, it was shown that the presence of cADPR decreased the threshold of caffeine or ryanodine required for  $\text{Ca}^{2+}$  release in rat pancreatic acinar cells by the same extent as that in the case of FK506 (28). This finding suggests that cADPR modulates the RyR in rat pancreatic acinar cells by the same mechanism as that by which FK506 modulates the RyR.

The FK506 ( $\leq 10\ \mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release from pancreatic microsomal vesicles was stimulated by the presence of cADPR (see Results). If FKBP is removed from the RyR by cADPR before the addition of FK506, it is unlikely that the FK506-induced  $^{45}\text{Ca}^{2+}$  release following pretreatment of the vesicles with cADPR is due to the dissociation of FKBP. FK506 is a compound with a macrocyclic lactone ring structure. It has been shown that rapamycin and ivermectin, macrocyclic lactone derivatives, increased the open probability of FKBP12-stripped RyRs in skeletal muscle (35,36). This finding suggests that the compounds activate the RyR by a mechanism other than dissociation of FKBP. The first phase of the FK506-induced  $^{45}\text{Ca}^{2+}$  release from the vesicles shown in Fig. 1 may be explained by a direct activation of the RyR by FK506.

In the rat cerebellum, FK506 dissociates FKBP12 from the  $\text{IP}_3\text{R}$  and decreases the threshold of  $\text{IP}_3$  required for  $\text{Ca}^{2+}$  release (18). The  $\text{EC}_{50}$  value for the dissociation of FKBP12 from the  $\text{IP}_3\text{R}$  complex has been shown to be 10-100 nM FK506 (23). This range is even lower than the  $K_m$  value ( $55\ \mu\text{M}$ ) for the second phase of the FK506-induced  $^{45}\text{Ca}^{2+}$  release from pancreatic microsomal vesicles. The second phase of the FK506-induced  $^{45}\text{Ca}^{2+}$  release may be due to a direct activation of the  $\text{IP}_3\text{R}$ , since heparin inhibited the FK506 ( $30\ \mu\text{M}$ )-induced  $^{45}\text{Ca}^{2+}$  release from the vesicles (Fig. 2) and the  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  pool of the vesicles was emptied by the addition of FK506 ( $100\ \mu\text{M}$ ) (Fig. 3). The finding that the second phase of the FK506-induced  $^{45}\text{Ca}^{2+}$  release from the vesicles represents a sigmoid curve (Fig. 1C) suggests allosteric changes in the  $\text{IP}_3\text{R}$  by the binding of FK506. There is a possibility that the allosteric effect induces greater release from the receptor by FK506 [the maximum value (60%) of the second phase of the FK506-induced  $^{45}\text{Ca}^{2+}$  release vs. the maximum percentage ( $\sim 25\%$ ) of the  $\text{IP}_3$ -induced  $^{45}\text{Ca}^{2+}$  release (37)].

Further studies using an FKBP antibody are needed to clarify whether the  $\text{Ca}^{2+}$  release from the ER induced by FK506 in rat pancreatic acinar cells is due to mechanisms without dissociation of the FKBP.

## Acknowledgments

The authors are grateful to Fujisawa Pharmaceutical Co., Ltd. for providing FK506.

## References

- Guillemette G, Balla T, Baukal AJ, Spät A and Catt KJ: Intracellular receptors for inositol 1,4,5-trisphosphate in angiotensin II target tissues. *J Biol Chem* 262: 1010-1015, 1987.
- Worley PF, Baraban JM, Supattapone S, Wilson VS and Snyder SH: Characterization of inositol trisphosphate receptor binding in brain: regulation by pH and calcium. *J Biol Chem* 262: 12132-12136, 1987.
- Supattapone S, Worley PF, Baraban JM and Snyder SH: Solubilization, purification, and characterization of an inositol trisphosphate receptor. *J Biol Chem* 263: 1530-1534, 1988.
- Chadwick CC, Saito A and Fleischer S: Isolation and characterization of the inositol trisphosphate receptor from smooth muscle. *Proc Natl Acad Sci USA* 87: 2132-2136, 1990.
- Furuichi T, Yoshikawa S, Miyawaki A, Wada K, Maeda N and Mikoshiba K: Primary structure and functional expression of the inositol 1,4,5-trisphosphate-binding protein  $\text{P}_{400}$ . *Nature* 342: 32-38, 1989.
- Mignery GA, Südhof TC, Takei K and De Camilli P: Putative receptor for inositol 1,4,5-trisphosphate similar to ryanodine receptor. *Nature* 342: 192-195, 1989.
- Inui M, Saito A and Fleischer S: Purification of the ryanodine receptor and identity with feet structures of junctional terminal cisternae of sarcoplasmic reticulum from fast skeletal muscle. *J Biol Chem* 262: 1740-1747, 1987.
- Inui M, Saito A and Fleischer S: Isolation of the ryanodine receptor from cardiac sarcoplasmic reticulum and identity with the feet structures. *J Biol Chem* 262: 15637-15642, 1987.
- Takeshima H, Nishimura S, Matsumoto T, Ishida H, Kangawa K, Minamino N, Matsuo H, Ueda M, Hanaoka M, Hirose T and Numa S: Primary structure and expression from complementary DNA of skeletal muscle ryanodine receptor. *Nature* 339: 439-445, 1989.
- Otsu K, Willard HF, Khanna VK, Zorzato F, Green NM and MacLennan DH: Molecular cloning of cDNA encoding the  $\text{Ca}^{2+}$  release channel (ryanodine receptor) of rabbit cardiac muscle sarcoplasmic reticulum. *J Biol Chem* 265: 13472-13483, 1990.
- Shoshan-Barmatz V, Pressley TA, Higham S and Kraus-Friedmann N: Characterization of high-affinity ryanodine-binding sites of rat liver endoplasmic reticulum: differences between liver and skeletal muscle. *Biochem J* 276: 41-46, 1991.
- Lilly LB and Gollan JL: Ryanodine-induced calcium release from hepatic microsomes and permeabilized hepatocytes. *Am J Physiol* 268: G1017-G1024, 1995.
- Ozawa T and Nishiyama A: Characterization of ryanodine-sensitive  $\text{Ca}^{2+}$  release from microsomal vesicles of rat parotid acinar cells: regulation by cyclic ADP-ribose. *J Membr Biol* 156: 231-239, 1997.
- Ozawa T: Ryanodine-sensitive  $\text{Ca}^{2+}$  release mechanism of rat pancreatic acinar cells is modulated by calmodulin. *Biochim Biophys Acta* 1452: 254-262, 1999.
- Ozawa T: Ryanodine-sensitive  $\text{Ca}^{2+}$  release mechanism in non-excitable cells (review). *Int J Mol Med* 7: 21-25, 2001.
- Timmerman AP, Ogunbumni E, Freund E, Wiederrecht G, Marks AR and Fleischer S: The calcium release channel of sarcoplasmic reticulum is modulated by FK-506-binding protein: dissociation and reconstitution of FKBP-12 to the calcium release channel of skeletal muscle sarcoplasmic reticulum. *J Biol Chem* 268: 22992-22999, 1993.
- Brillantes A-MB, Ondrias K, Scott A, Kobrinsky E, Ondriasová E, Moschella MC, Jayaraman T, Landers M, Ehrlich BE and Marks AR: Stabilization of calcium release channel (ryanodine receptor) function by FK506-binding protein. *Cell* 77: 513-523, 1994.
- Cameron AM, Steiner JP, Sabatini DM, Kaplin AI, Walensky LD and Snyder SH: Immunophilin FK506 binding protein associated with inositol 1,4,5-trisphosphate receptor modulates calcium flux. *Proc Natl Acad Sci USA* 92: 1784-1788, 1995.
- Collins JH: Sequence analysis of the ryanodine receptor: possible association with a 12K, FK506-binding immunophilin/protein kinase C inhibitor. *Biochem Biophys Res Commun* 178: 1288-1290, 1991.
- Jayaraman T, Brillantes A-M, Timmerman AP, Fleischer S, Erdjument-Bromage H, Tempst P and Marks AR: FK506 binding protein associated with the calcium release channel (ryanodine receptor). *J Biol Chem* 267: 9474-9477, 1992.
- Lam E, Martin MM, Timmerman AP, Sabers C, Fleischer S, Lukas T, Abraham RT, O'Keefe SJ, O'Neill EA and Wiederrecht GJ: A novel FK506 binding protein can mediate the immuno-

- suppressive effects of FK506 and is associated with the cardiac ryanodine receptor. *J Biol Chem* 270: 26511-26522, 1995.
22. Timerman AP, Onoue H, Xin HB, Barg S, Copello J, Wiederrecht G and Fleischer S: Selective binding of FKBP12.6 by the cardiac ryanodine receptor. *J Biol Chem* 271: 20385-20391, 1996.
  23. Cameron AM, Steiner JP, Roskams AJ, Ali SM, Ronnett GV and Snyder SH: Calcineurin associated with the inositol 1,4,5-trisphosphate receptor-FKBP12 complex modulates  $\text{Ca}^{2+}$  flux. *Cell* 83: 463-472, 1995.
  24. Cameron AM, Nucifora FC, Fung ET, Livingston DJ, Aldape RA, Ross CA and Snyder SH: FKBP12 binds the inositol 1,4,5-trisphosphate receptor at leucine-proline (1400-1401) and anchors calcineurin to this FK506-like domain. *J Biol Chem* 272: 27582-27588, 1997.
  25. Streb H, Irvine RF, Berridge MJ and Schulz I: Release of  $\text{Ca}^{2+}$  from a nonmitochondrial intracellular store in pancreatic acinar cells by inositol-1,4,5-trisphosphate. *Nature* 306: 67-69, 1983.
  26. Streb H, Bayerdörffer E, Haase W, Irvine RF and Schulz I: Effect of inositol-1,4,5-trisphosphate on isolated subcellular fractions of rat pancreas. *J Membr Biol* 81: 241-253, 1984.
  27. Dehlinger-Kremer M, Zeuzem S and Schulz I: Interaction of caffeine-,  $\text{IP}_3$ - and vanadate-sensitive  $\text{Ca}^{2+}$  pools in acinar cells of the exocrine pancreas. *J Membr Biol* 119: 85-100, 1991.
  28. Ozawa T: Elucidation of the ryanodine-sensitive  $\text{Ca}^{2+}$  release mechanism of rat pancreatic acinar cells: modulation by cyclic ADP-ribose and FK506. *Biochim Biophys Acta* 1693: 159-166, 2004.
  29. Ozawa T, Thévenod F and Schulz I: Characterization of two different  $\text{Ca}^{2+}$  uptake and  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  release mechanisms in microsomal  $\text{Ca}^{2+}$  pools of rat pancreatic acinar cells. *J Membr Biol* 144: 111-120, 1995.
  30. Bradford MM: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248-254, 1976.
  31. Lee HC: Potentiation of calcium- and caffeine-induced calcium release by cyclic ADP-ribose. *J Biol Chem* 268: 293-299, 1993.
  32. Ghosh TK, Eis PS, Mullaney JM, Ebert CL and Gill DL: Competitive, reversible, and potent antagonism of inositol 1,4,5-trisphosphate-activated calcium release by heparin. *J Biol Chem* 263: 11075-11079, 1988.
  33. Noguchi N, Takasawa S, Nata K, Tohgo A, Kato I, Ikehata F, Yonekura H and Okamoto H: Cyclic ADP-ribose binds to FK506-binding protein 12.6 to release  $\text{Ca}^{2+}$  from islet microsomes. *J Biol Chem* 272: 3133-3136, 1997.
  34. Tang W-X, Chen Y-F, Zou A-P, Campbell WB and Li P-L: Role of FKBP12.6 in cADPR-induced activation of reconstituted ryanodine receptors from arterial smooth muscle. *Am J Physiol* 282: H1304-H1310, 2002.
  35. Ahern GP, Junankar PR and Dulhunty AF: Ryanodine receptors from rabbit skeletal muscle are reversibly activated by rapamycin. *Neurosci Lett* 225: 81-84, 1997.
  36. Ahern GP, Junankar PR, Pace SM, Curtis S, Mould JA and Dulhunty AF: Effects of ivermectin and midcamycin on ryanodine receptors and the  $\text{Ca}^{2+}$ -ATPase in sarcoplasmic reticulum of rabbit and rat skeletal muscle. *J Physiol* 514: 313-326, 1999.
  37. Ozawa T and Schulz I:  $\text{H}^+$  uptake increases GTP-induced connection of inositol 1,4,5-trisphosphate and caffeine-sensitive calcium pools in pancreatic microsomal vesicles. *Biochem Biophys Res Commun* 180: 755-764, 1991.