

# The novel anti-adipogenic effect and mechanisms of action of SGI-1776, a Pim-specific inhibitor, in 3T3-L1 adipocytes

YU-KYOUNG PARK<sup>1\*</sup>, VICTOR SUKBONG HONG<sup>2\*</sup>, TAE-YOON LEE<sup>3\*</sup>, JINHO LEE<sup>2</sup>, JONG-SOON CHOI<sup>4,5</sup>,  
DONG-SOON PARK<sup>6</sup>, GI-YOUNG PARK<sup>6</sup> and BYEONG-CHURL JANG<sup>1</sup>

<sup>1</sup>Department of Molecular Medicine, College of Medicine, <sup>2</sup>Department of Chemistry, College of Natural Sciences, Keimyung University, Daegu 704-701; <sup>3</sup>Department of Microbiology, College of Medicine, Yeungnam University, Daegu 705-717; <sup>4</sup>Division of Life Science, Korea Basic Science Institute, Daejeon 305-333; <sup>5</sup>Graduate School of Analytical Science and Technology, Chungnam National University, Daejeon 305-764; <sup>6</sup>Department of Rehabilitation Medicine, Catholic University of Daegu School of Medicine, Daegu 705-718, Republic of Korea

Received June 8, 2015; Accepted October 27, 2015

DOI: 10.3892/ijmm.2015.2415

**Abstract.** The proviral integration site for moloney murine leukemia virus (Pim) kinases, consisting of Pim-1, Pim-2 and Pim-3, belongs to a family of serine/threonine kinases that are involved in controlling cell growth and differentiation. Pim kinases are emerging as important mediators of adipocyte differentiation. SGI-1776, an inhibitor of Pim kinases, is widely used to assess the physiological roles of Pim kinases, particularly cell functions. In the present study, we examined the effects of SGI-1776 on adipogenesis. The anti-adipogenic effect of SGI-1776 was measured by Oil Red O staining and AdipoRed assays. The effect of SGI-1776 on the growth of 3T3-L1 adipocytes was determined by cell count analysis. The effects of SGI-1776 on the protein and mRNA expression of adipogenesis-related proteins and adipokines in 3T3-L1 adipocytes were also evaluated by western blot analysis and RT-PCR, respectively. Notably, SGI-1776 markedly inhibited lipid accumulation during the differentiation of 3T3-L1 preadipocytes into adipocytes. On a mechanistic level, SGI-1776 inhibited not only the expression of CCAAT/enhancer-binding protein- $\alpha$  (C/EBP- $\alpha$ ), peroxisome proliferator-activated receptor- $\gamma$  (PPAR- $\gamma$ ) and fatty

acid synthase (FAS), but also the phosphorylation of signal transducer and activator of transcription-3 (STAT-3). Moreover, SGI-1776 decreased the expression of adipokines, including the expression of leptin and regulated on activation, normal T cell expressed and secreted (RANTES) during adipocyte differentiation. These findings demonstrate that SGI-1776 inhibits adipogenesis by downregulating the expression and/or phosphorylation levels of C/EBP- $\alpha$ , PPAR- $\gamma$ , FAS and STAT-3.

## Introduction

Obesity is a high risk factor for the development of a number of human diseases, including insulin resistance, type 2 diabetes, hyperlipidemia and cancer (1). Obesity is induced by several factors, such as genetic and endocrine abnormalities, certain medicines, a low metabolic rate, nutritional and environmental factors, as well as imbalanced energy homeostasis (2,3). Previous studies have indicated that adipose tissue serves as an energy reservoir and also plays a critical role in the control of energy metabolism by secreting adipokines (3,4). However, there is also strong evidence that the abnormal expansion/accumulation of adipose tissue, which is largely attributable to excessive adipocyte differentiation and an increase in the number and size of fat cells, is closely linked to the development of obesity (5,6). Thus, any compound that inhibits excessive adipocyte differentiation and adipocyte hyperplasia/hypertrophy may have preventive and therapeutic potential against obesity.

It has previously been reported that the differentiation of preadipocytes into mature adipocytes, also termed adipogenesis, is controlled by numerous cellular proteins, transcription factors, adipocyte-specific genes, lipogenic enzymes and signaling proteins. For instance, it has previously been demonstrated that the expression of the family of CCAAT/enhancer-binding proteins (C/EBPs; C/EBP- $\alpha$ , - $\beta$  and - $\delta$ ) and peroxisome proliferator-activated receptors (PPARs; PPAR- $\gamma$ , - $\alpha$  and - $\beta$ ) is critical for adipocyte differentiation (7,8). Moreover, there is evidence indicating that Janus-activated protein kinase-2 (JAK-2)/signal transducer

---

*Correspondence to:* Professor Byeong-Churl Jang, Department of Molecular Medicine, College of Medicine, Keimyung University, 1095 Dalgubeoldae-ro, Dalseo-gu, Daegu 704-701, Republic of Korea  
E-mail: jangbc123@gw.kmu.ac.kr

Professor Gi-Young Park, Department of Rehabilitation Medicine, Catholic University of Daegu School of Medicine, 33 Duryugongwon-ro 17-gil, Nam-Gu, Daegu 705-718, Republic of Korea  
E-mail: gypark@cu.ac.kr

\*Contributed equally

**Key words:** SGI-1776, adipogenesis, peroxisome proliferator-activated receptor- $\gamma$ , CCAAT/enhancer-binding protein- $\alpha$ , signal transducer and activator of transcription-3, fatty acid synthase

and activator of transcription (STAT)-3 and STAT-5 signaling complexes are associated with adipocyte differentiation (9,10). In addition, a number of signaling proteins and factors, including protein kinase A (PKA), protein kinase C (PKC), extracellular signal-regulated protein kinase-1/2 (ERK-1/2) and adenosine 3',5'-cyclic monophosphate (cAMP), have been found to be of importance for controlling adipocyte differentiation (11-13).

SGI-1776, *N*-[(1-methylpiperidin-4-yl)methyl]-3-[3-(trifluoromethoxy)phenyl]imidazo[1,2-*b*]pyridazin-6-amine, has been shown to inhibit three members of the Pim kinase family (Pim-1, Pim-2 and Pim-3) (14). Pim kinases are constitutively active serine/threonine kinases that are known to be over-expressed in hematological malignancies, such as acute myeloid leukemia and multiple myeloma (15). Pim kinases have multiple substrates that are involved in transcription, protein translation, cell proliferation and apoptosis. Due to its ability to inhibit Pim kinase, SGI-1776 has been tested as a possible treatment for a number of hematological malignancies (16). Of note, it has been previously suggested that Pim-2 is substantially expressed in adipocytes (17) and the expression of Pim-1 in adipose tissue may be used as a marker of adipocytic differentiation (18). There is also compelling evidence indicating that Pim kinases are downstream target genes of STAT-3 (19). In a recent studies of ours, we demonstrated that a meridianin C derivative, which inhibits protein kinases, including Pim kinases (20), markedly inhibited adipogenesis (21). This research led us to further hypothesize that SGI-1776, a Pim-specific inhibitor would suppress adipogenesis to a greater extent than a meridianin C derivative.

In the present study, we investigated the anti-adipogenic effects of SGI-1776 on 3T3-L1 adipocytes. To the best of our knowledge, this is the first study which demonstrates that SGI-1776 exerts a prominent anti-adipogenic effect without affecting the viability of adipocytes and that the anti-adipogenic effect of SGI-1776 is largely associated with the decreased expression and/or phosphorylation levels of C/EBP- $\alpha$ , PPAR- $\gamma$ , fatty acid synthase (FAS) and STAT-3.

## Materials and methods

**Materials.** Polyclonal C/EBP- $\alpha$  (sc-61), monoclonal PPAR- $\gamma$  (sc-7273), monoclonal STAT-3 (sc-8019), monoclonal phosphorylated (p)-STAT-3 (p-STAT-3; sc-8059), polyclonal STAT-5 (sc-835) and polyclonal p-STAT-5 (sc-101806) antibodies were all purchased from Santa Cruz Biotechnology, Inc. (Delaware, CA, USA). Monoclonal FAS (610962) antibody was purchased from BD Biosciences (San Jose, CA, USA). Monoclonal  $\beta$ -actin (A5441) antibody, 3-isobutyl-1-methylxanthine (IBMX), dexamethasone and insulin were all purchased from Sigma (St. Louis, MO, USA). SGI-1776 was purchased from Merck Millipore (Darmstadt, Germany).

**Cell culture and differentiation.** As previously described (27), 3T3-L1 preadipocytes (ATCC, Manassas, VA, USA) were grown to the contact inhibition stage and remained in the post-confluent stage for 2 days in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS; Gibco, Grand Island, NY, USA) and penicillin-streptomycin (Welgene, Daegu, Korea). Differentiation was then induced by changing the medium to DMEM supplemented

with 10% FBS (Welgene, Daegu, Korea) plus a cocktail of hormones (MDI), containing 0.5 mM IBMX (M), 0.5  $\mu$ M dexamethasone (D) and 5  $\mu$ g/ml insulin (I) in the presence or absence of SGI-1776 or butein (Sigma) at the indicated concentrations. Following 48 h MDI induction, the differentiation medium was replaced with DMEM supplemented with 10% FBS and 5  $\mu$ g/ml insulin in the presence or absence of SGI-1776 at the indicated concentrations. The cells were then fed every other day with DMEM containing 10% FBS in the presence or absence of SGI-1776 or butein at the indicated concentrations until day 8. On day 8, the preadipocytes became mature adipocytes that were rounded and filled with many oil droplets.

**Oil Red O staining.** On day 8 of differentiation, the mock- (not treated with SGI-1776 or butein) or SGI-1776- or butein-treated 3T3-L1 cells were washed twice with phosphate-buffered saline (PBS), fixed with 10% formaldehyde for 2 h at room temperature, washed with 60% isopropanol and dried completely. The fixed cells were then stained with Oil Red O (Sigma, St. Louis, MO, USA) working solution for 1 h at room temperature and were then washed twice with distilled water. Lipid droplets were observed under a light microscope (TS100; Nikon, Tokyo, Japan).

**Cell count assay.** The 3T3-L1 preadipocytes seeded in 24-well plates were similarly grown under the above-mentioned differentiation conditions. On day 8 of differentiation, the mock- or SGI-1776-treated 3T3-L1 cells, which cannot be stained with trypan blue dye, were counted under a microscope. The cell count assay was carried out in triplicate. Data are presented as the means  $\pm$  standard error of 3 independent experiments.

**Quantification of intracellular triglyceride (TG) content by AdipoRed assay.** On day 8 of differentiation, the lipid content in the mock- or SGI-1776- or butein-treated 3T3-L1 cells was measured using a commercially available AdipoRed Assay Reagent kit according to the manufacturer's instructions (Lonza, Basel, Switzerland). Following 10 min of incubation, fluorescence was measured using a multilabel reader (Victor3; PerkinElmer, Waltham, MA, USA) with the excitation set at 485 nm and the emission at 572 nm.

**Preparation of whole cell lysates.** As previously described (27), at a designated time point, the 3T3-L1 cells were washed twice with PBS and exposed to modified RIPA buffer [50 mM Tris-Cl (pH 7.4), 150 mM NaCl, 0.1% sodium dodecyl sulfate, 0.25% sodium deoxycholate, 1% Triton X-100, 1% Nonidet P-40, 1 mM EDTA, 1 mM EGTA, proteinase inhibitor cocktail (1X)]. The cell lysates were collected and centrifuged at 12,000 rpm for 20 min at 4°C. The supernatant was saved, and the protein concentrations were determined using Bradford reagent (Bio-Rad, Hercules, CA, USA).

**Western blot analysis.** Proteins (50  $\mu$ g) were separated by SDS-PAGE (10%) and transferred onto nitrocellulose membranes (Millipore, Bedford, MA, USA). The membranes were washed with Tris-buffered saline (10 mM Tris-Cl, 150 mM NaCl, pH 7.5), supplemented with 0.05% (v/v) Tween-20 (TBST), followed by blocking with TBST containing 5% (w/v) non-fat dried milk. The membranes were incubated

overnight with antibodies specific for C/EBP- $\alpha$ , PPAR- $\gamma$ , STAT-3, p-STAT-3, STAT-5, p-STAT-5, FAS or  $\beta$ -actin at 4°C. The membranes were then exposed to secondary antibodies conjugated to horseradish peroxidase for 2 h at room temperature and further washed 3 times with TBST. Immunoreactivity was detected using ECL reagents. Equal protein loading was assessed by the expression levels of actin.

#### Reverse transcription-polymerase chain reaction (RT-PCR).

At a designated time point, total cellular RNA in the mock- or SGI-1776-treated 3T3-L1 cells was isolated using RNazol-B (Tel-Test, Inc., Friendswood, TX, USA). Three micrograms of total RNA were reverse transcribed using a random hexadeoxynucleotide primer and reverse transcriptase. Single-stranded cDNA was amplified by PCR using specific primers. The primer sequences used for amplification were as follows: C/EBP- $\alpha$  sense, 5'-TTACAACAGGCCAGGTTTCC-3' and antisense, 5'-CTCTGGGATGGATCGATTGT-3'; PPAR- $\gamma$  sense, 5'-GGT GAACTCTGGGAGATTC-3' and antisense, 5'-CAACCA TTGGGTCAGCTCTC-3'; FAS sense, 5'-TTGCTGGCAC TACAGAATGC-3' and antisense, 5'-AACAGCCTCAGAG CGACAAT-3'; leptin sense, 5'-CCAAAACCCATCAA GACC-3' and antisense, 5'-CTCAAAGCCACCACCTCTGT-3'; adiponectin sense, 5'-GGAGATGCAGGTCTTCTTGGT-3' and antisense, 5'-TCCTGATACTGGTCGTAGGTGAA-3'; regulated on activation, normal T cell expressed and secreted (RANTES) sense, 5'-TCCAATCTTGCAGTCGTGTTTG-3' and antisense, 5'-TCTGGGTTGGCACACACTTG-3'; monocyte chemoattractant protein-1 (MCP-1) sense, 5'-TCCAATCTT GCAGTCGTGTTTG-3' and antisense, 5'-TCTGGGTTG GCACACACTTG-3'; actin antisense, 5'-GGTAGGAACA CGGAAGGCCA-3'. The mRNA expression levels of actin were used as an internal control to evaluate the relative mRNA expression of adipocyte-specific genes and adipokines.

**Statistical analysis.** Cell count analysis was completed in triplicate and repeated 3 times. Data are expressed as the means  $\pm$  standard error (SE). The significant differences between groups were determined by one-way ANOVA (Laerd Statistics, Chicago, IL, USA). All significance testing was based upon a P-value  $<0.05$ , indicating a statistically significant difference.

## Results

**SGI-1776 exerts potent anti-adipogenic effects.** Primarily, we wished to determine whether the Pim-selective inhibitor, SGI-1776, inhibits intracellular lipid accumulation during the differentiation of 3T3-L1 preadipocytes into adipocytes by Oil Red O staining. The timescale of 3T3-L1 preadipocyte differentiation is illustrated in Fig. 1A. Many lipid droplets were formed in the differentiated 3T3-L1 preadipocytes cultured in the induction medium on day 8 of cell differentiation, compared with the undifferentiated cells (preadipocytes), in which no lipid droplets had formed (Fig. 1B, upper panels). However, treatment with SGI-1776 for 8 days markedly reduced the amount of lipid droplets in a concentration-dependent manner, as compared with the mock-treated cells. Evidently, treatment with 10  $\mu$ M SGI-1776 completely blocked the accumulation of lipid droplets. The SGI-1776-mediated lipid-reducing effect was also confirmed by light microscopy (Fig. 1B, lower panels).

In addition, we noted that SGI-1776 decreased the levels of TG during adipocyte differentiation in a dose-dependent manner, with maximum reduction being achieved with an inhibitor concentration of 10  $\mu$ M (Fig. 1C). However, using a cell count assay, it was observed that SGI-1776 induced a slight decrease in viability at 10  $\mu$ M, but not at 5  $\mu$ M (Fig. 1D). Certain compounds, such as butein (40  $\mu$ M), have also previously been shown to exert anti-adipogenic effects on 3T3-L1 adipocytes (22). In this study, we thus compared the anti-adipogenic effect of SGI-1776 (5  $\mu$ M) with that of butein (40  $\mu$ M) used as a positive control. In agreement with the results of previous studies, treatment with butein at 40  $\mu$ M markedly inhibited both lipid droplet formation (Fig. 1E) and TG synthesis (Fig. 1F) during adipocyte differentiation. Of note, treatment with SGI-1776 at 5  $\mu$ M exerted more potent inhibitory effects on both lipid droplet formation and TG synthesis than butein (40  $\mu$ M). Therefore, the concentration of 5  $\mu$ M of SGI-1776 was selected for use in further experiments due to the potent inhibitory effects it exerted on both lipid accumulation and TG synthesis without affecting the number of adipocytes.

**Inhibition of the early adipogenic process is critical for the SGI-1776-mediated anti-adipogenic effects.** We subsequently sought to determine which stage(s) of adipocyte differentiation is inhibited by SGI-1776. To this end, the 3T3-L1 preadipocytes were incubated with induction medium (MDI, insulin and FBS) in the presence or absence of SGI-1776 (5  $\mu$ M) for the indicated periods of time, as illustrated in Fig. 2A. As expected, the mock-treated cells incubated with MDI, insulin and FBS for 2, 5 and 8 days exhibited high lipid accumulation (treatment a). The 3T3-L1 preadipocytes treated with SGI-1776 from day 0 to 2 (treatment b), or from day 0 to 5 (treatment c) showed a comparatively low lipid content. The preadipocytes treated with SGI-1776 from day 0 to 8 exhibited the highest inhibition of lipid droplet accumulation (treatment d). Of note, differentiating 3T3-L1 cells treated with SGI-1776 from day 2 to 5, from day 2 to 8 or from day 5 to 8 (treatment e, f and g) exhibited a much weaker inhibition of adipogenic differentiation, compared with treatment b, c and d (Fig. 2B). The results of the AdipoRed assay further demonstrated that treatment b, c and d resulted in the highest reduction of TG during adipocyte differentiation (Fig. 2C). Treatment f also substantially reduced the TG content during adipocyte differentiation, whereas treatment e and g led to a slight reduction in the TG content. These results collectively suggest that SGI-1776 inhibits adipogenesis particularly at an early phase of differentiation.

**SGI-1776 decreases the expression of C/EBP- $\alpha$  and PPAR- $\gamma$  and the phosphorylation levels of STAT-3 during adipocyte differentiation.** To elucidate the mechanisms underlying the SGI-1776-mediated anti-adipogenic effect, we examined the effects of SGI-1776 (5  $\mu$ M) on the expression and/or activity (phosphorylation) levels of adipogenic transcription factors, including the families of C/EBPs, PPARs and STATs. Western blot analysis revealed that SGI-1776 markedly inhibited the adipogenesis-dependent protein expression of C/EBP- $\alpha$  and PPAR- $\gamma$  during adipocyte differentiation, particularly on day 5 and 8 (Fig. 3A and E). The results from RT-PCR also revealed that SGI-1776 suppressed the mRNA expression of C/EBP- $\alpha$  and PPAR- $\gamma$  during adipocyte differentiation (Fig. 3B and F).




Figure 1. Effects of SGI-1776 on lipid accumulation, triglyceride (TG) synthesis and cell growth during adipocyte differentiation. (A) The experimental schema of 3T3-L1 preadipocyte differentiation. (B) Microscopy of lipid droplets of 3T3-L1 preadipocytes or differentiated adipocytes on day 8 visualized by Oil Red O staining. Phase-contrast images of the cells were also acquired following treatment (B, lower panels). Each image in (B) is representative of 3 independent experiments. (C) Quantification of the TG content in SGI-1776-treated 3T3-L1 adipocytes on day 8 by AdipoRed assay. Values represent the means  $\pm$  SE of data from 3 independent experiments with 3 replicates. \* $P < 0.05$  vs. control (mock, no SGI-1776 treatment). (D) Measurement of cell proliferation in SGI-1776-treated 3T3-L1 adipocytes on day 8 by cell count assay. 3T3-L1 preadipocytes were grown under the above-mentioned 3T3-L1 preadipocyte differentiation conditions as in (C). On day 8, SGI-1776-treated 3T3-L1 cells, which cannot be stained with trypan blue dye, were counted under a microscope. The cell count assay was undertaken in triplicate. Data represent the means  $\pm$  SE of 3 independent experiments. \* $P < 0.05$  vs. control (no chemical). (E) Microscopy of lipid droplets of the control, SGI-1776- or butein-treated 3T3-L1 adipocytes on day 8 by Oil Red O staining. Phase-contrast images of the cells were also acquired following treatment (E, lower panels). Each image in (E) is representative of 3 independent experiments. (F) Quantification of TG content in the control, SGI-1776- or butein-treated 3T3-L1 adipocytes on day 8 by AdipoRed assay. Values represent the means  $\pm$  SE of data from 3 independent experiments with 3 replicates. \* $P < 0.05$  vs. control (no treatment).

These results suggest that SGI-1776 downregulates PPAR- $\gamma$  and C/EBP- $\alpha$  expression at the transcriptional level. Moreover, SGI-1776 markedly decreased the phosphorylation levels of STAT-3 on day 2 during adipocyte differentiation (Fig. 3C

and G). By contrast, SGI-1776 did not markedly affect the phosphorylation levels of STAT-5. Notably, SGI-1776 also had the capacity to suppress STAT-3 phosphorylation during the early stages of adipogenesis induced by MDI at the treatment




Figure 2. Effect of SGI-1776 on different processes of 3T3-L1 adipocyte differentiation. (A) Schematic outline of the experimental setup: the double-headed arrows indicate the length of treatment. 3T3-L1 cells were treated with or without SGI-1776 for the indicated periods of time. (B) 3T3-L1 preadipocytes were induced to differentiate with induction medium containing MDI, insulin and fetal bovine serum (FBS) in the presence or absence of SGI-1776, as described in Materials and methods (a-g) at the indicated time points. The cellular lipid content at each time-point was assessed by Oil Red O staining (left panels). Phase-contrast images of the cells were also acquired following treatment (right panels). Each images is representative of 3 independent experiments. (C) Quantification of triglyceride (TG) content in control or SGI-1776-treated 3T3-L1 adipocytes on day 8 by AdipoRed assay. Values represent the means  $\pm$  SE of data from 3 independent experiments with 3 replicates. \* $P < 0.05$  vs. control (no treatment).

times of 2, 4, 8 and 24 h (Fig. 3D and H). SGI-1776 did not markedly alter the total protein levels of STAT-3 and STAT-5 during adipocyte differentiation at the designated time points.

*SGI-1776 downregulates the protein and/or mRNA expression of FAS, leptin and RANTES during adipocyte differentiation.* The expression and secretion of adipokines occur during preadipocyte differentiation (28-30). Therefore, we examined whether SGI-1776 (5  $\mu$ M) regulates the expression of adipocyte-specific genes and/or adipokines during adipocyte differentiation. SGI-1776 markedly suppressed the protein and mRNA expression levels of FAS on days 5 and 8 during adipocyte differentiation (Fig. 4A-C). Furthermore, the results from RT-PCR demonstrated that SGI-1776 markedly reduced the insulin- and FBS-induced mRNA expression of leptin and RANTES on days 5 and 8 (Fig. 4B-D). However, SGI-1776 did not greatly affect the mRNA levels of adiponectin or MCP-1 on days 5 and 8 (Fig. 4B and D).

## Discussion

Excessive adipocyte differentiation causes the abnormal expansion/accumulation of adipose tissue, leading to the high

secretion of adipokines, which have been implicated in inflammation, insulin resistance and metabolic disorders (5,6). Thus, inhibitors of adipocyte differentiation likely have therapeutic potential as anti-obesity drugs. Recently, we demonstrated that a meridianin C derivative, which inhibits protein kinases including Pim kinases (20), markedly blocks adipocytic differentiation (21). In the present study, we focused on the role of Pim kinases in 3T3-L1 adipogenesis. We noted that when the 3T3-L1 cells differentiated in the presence of a potent and selective Pim inhibitor, such as SGI-1776, adipogenesis was severely inhibited.

Mature adipocytes have a spherical shape and are filled with a high number of intracellular lipid droplets, and these can be distinguished from fibroblast-like preadipocytes (23). The present study demonstrated that SGI-1776 (5  $\mu$ M) markedly inhibited lipid accumulation during the differentiation of 3T3-L1 preadipocytes into adipocytes by 8 days of differentiation (Fig. 1B and C), suggesting that it exerts a potent anti-adipogenic effect. Cell count analysis further revealed that treatment with SGI-1776 (5  $\mu$ M) for 8 days did not markedly affect the viability of 3T3-L1 adipocytes (Fig. 1D).

As previously mentioned, the induction of adipocyte differentiation is largely dependent on the expression and activity of




Figure 3. Effect of SGI-1776 on the cellular expression and/or phosphorylation levels of CCAAT/enhancer-binding protein- $\alpha$  (C/EBP- $\alpha$ ), peroxisome proliferator-activated receptor- $\gamma$  (PPAR- $\gamma$ ), signal transducer and activator of transcription (STAT)-3 and STAT-5 during 3T3-L1 adipocyte differentiation. (A-C) 3T3-L1 preadipocytes were induced to differentiate with induction medium containing MDI, insulin and fetal bovine serum (FBS) in the presence or absence of SGI-1776, and harvested on days 2, 5 and 8, respectively. Cellular protein and mRNA at the indicated time points were extracted and analyzed by (A and C) western blot analysis and (B) RT-PCR. Each blot in (A-C) is representative of 3 independent experiments. (D) 3T3-L1 preadipocytes were treated with induction medium containing MDI in the presence or absence of SGI-1776 and harvested at 2, 4, 8 and 24 h, respectively. The cellular protein at the indicated time points was extracted and analyzed by western blot analysis. Each blot in (D) is representative of 3 independent experiments. (E and F) The densitometric data of (A and B), respectively, that show C/EBP- $\alpha$  and PPAR- $\gamma$  protein and mRNA levels normalized to actin protein and mRNA levels as percentages of the value in the presence or absence of SGI-1776 at the indicated days. Values represent the means  $\pm$  SE of data from 3 independent experiments with 3 replicates. \* $p < 0.05$  compared to the value of SGI-1776-free control on the indicated days. (G and H) The densitometric data of (C and D), respectively, that show p-STAT-3, p-STAT-1 and p-STAT-5 protein levels normalized to total expression levels of each protein as percentages of the value in the presence or absence of SGI-1776 at the indicated days. \* $p < 0.05$  compared to the value of SGI-1776-free control on the indicated days.

adipogenesis-related transcription factors, such as C/EBP- $\alpha$  and PPAR- $\gamma$ . It has been previously demonstrated that blocking their expression and/or activity with either pharmacological inhibitors or small interfering RNA leads to the inhibition of lipid accumulation during adipocyte differentiation, and that

the knockdown of C/EBP- $\alpha$  or PPAR- $\gamma$  decreases or impairs the differentiation of white adipose tissue in mice (7,8,24). The present study demonstrated that SGI-1776 inhibited the protein and mRNA expression of C/EBP- $\alpha$  and PPAR- $\gamma$  during adipocyte differentiation (Fig. 3A, B, E and F). Thus,




Figure 4. Effect of SGI-1776 on fatty acid synthase (FAS) protein and mRNA expression levels of FAS and adipokines [leptin, adiponectin, regulated on activation, normal T cell expressed and secreted (RANTES) and monocyte chemoattractant protein-1 (MCP-1)] during 3T3-L1 adipocyte differentiation. (A and B) 3T3-L1 preadipocytes were induced to differentiate with induction medium containing MDI, insulin and fetal bovine serum (FBS) in the presence or absence of SGI-1776, and harvested at days 2, 5 and 8, respectively. Cellular protein and mRNA at the indicated time points were extracted and analyzed by (A) western blot analysis and (B) RT-PCR. Each blot in (A and B) is a representative of 3 independent experiments. (C and D) The densitometric data of (A and B), respectively, that show FAS protein and mRNA levels and/or leptin, adiponectin, RANTES and MCP-1 mRNA levels normalized to actin protein and/or mRNA levels as percentages of the value in the presence or absence of SGI-1776 on the indicated days. Values represent the means  $\pm$  SE of data from 3 independent experiments with 3 replicates. \* $p < 0.05$  compared to the value of SGI-1776-free control on the indicated days.

the SGI-1776-mediated anti-adipogenic effect appears to be closely associated with the reduced expression of C/EBP- $\alpha$  and PPAR- $\gamma$ . Previously, it has also been reported that the family of STATs, including STAT-1, STAT-3, STAT-5 and STAT-6, is expressed in both 3T3-L1 preadipocytes and adipocytes (9), and that several members of the STAT family, including STAT-3 and STAT-5, are critical for 3T3-L1 adipocyte differentiation (10,21,25-27). Of note, previous studies have further demonstrated that STAT-3 is rapidly (within hours) activated during adipocyte differentiation, and that active STAT-3 plays an important role in the early stages of adipogenesis (10,21,27). In the present study, we noted that SGI-1776 blocked not only the early adipogenic process (Fig. 2B and C), but also MDI (but neither insulin nor FBS) signaling to activate STAT-3 during the early stage of adipogenesis (Fig. 3C, D and H). It is thus likely that the STAT-3-dependent blockage of early adipogenesis is also crucial for the SGI-1776-mediated anti-adipogenic effect. Previous studies have indicated that STAT-3 regulates adipocyte differentiation via PPAR- $\gamma$ , and that STAT-3 activity is also important for the adipogenesis-induced expression of

C/EBP- $\alpha$  and PPAR- $\gamma$  (26,27). Hence, it is suggested in this study that SGI-1776 inhibits adipogenesis primarily through the inhibition of STAT-3 at the early adipogenic process, which subsequently downregulates the expression of C/EBP- $\alpha$  and PPAR- $\gamma$ .

Previous research has demonstrated that the increased expression and activity of C/EBPs and PPARs is necessary for the expression of adipocyte-specific genes and adipokines, including FAS, leptin, adiponectin, MCP-1 and RANTES (28-30). Of those, FAS is a lipogenic enzyme involved in fatty acid synthesis, and its expression is highly upregulated in cells or tissues with high rates of fatty acid synthesis (31). In addition, it has previously been shown that FAS mRNA and protein expression are substantially increased during 3T3-L1 preadipocyte differentiation (21,27,32). In this study, we noted that SGI-1776 also had the ability to markedly inhibit the insulin- and FBS-mediated induction of FAS at both the protein and mRNA level in 3T3-L1 adipocytes (Fig. 4). These results suggest that SGI-1776 exerts an anti-lipogenic effect by suppressing FAS expression, which further contributes to the SGI-1776-mediated anti-adipogenic

effect. It has been well documented that the maturation of adipocyte differentiation is characterized by the capacity of mature adipocytes to express and secrete adipokines, which are involved in the endocrine control of energy homeostasis (33,34). Therefore, considering that SGI-1776 inhibits the mRNA expression of leptin and RANTES, but not adiponectin, during the adipocyte differentiation process (Fig. 4B and D), we suggest that SGI-1776 further blocks the adipocyte differentiation maturation process.

In conclusion, in this study, we demonstrate firstly that a pan-Pim kinase inhibitor, SGI-1776, exerts potent anti-adipogenic effects in 3T3-L1 cells that are related to the reduced expression and/or phosphorylation levels of PPAR- $\gamma$ , C/EBP- $\alpha$ , FAS and STAT-3. The findings of this study thus suggest that the Pim-specific inhibitor, SGI-1776, be considered a potential therapeutic agent for the prevention and treatment of obesity in which excessive adipogenesis plays pathological roles.

### Acknowledgements

This study was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea Government (MSIP) (no. 2014R1A5A2010008) and also in part by the Korea Basic Science Institute grants (no. D35400) and a Keimyung University Research Grant of 2015.

### References

- Kopelman PG: Obesity as a medical problem. *Nature* 404: 635-643, 2000.
- Ahima RS and Flier JS: Adipose tissue as an endocrine organ. *Trends Endocrinol Metab* 11: 327-332, 2000.
- Havel PJ: Control of energy homeostasis and insulin action by adipocyte hormones: leptin, acylation stimulating protein, and adiponectin. *Curr Opin Lipidol* 13: 51-59, 2002.
- Spiegelman BM and Flier JS: Obesity and the regulation of energy balance. *Cell* 104: 531-543, 2001.
- Bays HE: 'Sick fat', metabolic disease, and atherosclerosis. *Am J Med* 122 (Suppl): S26-S37, 2009.
- Melnikova I and Wages D: Anti-obesity therapies. *Nat Rev Drug Discov* 5: 369-370, 2006.
- Cao Z, Umek RM and McKnight SL: Regulated expression of three C/EBP isoforms during adipose conversion of 3T3-L1 cells. *Genes Dev* 5: 1538-1552, 1991.
- Farmer SR: Transcriptional control of adipocyte formation. *Cell Metab* 4: 263-273, 2006.
- Stephens JM, Morrison RF and Pilch PF: The expression and regulation of STATs during 3T3-L1 adipocyte differentiation. *J Biol Chem* 271: 10441-10444, 1996.
- Zhang K, Guo W, Yang Y and Wu J: JAK2/STAT3 pathway is involved in the early stage of adipogenesis through regulating C/EBP $\beta$  transcription. *J Cell Biochem* 112: 488-497, 2011.
- Martini CN, Plaza MV and Vila MC: PKA-dependent and independent cAMP signaling in 3T3-L1 fibroblasts differentiation. *Mol Cell Endocrinol* 298: 42-47, 2009.
- Zhou Y, Wang D, Li F, Shi J and Song J: Different roles of protein kinase C- $\beta$ 1 and - $\delta$  in the regulation of adipocyte differentiation. *Int J Biochem Cell Biol* 38: 2151-2163, 2006.
- Prusty D, Park BH, Davis KE and Farmer SR: Activation of MEK/ERK signaling promotes adipogenesis by enhancing peroxisome proliferator-activated receptor gamma (PPARgamma) and C/EBPalpha gene expression during the differentiation of 3T3-L1 preadipocytes. *J Biol Chem* 277: 46226-46232, 2002.
- Chen LS, Redkar S, Taverna P, Cortes JE and Gandhi V: Mechanisms of cytotoxicity to Pim kinase inhibitor, SGI-1776, in acute myeloid leukemia. *Blood* 118: 693-702, 2011.
- Nawijn MC, Alendar A and Berns A: For better or for worse: the role of Pim oncogenes in tumorigenesis. *Nat Rev Cancer* 11: 23-34, 2011.
- Cervantes-Gomez F, Chen LS, Orlowski RZ and Gandhi V: Biological effects of the Pim kinase inhibitor, SGI-1776, in multiple myeloma. *Clin Lymphoma Myeloma Leuk* 13 (Suppl 2): S317-S329, 2013.
- Behan JW, Yun JP, Proektor MP, Ehsanipour EA, Arutyunyan A, Moses AS, Avramis VI, Louie SG, Butturini A, Heisterkamp N and Mittelman SD: Adipocytes impair leukemia treatment in mice. *Cancer Res* 69: 7867-7874, 2009.
- Nga ME, Swe NN, Chen KT, Shen L, Lilly MB, Chan SP, Salto-Tellez M and Das K: PIM-1 kinase expression in adipocytic neoplasms: diagnostic and biological implications. *Int J Exp Pathol* 91: 34-43, 2010.
- Meloche J, Paulin R, Courboulain A, Lambert C, Barrier M, Bonnet P, Bissierier M, Roy M, Sussman MA, Agharazii M and Bonnet S: RAGE-dependent activation of the oncoprotein Pim1 plays a critical role in systemic vascular remodeling processes. *Arterioscler Thromb Vasc Biol* 31: 2114-2124, 2011.
- More KN, Jang HW, Hong VS and Lee J: Pim kinase inhibitory and antiproliferative activity of a novel series of meridianin C derivatives. *Bioorg Med Chem Lett* 24: 2424-2428, 2014.
- Park YK, Lee TY, Choi JS, Hong VS, Lee J, Park JW and Jang BC: Inhibition of adipogenesis and leptin production in 3T3-L1 adipocytes by a derivative of meridianin C. *Biochem Biophys Res Commun* 452: 1078-1083, 2014.
- Song NJ, Yoon HJ, Kim KH, Jung SR, Jang WS, Seo CR, Lee YM, Kweon DH, Hong JW, Lee JS, *et al*: Butein is a novel anti-adipogenic compound. *J Lipid Res* 54: 1385-1396, 2013.
- Rosen ED and Spiegelman BM: Molecular regulation of adipogenesis. *Annu Rev Cell Dev Biol* 16: 145-171, 2000.
- Linhardt HG, Ishimura-Oka K, DeMayo F, Kibe T, Repka D, Poindexter B, Bick RJ and Darlington GJ: C/EBPalpha is required for differentiation of white, but not brown, adipose tissue. *Proc Natl Acad Sci USA* 98: 12532-12537, 2001.
- Shang CA and Waters MJ: Constitutively active signal transducer and activator of transcription 5 can replace the requirement for growth hormone in adipogenesis of 3T3-F442A preadipocytes. *Mol Endocrinol* 17: 2494-2508, 2003.
- Wang D, Zhou Y, Lei W, Zhang K, Shi J, Hu Y, Shu G and Song J: Signal transducer and activator of transcription 3 (STAT3) regulates adipocyte differentiation via peroxisome-proliferator-activated receptor gamma (PPARgamma). *Biol Cell* 102: 1-12, 2010.
- Park YK, Lee J, Hong VS, Choi JS, Lee TY and Jang BC: Identification of KMU-3, a novel derivative of gallic acid, as an inhibitor of adipogenesis. *PLoS One* 9: e109344, 2014.
- Christy RJ, Yang VW, Ntambi JM, Geiman DE, Landschulz WH, Friedman AD, Nakabeppu Y, Kelly TJ and Lane MD: Differentiation-induced gene expression in 3T3-L1 preadipocytes: CCAAT/enhancer binding protein interacts with and activates the promoters of two adipocyte-specific genes. *Genes Dev* 3: 1323-1335, 1989.
- Gustafson B, Jack MM, Cushman SW and Smith U: Adiponectin gene activation by thiazolidinediones requires PPAR gamma 2, but not C/EBP alpha-evidence for differential regulation of the aP2 and adiponectin genes. *Biochem Biophys Res Commun* 308: 933-939, 2003.
- Hwang CS, Loftus TM, Mandrup S and Lane MD: Adipocyte differentiation and leptin expression. *Annu Rev Cell Dev Biol* 13: 231-259, 1997.
- Lakshmanan MR, Nepokroeff CM and Porter JW: Control of the synthesis of fatty-acid synthetase in rat liver by insulin, glucagon, and adenosine 3':5' cyclic monophosphate. *Proc Natl Acad Sci USA* 69: 3516-3519, 1972.
- Rubin CS, Hirsch A, Fung C and Rosen OM: Development of hormone receptors and hormonal responsiveness in vitro. Insulin receptors and insulin sensitivity in the preadipocyte and adipocyte forms of 3T3-L1 cells. *J Biol Chem* 253: 7570-7578, 1978.
- Gregoire FM, Smas CM and Sul HS: Understanding adipocyte differentiation. *Physiol Rev* 78: 783-809, 1998.
- Fève B: Adipogenesis: Cellular and molecular aspects. *Best Pract Res Clin Endocrinol Metab* 19: 483-499, 2005.