

Serrated polyposis associated with a family history of colorectal cancer and/or polyps: The preferential location of polyps in the colon and rectum defines two molecular entities

PATRÍCIA SILVA^{1*}, CRISTINA ALBUQUERQUE^{1*}, PEDRO LAGE^{2,3}, VANESSA FONTES¹, RICARDO FONSECA⁴, INÊS VITORIANO¹, BRUNO FILIPE¹, PAULA RODRIGUES³, SUSANA MOITA¹, SARA FERREIRA^{2,3}, RITA SOUSA^{2,5}, ISABEL CLARO^{2,3}, CARLOS NOBRE LEITÃO^{2,6}, PAULA CHAVES⁴ and ANTÓNIO DIAS PEREIRA²

¹Molecular Pathobiology Research Unit (UIPM), ²Gastroenterology Service, ³Familial Cancer Risk Clinic,

⁴Pathology Service, Portuguese Institute of Oncology of Lisbon Francisco Gentil, E.P.E. (IPOLFG, EPE), Lisbon, Portugal

Received November 13, 2015; Accepted March 11, 2016

DOI: 10.3892/ijmm.2016.2666

Abstract. Serrated polyposis (SPP) is characterized by the development of multiple serrated polyps and an increased predisposition to colorectal cancer (CRC). In the present study,

we aimed to characterize, at a clinical and molecular level, a cohort of SPP patients with or without a family history of SPP and/or polyps/CRC (SPP-FHP/CRC). Sixty-two lesions from 12 patients with SPP-FHP/CRC and 6 patients with sporadic SPP were included. The patients with SPP-FHP/CRC presented with an older mean age at diagnosis ($p=0.027$) and a more heterogeneous histological pattern of lesions ($p=0.032$) than the patients with sporadic SPP. We identified two molecular forms of SPP-FHP/CRC, according to the preferential location of the lesions: proximal/whole-colon or distal colon. Mismatch repair (MMR) gene methylation [mutS homolog 6 (MSH6)/mutS homolog 3 (MSH3)] or loss of heterozygosity (LOH) of D2S123 (flanking *MSH6*) were detected exclusively in the former ($p=3.0 \times 10^{-7}$), in most early lesions. Proximal/whole-colon SPP-FHP/CRC presented a higher frequency of O-6-methylguanine-DNA methyltransferase (*MGMT*) methylation/LOH, microsatellite instability (MSI) and Wnt mutations (19/29 vs. 7/17; 16/23 vs. 1/14, $p=2.2 \times 10^{-4}$; 15/26 vs. 2/15, $p=0.006$; 14/26 vs. 4/20, $p=0.02$) but a lower frequency of B-raf proto-oncogene, serine/threonine kinase (*BRAF*) mutations (7/30 vs. 12/20, $p=0.0089$) than the distal form. CRC was more frequent in cases of Kirsten rat sarcoma viral oncogene homolog (*KRAS*)-associated proximal/whole-colon SPP-FHP/CRC than in the remaining cases (4/4 vs. 1/8, $p=0.01$). Thus, SPP-FHP/CRC appears to be a specific entity, presenting two forms, proximal/whole-colon and distal, which differ in the underlying tumor initiation pathways. Early *MGMT* and MMR gene deficiency in the former may underlie an inherited susceptibility to genotoxic stress.

Correspondence to: Dr Cristina Albuquerque, Molecular Pathobiology Research Unit (UIPM), Portuguese Institute of Oncology of Lisbon Francisco Gentil, E.P.E. (IPOLFG, EPE), Rua Professor Lima Basto, 1099-023 Lisbon, Portugal
E-mail: mc.albuquerque@sapo.pt

Present addresses: ⁵Garcia de Orta Hospital, E.P.E., Almada, Portugal; ⁶Lusíadas Lisbon Hospital, Lisbon, Portugal

*Contributed equally

Abbreviations: AD, adenomatous; APC, adenomatous polyposis coli; AD/S, adenomatous/serrated; *BRAF*, B-raf proto-oncogene, serine/threonine kinase; Ca, carcinoma; CIMP, CpG island methylator phenotype; CRC, colorectal cancer; *CTNBI*, catenin beta 1; *KRAS*, Kirsten rat sarcoma viral oncogene homolog; HCM, hyperplastic colonic mucosa; HP, hyperplastic polyp; LOH, loss of heterozygosity; MDE, mutation detection enhancement; *MGMT*, O-6-methylguanine-DNA methyltransferase; *MLH1*, mutL homolog 1; MMR, mismatch repair; MSI, microsatellite instability; MSI-H, microsatellite instability-high; MSI-L, microsatellite instability-low; MSS, microsatellite stable; *MSH3*, mutS homolog 3; *MSH6*, mutS homolog 6; NCM, normal colonic mucosa; *NRAS*, neuroblastoma RAS viral (v-ras) oncogene homolog; PCR, polymerase chain reaction; PTT, protein truncation test; SCa, serrated carcinoma; SE, serrated; SPP, serrated polyposis; SPP-FHP/CRC, SPP associated with a family history of SPP and/or polyps/CRC (multiple or diagnosed at a young age) in first-degree relatives; SSA, sessile serrated adenoma; SSCP, single-strand conformational polymorphism; TA, tubular adenoma; TSA, traditional serrated adenoma; TVA, tubulovillous adenoma

Key words: colorectal cancer, genotoxic susceptibility, O-6-methylguanine-DNA methyltransferase, mismatch repair gene methylation, serrated polyposis, proximal vs. distal colon

Introduction

Serrated polyposis (SPP), which was previously known as hyperplastic polyposis, is characterized by the presence of multiple colorectal epithelial polyps with a serrated architecture, termed serrated (SE) polyps, as well as an increased predisposition to colorectal cancer (CRC) (1-3). SE polyps differ from adenomatous (AD) polyps and are comprised of various lesions,

namely: hyperplastic polyps (HPs), non-dysplastic lesions with normal proliferation and architecture but elongated crypts with a saw-toothed appearance; sessile serrated adenomas (SSAs), lesions that present abnormal proliferation and architecture and may or may not include dysplasia; and traditional serrated adenomas (TSAs) that are dysplastic polyps with prominent serration (1,4-6). The presence of multiple SE polyps has also been associated with other hereditary conditions, namely serrated pathway syndrome or Jass syndrome (7-9).

It has been proposed that these SE lesions arise through the serrated pathway rather than through the adenoma-carcinoma sequence pathway (7,10-13). It has also been suggested that the HP is the precursor lesion in this pathway, with SSA as an intermediate step which then progresses to an adenocarcinoma with or without microsatellite instability (MSI or MSS, respectively). At the molecular level, SE lesions associated with those hereditary serrated syndromes share some genetic alterations, namely the presence of B-raf proto-oncogene, serine/threonine kinase (*BRAF*) mutations and the methylator phenotype, termed CpG island methylator phenotype (CIMP) (14-16), although these are also common to the sporadic SE lesions.

However, the analysis of SPP lesions has revealed specific features which are distinct from the SE lesions occurring in a sporadic context; accordingly, in SPP, HPs, TSAs and CRC are preferentially located in the proximal colon, i.e. proximal to the splenic flexure (17-19). Moreover, patients with SPP have previously been found to present extensive DNA methylation in the normal mucosa of the proximal colon (20), suggesting the involvement of widespread gene promoter methylation (CIMP) (7,14,21). This phenotype appears to be related to MutL homolog 1 (*MLH1*) methylation, which has been associated with MSI status, rather than O-6-methylguanine-DNA methyltransferase (*MGMT*) methylation. However, MSI appears to be less frequent in SPP lesions than in sporadic lesions (14,22). Similarly, SPP lesions also exhibited a lower frequency of Kirsten rat sarcoma viral oncogene homolog (*KRAS*) mutations when compared with sporadic SE lesions (21,22). However, it was also found that most adenomas and CRCs from patients with SPP exhibited a classic morphology and that few of these had *BRAF* or *KRAS* mutations, although SE lesions presented a high frequency of mutations in these genes (22). Previous research has suggested that tumorigenesis associated with SPP may not necessarily follow the serrated pathway and may follow an alternate pathway, involving TSAs and tubulovillous adenomas (TVAs) as intermediate lesions that progress to MSS adenocarcinomas with *KRAS* mutations or even a traditional pathway with adenomatous polyposis coli (*APC*) mutations as the initiating events (10).

A review of published case studies of SPP reported that approximately 10-50% of patients with SPP have been described as having a family history of CRC (23). In agreement with these findings, several studies have described an increased risk of CRC in the first-degree relatives of probands diagnosed with SPP compared with the general population (2,17,18,24-28). These studies have contributed to the notion that there are SPP cases where heredity may play a role in the development of CRC and/or polyps. Indeed, another review on this subject reinforces the concept that familial SPP exists and also the importance of defining the genetic basis of familial SPP and of studying these families in a systematic manner (29).

Thus, in the present study, we aimed to characterize, at the clinical and molecular level, SE and AD lesions from a cohort of patients with SPP who had been stratified into two groups: patients with or without a family history of SPP and/or polyps/CRC in first-degree relatives, in order to elucidate the information available regarding this new SPP entity with an apparent hereditary component.

Patients and methods

Patients and specimens. Eighteen patients diagnosed with SPP according to the WHO diagnostic criteria (1) were included in this study: 12 patients with SPP associated with a family history of SPP and/or polyps/CRC (multiple or diagnosed at a young age) in first-degree relatives (designated herein as SPP-FHP/CRC) (11 index and one affected relative diagnosed simultaneously with the index patient), and 6 index patients without a family history of SPP/polyps/CRC (designated herein as sporadic SPP) from the familial colorectal cancer registry of the Portuguese Institute of Oncology of Lisbon Francisco Gentil (Lisbon, Portugal). No evidence of SPP and/or polyps/CRC was found in the first-degree relatives of the patients with sporadic SPP, either by regular colonoscopy examination or by the absence of symptoms. The patients were classified as presenting a family history of polyps in first-degree relatives (5/11), if at least one relative had been diagnosed with polyps at or under 52 years of age or with >10 polyps. We cannot exclude the possibility that some of these families, namely PH4 or PH6, may have Jass syndrome instead of SPP, due to the presence of a mixture of AD and SE lesions (7,9). All patients had developed >10 lesions prior to the date of recruitment.

Sixty-two lesions were included here: 1 hyperplastic colonic mucosa (HCM), 25 HPs, 8 TSAs, 11 SSAs, 1 adenomatous/serrated (AD/S) carcinoma (Ca), 1 serrated carcinoma (SCa), 8 tubular adenomas (TAs), 2 TVAs and 5 Ca. Two normal colonic mucosa (NCM) samples were also included in this study. Fresh colorectal lesions were obtained from colectomy or colonoscopy specimens from patients who underwent surgery or colonoscopy in the Portuguese Institute of Oncology of Lisbon Francisco Gentil. Sections from corresponding areas of the specimens submitted for diagnosis were divided into two parts: one was snap frozen in liquid nitrogen immediately after resection, while the other was formalin-fixed and paraffin-embedded.

Histological characterization, according to the WHO guidelines (1), was performed by experienced pathologists (R.F. and P.C.). The study was conducted in accordance with local ethical standards and in agreement with the Helsinki Declaration of 1975, as revised in 1983. Informed consent for diagnosis and additional investigational studies, which may result in improving the knowledge about the pathogenesis of the disease, was obtained from patients included in this study. Moreover, biological material used for DNA isolation was obtained from archival sections from colorectal adenomas and carcinomas specimens, submitted for diagnosis (histological classification) and derived from patients who underwent surgery or colonoscopy in the Portuguese Institute of Oncology of Lisbon Francisco Gentil; only somatic analysis was performed and samples are truly anonymized.

Methods

DNA isolation. DNA was isolated from fresh-frozen and/or paraffin-embedded tumor tissue and matched normal tissue. DNA was isolated from the paraffin-embedded tissues by proteinase K digestion, which was followed by phenol/chloroform extraction and ethanol precipitation, as previously described (30). DNA from the fresh frozen tissue was isolated by proteinase K digestion, followed by precipitation with a saturated NaCl solution and ethanol as previously described (31).

Somatic mutation analysis of the APC gene. APC mutation analysis was performed using the protein truncation test (PTT) for exon 15, as previously reported (30). Briefly, APC exon 15 was divided into four overlapping fragments that were amplified by polymerase chain reaction (PCR) and, subsequently, *in vitro* transcription and translation were performed using a TnT T7-coupled reticulocyte lysate system (Promega, Madison, WI, USA). In negative cases, the mutational cluster region was subsequently analyzed by automated sequencing in order to search for missense mutations. For the samples obtained from paraffin-embedded tissues, APC mutations were analyzed by single-strand conformational polymorphism (SSCP) or by automated sequencing (32).

Somatic mutation analysis of catenin beta 1 (CTNNB1). Genomic DNA from each tumor sample was amplified by PCR for SSCP analysis of exon 3 of the CTNNB1 gene. The amplified products were analyzed in a mutation detection enhancement (MDE) gel and visualized by silver staining (33).

Somatic mutation analysis of the AXIN2 gene. AXIN2 mutation analysis was performed by amplification of a repetitive sequence containing the (G)₇, (C)₆ and (C)₅ tracts (where a considerable mutation frequency was described) (34), followed by electrophoresis in 7% polyacrilamide gel containing formamide and urea and visualization by silver staining (33).

Somatic mutation analysis of BRAF, KRAS and neuroblastoma RAS viral (v-ras) oncogene homolog (NRAS) genes. BRAF (exon 15: forward primer 5'-TCATAATGCTTGCTCTGATAGGA-3'; reverse primer 5'-GGCCAAAATTTAATCAGTGGGA-3'), KRAS (exon 2: forward primer 5'-GTGTGACATGTTCTAATATAGTCA-3'; reverse primer 5'-GAATGGTCCTGCACCAGTAA) (35) and NRAS (exons 2 and 3-primers were kindly provided by Dr Branca Cavaco), were amplified by PCR. The DNA samples were amplified in a standard PCR buffer (Invitrogen, Waltham, MA, USA). Mutations in these genes were analyzed by automated sequencing.

Sequencing analysis. After amplification, PCR products were purified with Illustra GFX™ PCR DNA and Gel Band purification kit (GE Healthcare, Little Chalfont, UK) according to the manufacturer's instructions. Sequencing reactions were performed using the BigDye Terminator Cycle Sequencing kit and the respective products were analyzed on the ABI PRISM™ 310 Genetic Analyzer (both from Applied Biosystems, Foster City, CA, USA) using Sequencing Analysis software. The pathogenic relevance of missense variants was evaluated by comparing aminoacid sequences using PolyPhen software (<http://genetics.bwh.harvard.edu/pph/>) and SIFT software (<http://sift.jcvi.org/>).

MSI/loss of heterozygosity (LOH) analysis. MSI status was analyzed using the Bethesda panel of reference markers (36). Each colonic lesion and paired normal DNA were amplified

by PCR for each of the microsatellite markers and analyzed in the ABI Prism™ 310 Genetic Analyzer using GeneScan software (Applied Biosystems). The lesions were classified as MSI-high (H) when showing MSI in two or more of the five markers, MSI-low (L) when MSI was detected in one of the markers, and MSS when none of the markers revealed instability (37). In cases exhibiting MSI-L, BAT-40 and MYCL markers were also analyzed and the lesions were classified as MSI-H when MSI was detected in >40% of the 7 markers analyzed; otherwise they were classified as MSI-L, as previously described (37).

A total of 4 dinucleotide markers flanking MGMT (D10S1703, D10S1676, D10S169 and D10S1651) were analyzed for each colonic lesion and paired normal DNA in order to evaluate the presence of LOH. Each lesion was subsequently scored as demonstrating LOH if the ratio between the areas of the normal and the tumor alleles was >1.5 or <0.67.

Regarding the D5S346 marker, LOH was evaluated (and confirmed using the D5S1965 marker) as indicative of loss of the APC gene. LOH of D2S123 and D17S250 was also evaluated as described above.

Methylation analysis. The analysis of MGMT and mismatch repair (MMR) gene promoter methylation was performed by methylation-specific multiplex ligation-dependent probe amplification (MS-MLPA) (38) using the SALSA MS-MLPA KIT ME011 MMR, (MRC-Holland, Amsterdam, The Netherlands). MS-MLPA reactions were performed as described by the manufacturer. The samples were analyzed using GeneScan software on the ABI Prism™ 310 Genetic Analyzer (Applied Biosystems). The results were normalized using MRC Coffalyser MLPA-DAT software v.9.4 (MRC-Holland). A ratio of 0.15 or higher, corresponding to 15% of methylated DNA, was indicative of promoter methylation as described elsewhere (32,39).

Statistical analysis. Fisher's exact test (using a two-sided or 2x3 table) and the χ^2 test (<http://www.quantitativeskills.com/sisa/index.htm>) were used to compare categorical variables, and the Student's t-test (<http://www.physics.csbsju.edu/stats/t-test.html>) was used to compare continuous variables. A p-value <0.05 was considered to indicate a statistically significant difference.

Results

Clinical characterization. Table I summarizes the clinical features of the 18 patients included in this study, stratified into two groups: SPP-FHP/CRC and sporadic SPP. The average age at diagnosis (i.e. the age at which they presented symptoms) among our cohort of SPP patients (n=18) was 55±11 years (range 25-80); however, it was significantly higher in the SPP-FHP/CRC group than in the sporadic SPP group [60±10 years (range 41-80) vs. 46±15 years (range 25-68), p=0.027 (Student's t-test)] (Table II).

The number of lesions was higher in the SPP-FHP/CRC group than in the sporadic SPP group (threshold, ≥40 lesions): [10/12 (83%) vs. 3/6 (50%) patients, respectively]. With respect to histological features, AD lesions were more frequent in the SPP-FHP/CRC group than in the patients with sporadic SPP [10/12 (83%) vs. 1/6 (17%), p=0.013]. Moreover, the patients

Table 1. Clinical features of the patients evaluated in this study, histological characterization of the respective lesions and family history of SPP and/or polyyps/CRC in first-degree relatives.

Family	Patient ID	Age at diagnosis in years	Gender	Total number of lesions	Type of lesions ^a	Preferential location of lesions ^b	CRC	WHO diagnostic criteria ^c	No. of affected individuals in the family	Family history ^d (age at diagnosis in years)
SPP-FHP/CRC										
PH1	A756	62	M	>100	HP+TALGD+TSALGD+SSA	Whole colon ^e	No	3	3	Son, >45 SE polyyps (42); Son, 2 SE polyyps (37)
PH3	A755	41	F	>100	HP+TSALGD+TVALGD+SSA+TALGD	Whole colon ^e	No	3	3	Mother, CRC (67); maternal aunt, CRC (58)
PH4	CA638	66	M	19	TSA (LGD+HGD)+VA	Whole colon	Yes ^f	1	5	Twin brother, 20 AD and SE polyyps (70); son, 2 AD polyyps (37); nephew, 18 AD polyyps (54)
PH5	CA636	69	M	50	HP+TALGD+TSALGD+TVALGD ^g	Proximal	Yes ^f	3	3	Sister, 2 AD polyyps (67); son, 1 SE polyp (52)
PH5	A193	69	F	40	HP+TALGD+TVAHGD	Proximal	Yes	3 plus 1	3	Father, CRC (57); sister, 7 polyyps (61); daughter, 1 SE polyp (39); son, 2 AD polyyps (35)
PH6	A478	64	M	50-100	HP+TSALGD+TALGD+SSA	Distal ^e	No	3 plus 1	5	Mother, 2 AD polyyps (75); brother, 8 SE and AD polyyps (47)
PH7	A759	50	M	50-100	HP+TA	Whole colon	Yes	3	3	Maternal grandfather, CRC; mother, CRC (73); sister, 3 AD polyyps (61)
PH8	A760	59	F	45	HP+SSA+TSALGD	Proximal	No	1 plus 3	4	Brother, 1 AD polyp (44)
PH12	A758	57	M	40	HP	Distal	Yes	3	2	Daughter, 2 SE polyyps (54); daughter, 11 SE and 1 AD polyyps (42); son, 1 AD polyp (44)
PH14	A686	80	M	49	HP+TSALGD+TALGD	Distal	No	3	4	Mother, CRC; sister, 6 polyyps (68)
PH19	A993	58	F	17	HP+TALGD	Distal	No	1	3	Paternal uncle, polyyps (69); father, 1 AD polyp (77); brother, 1 AD polyp (50)
PH33	A983	49	M	45	HP+TSALGD+TALGD+SSA	Distal	No	1 plus 3	4	
Sporadic SPP										
PH9	A500	48	F	40	HP+SSA+TSALGD	Whole colon	No	1 plus 3	1	

Table I. Continued.

Family	Patient ID	Age at diagnosis in years	Gender	Total number of lesions	Type of lesions ^a	Preferential location of lesions ^b	CRC	WHO diagnostic criteria ^c	No. of affected individuals in the family	Family history ^d (age at diagnosis in years)
PH10	A989	25	F	<100	HP	NA	Yes	3 plus 1	1	^h
PH11	A757	53	M	50	HP+TSA	Distal	Yes ^f	3	1	^h
PH16	A990	68	M	15	HP+SSA	Whole colon	No	1	1	^h
PH18	A992	46	F	10	TSALGD+HP	Distal	No	1	1	^h
PH22	A951	33	F	31	HP+TALGD	Distal	No	1 plus 3	1	^h

^aThe type of lesion(s) that was prevalent ($\geq 70\%$) in each patient is indicated in bold. ^bWe considered a proximal or distal preferential location of the lesions when at least 70% of the lesions (majority serrated) were located in the proximal or distal colon, respectively. ^cWHO clinical criteria for the identification of serrated polyposis (SPP), as revised in 2010: i) At least five serrated polyps proximal to the sigmoid colon, two of which are greater than 10 mm in diameter. ii) Any number of serrated polyps occurring proximal to the sigmoid colon in an individual who has a first-degree relative with SPP. iii) More than 20 serrated polyps of any size distributed throughout the colon. ^dAll families showed a dominant inheritance pattern with the exception of PH12, that revealed a recessive pattern. ^eThe polyps are larger in the proximal colon. ^fAdenomatous/ serrated carcinoma (AD/S); ^gwith areas of traditional serrated adenoma (TSA). ^hNo evidence of SPP and/or polyps/CRC was found in first-degree relatives of the sporadic SPP patients, either by regular colonoscopy examination or by absence of symptoms. CRC, colorectal cancer; M, male; F, female; HP, hyperplastic polyp; TALGD, tubular adenoma with low grade dysplasia; TSALGD, traditional serrated adenoma with low grade dysplasia; TVALGD, tubulovillous adenoma with high grade dysplasia; SSA, sessile serrated adenoma; SE, serrated; TVALGD, tubulovillous adenoma with low grade dysplasia; LGD, low grade dysplasia; HGD, high grade dysplasia; VA, villous adenoma; AD, adenomatous; TA, tubular adenoma; NA, not available; FHP, family history of SPP.

with SPP-FHP/CRC presented a more heterogeneous spectrum of lesions in comparison with the patients with sporadic SPP. In agreement with these findings, the presence of three or more types of lesions was more frequent in the former group [9/12 (75%) vs. 1/6 (17%), $p=0.032$].

Regarding the location of the lesions in each patient, we observed that whereas some of the patients with SPP presented lesions dispersed uniformly throughout the whole colon, other patients with SPP presented a predominance of lesions in one of the two major segments, proximal or distal. Therefore, a prevalence of proximal or distal location of the lesions was considered when at least 70% of the lesions (majority SE) were located in the proximal or distal colon, respectively. In accordance, a preferential proximal location was observed in 3/12 (25%) of the SPP-FHP/CRC patients and in none of the patients with sporadic SPP (0/5). A preferential distal location was observed in 3/5 (60%) of the sporadic SPP patients and in 5/12 (42%) of the patients with SPP-FHP/CRC.

Molecular characterization. The molecular alterations found in each SPP-FHP/CRC lesion, namely mutations in RAS/RAF and Wnt signaling genes, MSI, *MGMT* and MMR methylation, LOH of *MGMT* locus and LOH at D2S123 and D17S250 markers, are presented in Table III. The spectra of these molecular alterations led us to observe that the somatic mutation/promoter hypermethylation spectra differs between those patients whose lesions were preferentially located in the proximal colon, or distributed throughout the whole colon, and those whose lesions were preferentially located in the distal colon, as shown in Table IV. This led us to stratify the patients with SPP-FHP/CRC into two groups, proximal/whole-colon and distal SPP-FHP/CRC.

Although the ratio between the different types of lesions in the two groups, proximal/whole-colon and distal, do not match, i.e. in the former a higher proportion of AD lesions have been analyzed [in a recent study, individuals with large and right-sided SE lesions also had significantly more AD lesions compared with those without such types of SE lesions (40)], we found differences with respect to the mutation spectrum, even considering only HPs or SSAs, which have been analyzed in both groups. Accordingly, HPs and SSAs from patients with proximal/whole-colon SPP-FHP/CRC presented RAS/RAF gene mutations less frequently [2/6 (33%) and 1/7 (14%) vs. 14/14 and 2/2, respectively, $p=0.003$ and $p=0.08$] whereas MMR gene methylation or LOH of D2S123 [flanking mutS homolog 6 (*MSH6*)] occurred more frequently (4/4 and 2/2 vs. 0/7 and 0/2, respectively, $p=0.003$ and $p=0.33$), when compared with the same type of lesions from distal SPP-FHP/CRC patients (Table V). Moreover, SE and AD lesions presented a similar spectrum of molecular alterations, especially among each patient. The exception were *BRAF* mutations that were significantly more frequent in SE lesions [17/36 (47%) vs. 2/14 (14%), $p=0.05$] (Table V). For each patient, either presenting proximal/whole-colon or distal SPP-FHP/CRC, the spectra of somatic molecular alterations detected in the lesions were similar regardless of the location of each specific lesion. For example, in a patient with a prevalence of proximal lesions, these presented a specific mutation pattern that was shared by the few distal lesions that were analyzed from the same patient and different from the

Table II. Comparison between clinical features in patients with sporadic SPP and those with SPP-FHP/CRC.

Clinical feature	SPP-FHP/CRC	Sporadic SPP	p-value
Age at diagnosis (years)	60±10	46±15	0.027 (Student's t-test)
Preferential location of lesions			
Whole colon	4/12 (33%)	2/5 (40%)	NS
Proximal	3/12 (25%)	0/5	NS
Distal	5/12 (42%)	3/5 (60%)	NS
≥40 lesions	10/12 (83%)	3/6 (50%)	NS
AD lesions	10/12 (83%)	1/6 (17%)	0.013^a
≥3 types of lesions	9/12 (75%)	1/6 (17%)	0.032^a

Statistically significant values are shown in bold. NS, non-significant ($p > 0.05$). ^aFisher's exact test (two-sided). AD, adenomatous; SPP, serrated polyposis; SPP-FHP/CRC, SPP associated with a family history of SPP and/or polyps/colorectal cancer (CRC) (multiple or diagnosed at a young age) in first-degree relatives.

proximal lesions from patients with a predominance of distal lesions and vice versa (Table IIIA).

The abovementioned findings led us to analyze the molecular data from the patients with SPP-FHP/CRC, who were stratified into two groups: proximal/whole-colon and distal SPP-FHP/CRC.

Proximal/whole-colon vs. distal SPP-FHP/CRC. Mutations in the Wnt genes, as well as MSI, were significantly more frequent in the patients with proximal/whole-colon SPP-FHP/CRC than in the patients with distal SPP-FHP/CRC [14/26 (54%) vs. 4/20 (20%), $p = 0.02$; 15/26 (58%) vs. 2/15 (13%), $p = 0.0059$] (Table IV). Interestingly, among the proximal/whole-colon SPP-FHP/CRC samples, Wnt gene mutations were significantly more frequent in the SE lesions than in the AD lesions [12/16 (75%) vs. 2/10 (20%), $p = 0.0091$]. In the SE lesions, Wnt gene mutations, as well as MSI, were more frequent in TSAs (4/4 and 3/4) and in SSAs (6/7 and 6/7) and rarely detected in HPs (1/4 and 1/5) ($p = 0.02$ and $p = 0.017$, respectively) (Table V).

Similarly, LOH of the *MGMT* locus and *MGMT* methylation were also more frequent in lesions from the patients with proximal/whole-colon SPP-FHP/CRC, in comparison with lesions from the patients with distal SPP-FHP/CRC [16/23 (70%) vs. 1/14 (7%), $p = 2.2 \times 10^{-4}$; 19/29 (65%) vs. 7/17 (41%), $p = 0.1$, respectively] (Table IV). The same difference was observed even considering SE lesions only (Table V). In particular, *MGMT* methylation was detected in all HP lesions (6/6) from proximal/whole-colon SPP-FHP/CRC, in contrast to the lower frequency observed in HPs from distal SPP-FHP/CRC (4/11, 36%) ($p = 0.017$).

Interestingly, among lesions which were deemed informative for methylation analysis and loss of LOH of D2S123, the presence of MMR gene methylation or of the D2S123 LOH (flanking *MSH6*) was only observed in proximal/whole-colon SPP-FHP/CRC lesions [17/18 (94%) vs. 0/11, $p = 3.0 \times 10^{-7}$]. These alterations were found in the majority of early lesions and in all histological types (Tables IIIA and IV). It is of note that *MLH1* methylation was not detected in any of the SPP-FHP/CRC lesions, being observed only in one lesion from

the sporadic SPP group (Table IIIB). Moreover, it is also of note that except for one lesion, MSI, either MSI-L or MSI-H was detected only in dinucleotide microsatellite markers.

By contrast to the abovementioned molecular alterations, in the SPP-FHP/CRC samples, *BRAF* and *KRAS* mutations were more frequent in the lesions located in the distal colon than those located in the proximal/whole-colon [12/20 (60%) vs. 7/30 (23%), $p = 0.0089$ (χ^2 test) and 6/20 (30%) vs. 5/32 (16%), respectively], although the latter was not statistically significant (Table IV). Considering only the SE lesions, mutations in the RAS/RAF genes were detected in all the lesions from the distal SPP-FHP/CRC, namely in all HPs (14/14), HCM (1/1) and in one NCM (Table IIIA), but in only 8/19 (42%) lesions from the proximal/whole-colon SPP-FHP/CRC ($p = 1.3 \times 10^{-4}$).

With respect to *KRAS/BRAF* mutations, two groups of patients were observed among those with either proximal/whole-colon or distal SPP-FHP/CRC: one group whose lesions presented *KRAS* mutations (PH4, PH5, PH7 and PH14) and another group, whose lesions presented *BRAF* mutations (PH3, PH8, PH19 and PH33) (Table IIIA). One patient presented either *BRAF*- or *KRAS*-positive lesions (family PH6). Notably, CRC was more frequent in the patients with proximal/whole-colon SPP-FHP/CRC associated with *KRAS* mutations than in the remaining patients [4/4 vs. 1/8 (12%), $p = 0.01$] (Tables I and IIIA).

Discussion

SPP-FHP/CRC and sporadic SPP differ at the clinical level.

The patients with SPP-FHP/CRC and sporadic SPP differed with respect to clinical and histological features. The older mean age at diagnosis (60 vs. 46 years old) of the former may underlie a slower process of tumorigenesis. This is in contrast to that which has been observed in relation to other hereditary CRC syndromes, namely in familial adenomatous polyposis and in Lynch syndrome, which are diagnosed at an earlier age (usually ≤ 50 years old) compared to the age at diagnosis in patients with sporadic CRC (> 60 years old) (32,41). The presence of a more heterogeneous histological pattern of lesions in patients with SPP-FHP/CRC, associated with the concomitant presence of

Table III. Results of the molecular analysis of SE and AD lesions from patients with SPP.

Family	Patient ID	Lesion ID	Type of lesion	Location of lesions	MSI	D2S123, D17S250 LOH	LOH of MGMT locus	Mutations					Promoter methylation		
								APC	CTNNB1	AXIN2	BRAF	KRAS	MGMT	MMR genes	
Preferential location of lesions - proximal/whole-colon															
PH1	A756	A756AS	TSALGD	NA	MSS'	NC, NC	Yes	c.4099C>T (p.Q1367X)	N	N	-	N	N	M	MSH6
		A756AT1	TALGD	NA	MSS'	NC, N	No	c.4123C>T (p.H1375Y)	N	N	N	N	M	MSH6	
		A756AT2	TALGD	NA	MSS'	NC, N	IC	c.4123C>T (p.H1375Y)	N	N	N	N	NM	NM	
		A756PH1	HP	NA	MSS'	NC, NC	Yes	N	N	-	N	N	M	MSH6, MSH3	
		A756PH2	SSA	NA	MSS'	NC, N	IC	c.4289delC (p.T1430TfsX43)	N	N	-	N	M	MSH6	
PH3	A755	A755AS1	TSALGD	Distal	MSI-H	NI, D17S250	IC	NC	N	N	c.1785T>G (p.F595C)	N	-	-	-
		A755AS2	TSA	Proximal	MSI-L	NI, D17S250	IC	c.4235G>A (p.G1412E)	c.115G>A (p.A39T)	N	N	c.1799T>A (p.V600E)	N	M	MSH6
		A755AS3	TSALGD	Proximal	MSI-H	NI, NI	Yes	NC	N	N	N	N	NM	MSH6	
		A755PH1	HP	Proximal	MSI-L	NI, D17S250	Yes	N	N	N	N	N	M	MSH6	
		A755PH2	SSA	Proximal	MSI-L	D2S123, D17S250	IC	N	N	N	N	N	NM	NM	
PH4	CA638	CA638AS	TSALGD	Distal	MSS	N, N	No	N	c.122C>T (p.T411)	N	N	N	c.35G>C (p.G12A)	M	MSH3
		CA638C	Ca (AD/S)	Rectum	MSS	N, N	No	LOH	N	N	N	N	c.35G>A (p.G12D)	M	NM
		CA636AT1	TALGD	Proximal	MSI-L	NI, D17S250	Yes	c.4262delG (p.S1421MfsX52)	N	-	N	N	M	NM	
		CA636C	Ca	Proximal	MSS	D2S123, N	Yes	NC	N	N	N	N	M	MLH3	
		CA636AT2	TALGD	Proximal	MSI-L	D2S123, NI	Yes	N	N	N	N	N	N	NM	NM
		CA636AT3	TALGD	Proximal	MSS	D2S123, D17S250	Yes	N	N	N	c.1811G>A (p.W604X)	N	N	M	NM

Table III. Continued.

Family	Patient ID	Lesion ID	Type of lesion	Location of lesions	MSI	D2S123, D17S250 LOH	LOH of MGMT locus	Mutations					Promoter methylation		
								APC	CTNNB1	AXIN2	BRAF	KRAS	MGMT	MMR genes	
PH5	A193	CA636ATV	TVALGD	Proximal	MSI-H	NI, NI	IC	c.4024T>G (p.L1342V)	N	N	N	N	c.38G>A (p.G13D)	M	NM
		A193AT	TALGD	Proximal	MSI-L	N, D17S250	IC	c.4189G>A (p.E1397K)	N	N	N	N	N	M	<i>MSH6</i>
		A193ATV	TVAHGD	Rectum	MSS	D2S123, N	Yes	c.4123C>T (p.H1375Y)	N	N	N	N	c.35G>A (p.G12D)	M	NM
		A193C	Ca	Proximal	MSS	D2S123, D17S250	Yes	N	N	N	N	N	N	M	NM
		A193PH1	HP	Proximal	MSS	D2S123, N	Yes	LOH, c.4123C>T (p.H1375Y)	N	N	N	N	N	M	NM
		A193PH2	HP	Proximal	MSS	D2S123, D17S250	Yes	NC	N	N	N	N	N	M	NM
PH7	A759	A759T	Ca	Proximal	MSI-H	NI, N	IC	N	c.133T>C (p.S45P)	c.1994delG (p.G665AfsX24)	N	c.34G>A (p.G12S)	NM	NM	
PH8	A760	A643PA	HP	Proximal	MSS	NI, N	No	N	N	N	c.1799T>A (p.V600E)	N	M	NM	
		A643PB	HP	Distal	MSS	NI, N	No	N	N	N	c.1799T>A (p.V600E)	N	M	NM	
		A760AS1	TSALGD	Proximal	MSS'	NC, NC	Yes	LOH	N	N	N	c.1799T>A (p.V600E)	-	-	
		A760PH1	SSA	NA	MSI-L	NI, D17S250	-	c.3926_3930 delAAAAGA (p.E1309DfsX2)	N	N	N	N	NM	NM	
		A760PH2	SSA	Proximal	MSI-H	NI, NC	Yes	NC	N	N	N	N	-	-	
		A760PH3	SSA	Proximal	MSI-L	NI, N	Yes	LOH	c.130C>T (p.P44S)	N	N	c.1799T>A (p.V600E)	NM	NM	
		A760PH4	SSA	NA	MSI-L	NI, NC	Yes	LOH	N	N	N	N	NM	NM	
		A760PH5	SSA	NA	MSI-L	NI, N	No	LOH	N	N	N	N	NM	NM	
		A760PH6	SSA	NA	MSS	NI, N	No	LOH	N	N	N	N	NM	NM	
Preferential location of lesions - distal															
PH6	A478	A478PA2	HP	NA	MSS	N, N	No	N	N	N	N	c.1799T>A (p.V600E)	M	NM	

Table III. Continued.

Family	Patient ID	Lesion ID	Type of lesion	Location of lesions	MSI	D2S123, D17S250 LOH	LOH of MGMT locus	Mutations					Promoter methylation		
								APC	CTNNB1	AXIN2	BRAF	KRAS	MGMT	MMR genes	
	A478PB	HP	NA	NA	MSS	N, N	-	N	N	N	c.1799T>A (p.V600E)	N	-	-	-
	A478PC	NCM	NA	NA	MSS	N, N	-	N	N	N	c.35G>T (p.G12V)	c.35G>T (p.G12V)	NM	NM	NM
	A478PD	NCM	NA	NA	MSS	N, N	-	-	-	N	N	-	-	-	-
	A478PE	HCM	NA	NA	MSS	N, N	-	N	N	N	N	c.35G>A (p.G12D)	NM	NM	NM
	A478PF	TALGD	NA	NA	MSS	N, N	-	N	N	N	N	N	NM	NM	NM
	A478PG	HP	NA	NA	MSS	N, N	No	N	N	N	c.1799T>A (p.V600E)	N	M	M	NM
	A478PH	HP	NA	NA	MSS	N, N	No	N	N	N	c.1799T>A (p.V600E)	N	NM	NM	NM
	A462PH1	SSA	NA	NA	MSS	N, N	No	N	N	N	c.1799T>A (p.V600E)	N	M	M	NM
	A462PH2	SSA	NA	NA	MSS	N, N	IC	N	N	N	c.1799T>A (p.V600E)	N	M	M	NM
PH12	A758	A758C	Ca	Distal	MSI-L	NI, NC	No	LOH	N	N	c.1799T>A (p.V600E)	N	M	M	NM
	A758PH	HP	Rectum	Rectum	MSI-L	NI, D17S250	IC	LOH	N	N	c.1799T>A (p.V600E)	N	M	M	NM
PH14	A686	A686P1A	HP	Distal	MSS	N, N	No	N	N	N	N	c.35G>A (p.G12D)	NM	NM	NM
	A686P2B	HP	Distal	Distal	MSS	N, N	No	N	N	N	N	c.35G>T (p.G12V)	M	M	NM
	A686P3C	HP	Distal	Distal	MSS	N, N	No	N	N	N	N	c.35G>A (p.G12D)	NM	NM	NM
	A686P4D	HP	Distal	Distal	MSS	N, N	No	N	N	N	N	c.35G>A (p.G12D)	NM	NM	NM
	A686P5E	HP	Distal	Distal	MSS	N, N	-	N	N	N	N	c.35G>T (p.G12V)	-	-	-
PH19	A993	A993PH1	HP	Rectum	MSS*	-	Yes	NC	c.115G>A (p.A39T)	N	N	c.1799T>A (p.V600E)	-	-	-

Table III. Continued.

Family	Patient ID	Lesion ID	Type of lesion	Location of lesions	MSI	D2S123, D17S250 LOH	LOH of <i>MGMT</i> locus	Mutations					Promoter methylation		
								<i>APC</i>	<i>CTNNB1</i>	<i>AXIN2</i>	<i>BRAF</i>	<i>KRAS</i>	<i>MGMT</i>	MMR genes	
PH33	A983	A983PA	HP	Proximal	MSS*	-	No	N	N	N	c.1799T>A (p.V600E)	N	NM	NM	NM
		A983PB	HP	Proximal	MSS*	-	No	N	N	N	c.1799T>A (p.V600E)	N	NM	NM	NM
		A983PC	HP	Distal	MSS*	-	No	N	N	N	c.1799T>A (p.V600E)	N	NM	NM	NM
		A983PD	TALGB	Distal	MSS*	-	No	c.4189G>T (p.E1397X)	N	N	N	N	NM	NM	NM

B, Results of the molecular analysis of SE and AD lesions from patients with SPP without a family history of SPP and/or polyps/CRC in first-degree relatives

Family	Patient ID	Lesion ID	Type of lesion	Location of lesions	MSI	D2S123, D17S250 LOH	<i>APC</i>	<i>CTNNB1</i>	<i>AXIN2</i>	<i>BRAF</i>	<i>KRAS</i>	Promoter methylation		
												<i>MGMT</i>	MMR genes	
Preferential location of lesions - Proximal/whole-colon														
PH9	A500	A500PA1	HP	Proximal	MSS	D2S123, NI	N	N	N	c.1799T>A (p.V600E)	N	NM	NM	<i>MLH1</i>
		A500PB2	HP	Proximal	MSS	N, NI	N	N	N	c.1799T>A (p.V600E)	N	NM	NM	NM
PH16	A990	A990ASS	SSA	NA	MSS*	-	NC	N	N	N	N	M	NM	NM
		A990PHI	HP	Rectum	MSS*	-	NC	N	N	N	N	NM	NM	<i>MSH3</i>
Preferential location of lesions - Distal														
PH10	A989	A989C	Ca	Distal	MSS*	-	NC	N	N	N	N	NM	NM	NM
PH11	A757	A757AS1	SCa	NA	MSS	N, N	c.4233delT (p.S1411RfsX4)	N	N	N	N	M	M	NM
PH18	A992	A992AS1	TSALGB	Distal	MSS*	-	NC	N	N	c.1799T>A (p.V600E)	N	-	-	-
		A992AS2	TSALGB	Distal	MSS*	-	NC	N	N	N	N	-	-	-
PH22	A951	A951P1	HP	Rectum	MSS*	-	N	N	N	c.1799T>A (p.V600E)	N	NM	NM	NM
		A951P2	HP	Distal	MSS*	-	N	N	N	c.1799T>A (p.V600E)	N	NM	NM	NM

SE, serrated; AD, adenomatous; MSI, microsatellite instability; LOH, loss of heterozygosity; MMR, mismatch repair; NA, not available; MSS*, microsatellite stable (missing one marker, D2S123); NC, not completed; N, no mutation; -, not performed; M, methylated; IC, inconclusive; MSI-H, microsatellite instability high; NI, not informative; MSI-L, microsatellite instability low; NCM, normal colonic mucosa; HCM, hyperplastic colonic mucosa; SCa, serrated carcinoma; *MSH*, mutS Homolog; SPP, serrated polyposis; CRC, colorectal cancer; MSS*, microsatellite stable (only performed for BAT-26, due to the lack of normal tissue for this sample). *APC* missense mutations were not taken into account, although p.H1375Y and p.E1397K mutations have been predicted by *in silico* analysis to be probably pathogenic (data not shown).

Table IV. Molecular characterization of lesions from patients with SPP-FHP/CRC, stratified by preferential location of the lesions in each patient.

Molecular characterization	Preferential location of lesions		p-value
	Proximal/whole-colon	Distal colon	
Total Wnt gene mutations	14/26 (54%)	4/20 (20%)	0.02^b
Total RAS/RAF gene mutations	12/30 (40%)	18/20 (90%)	3.7x10^{-4b}
<i>BRAF</i> gene mutations	7/30 (23%)	12/20 (60%)	0.0089 (χ^2 test)
<i>KRAS</i> gene mutations	5/32 (16%)	6/20 (30%)	NS
MSI ^a	15/26 (58%)	2/15 (13%)	0.0059^b
MMR gene methylation and/or LOH of D2S123	17/18 (94%)	0/11	3.0x10^{-7b}
<i>MGMT</i> gene methylation	19/29 (65%)	7/17 (41%)	NS
LOH of <i>MGMT</i> locus	16/23 (70%)	1/14 (7%)	2.2x10^{-4b}

^aExcept for one lesion, microsatellite instability (MSI), either microsatellite instability-low (MSI-L) or microsatellite instability-high (MSI-H), was detected only in dinucleotide microsatellite markers. ^bFisher's exact test (two-sided). SPP, serrated polyposis; CRC, colorectal cancer; SPP-FHP/CRC, SPP associated with a family history of SPP and/or polyps/CRC (multiple or diagnosed at a young age) in first-degree relatives; MMR, mismatch repair; LOH, loss of heterozygosity; *MGMT*, O-6-methylguanine-DNA methyltransferase. Statistically significant values are shown in bold. NS, non-significant (p>0.05).

both SE and AD lesions, compared with the lesions in patients with sporadic SPP, suggests the involvement of other pathways in the tumorigenic process associated with SPP-FHP/CRC, in addition to the serrated pathway of tumorigenesis.

Molecular alterations involved in tumor initiation distinguish between two forms of SPP-FHP/CRC: proximal/whole-colon and distal. Two forms of SPP-FHP/CRC appear to exist according to the preferential location of the lesions in the colon and rectum, proximal/whole-colon and distal, which differ with respect to the somatic events involved in tumor initiation. LOH and methylation of *MGMT*, MMR gene methylation and/or LOH of D2S123 and Wnt gene mutations appear to be the major somatic events that lead to tumor initiation in proximal/whole-colon SPP-FHP/CRC. By contrast, in distal SPP-FHP/CRC, *KRAS* or *BRAF* mutations were found in the majority of early lesions and thus seem to play a major role in tumor initiation.

We have previously shown that distinct Wnt gene mutations are selected in sporadic and hereditary CRC according to tumor location, i.e. proximal or distal colon (33,42). We and others have also proposed that this finding is the result of the selection of a specific level of β -catenin signaling, optimal for tumor formation, which differs along the colorectum, thus contributing to differences in lesion distribution in specific types of CRC, such as Lynch syndrome (42,43). Proving this, variable gradients in the number of stem cells and physiological Wnt activity have been demonstrated throughout the length of the intestinal tract (44). Thus, in a similar fashion, tumorigenic pathways may also differ between proximal and distal SPP-FHP/CRC.

MGMT and MMR alterations, followed by Wnt gene mutations, are involved in the initiation of proximal/whole-colon SPP-FHP/CRC. The exclusive detection of MMR

gene methylation (mainly of *MSH6*) and/or LOH of D2S123 (flanking *MSH6*) in proximal/whole-colon SPP-FHP/CRC, in the majority of early lesions and in all histological types, appears to suggest that the MMR system plays an important role in the initiation of proximal/whole-colon SPP-FHP/CRC, which is in agreement with the high frequency of MSI (either MSI-L or MSI-H) in these lesions (58%). Interestingly, MMR methylation and, consequently, MMR deficiency were not associated with *MLH1* methylation as has been previously observed in sporadic SE lesions located in the proximal colon (7), but rather with *MSH6* or mutS homolog 3 (*MSH3*) methylation. Accordingly, a high frequency of LOH of the *MSH3* locus has been recently described in sporadic MSI-L CRC, suggesting that the impairment of other MMR genes such as *MSH3* or *MSH6*, as observed in the present study, are involved in an MSI-L pathway, instead of *MLH1*, which is usually associated with the MSI-H serrated pathway (45). The detection of MSI almost exclusively in dinucleotide microsatellite markers is in agreement with this finding, since this type of MSI has been described to be a characteristic feature of MSI-L tumors (46). Interestingly, in the present study, MSI was detected more frequently at D2S123 followed by D17S250.

MGMT methylation and LOH of the *MGMT* locus were the most frequent alterations in proximal/whole-colon SPP-FHP/CRC, and *MGMT* methylation was detected in all HPs, commonly known as the precursor lesion (12). Therefore, we suggest that, similarly to MMR alterations, LOH and methylation of *MGMT* may also be early events in SPP-FHP/CRC proximal/whole-colon tumorigenesis. Notably, among the 17 lesions informative for both MMR and *MGMT* alterations in this form of SPP-FHP/CRC, in 16 (94%) both events were noted (Table IIIA). It is known that *MGMT* deficiency results in the inability to repair O6-methylguanine in the DNA, caused by genotoxic stress, which, once accumulated,

Table V. Molecular characterization of SPP-FHP/CRC lesions stratified by histological type of lesions and the preferential location of the lesions in each patient (proximal/whole-colon or distal).

	HCM	HP	TSA	SSA	SCa	SE lesions	TA	TVA	Ca	AD lesions	SE+AD lesions
Total Wnt gene mutations											
Proximal/whole-colon	-	1/4 (25%)^a	4/4^a	6/7 (86%)^a	1/1	12/16 (75%)^{b,c}	1/6 (17%)	0/2	1/2 (50%)	2/10 (20%)^b	14/26 (54%)
Distal colon	0/1	2/14 (14%)	-	0/2	-	2/17 (12%)^c	1/2 (50%)	-	1/1	2/3 (67%)	4/20 (20%)
Total RAS/RAF gene mutations											
Proximal/whole-colon	-	2/6 (33%)	4/5 (80%)	1/7 (14%)	1/1	8/19 (42%)^d	1/6 (17%)	2/2	1/3 (33%)	4/11 (36%)	12/30 (40%)
Distal colon	1/1	14/14	-	2/2	-	17/17^d	0/2	-	1/1	1/3 (33%)	18/20 (90%)
BRAF gene mutations											
Proximal/whole-colon	-	2/6 (33%)	3/5 (60%)	1/7 (14%)	0/1	6/19 (32%)^e	1/6 (17%)	0/2	0/3	1/11 (9%)	7/30 (23%)
Distal colon	0/1	9/14 (64%)	-	2/2	-	11/17 (65%)^e	0/2	-	1/1	1/3 (33%)	12/20 (60%)
KRAS gene mutations											
Proximal/whole-colon	-	0/6	1/6 (17%)	0/8	1/1	2/21 (10%)^f	0/6	2/2	1/3 (33%)	3/11 (27%)	5/32 (16%)
Distal colon	1/1	5/14 (36%)	-	0/2	-	6/17 (35%)^f	0/2	-	0/1	0/3	6/20 (30%)
MSI											
Proximal/whole-colon	-	1/5 (20%)^g	3/4 (75%)^g	6/7 (86%)^g	0/1	10/17 (59%)^h	3/4 (75%)	1/2 (50%)	1/3 (33%)	5/9 (56%)	15/26 (58%)
Distal colon	0/1	1/10 (10%)	-	0/2	-	1/13 (8%)^h	0/1	-	1/1	1/2 (50%)	2/15 (13%)
MMR gene methylation and/or LOH of D2S123											
Proximal/whole-colon	-	4/4	4/4	2/2	0/1	10/11 (91%)ⁱ	4/4	1/1	2/2	7/7	17/18 (94%)
Distal colon	0/1	0/7	-	0/2	-	0/10ⁱ	0/1	-	-	0/1	0/11
MGMT gene methylation											
Proximal/whole-colon	-	6/6	3/4 (75%)	1/7 (14%)	1/1	11/18 (61%)	4/6 (67%)	2/2	2/3 (67%)	8/11 (73%)	19/29 (65%)
Distal colon	0/1	4/11 (36%)^j	-	2/2	-	6/14 (43%)	0/2	-	1/1	1/3 (33%)	7/17 (41%)
LOH of MGMT locus											
Proximal/whole-colon	-	4/6 (67%)^k	3/4 (75%)	3/5 (60%)	0/1	10/16 (63%)^l	3/4 (75%)	1/1	2/2	6/7 (86%)	16/23 (70%)
Distal colon	-	1/11 (9%)^k	-	0/1	-	1/12 (8%)^l	0/1	-	0/1	0/2	1/14 (7%)

Statistically significant values are shown in bold. ^ap=0.02, Fisher's exact test (2x3). ^bp=0.0091, Fisher's exact test (two-sided). ^cp=0.0011, Fisher's exact test (two-sided). ^dp=1.3x10⁻⁴, Fisher's exact test (two-sided). ^ep=0.049, Fisher's exact test (two-sided). ^fp=0.062, Fisher's exact test (two-sided). ^gp=0.017, Fisher's exact test (2x3). ^hp=0.0049, Fisher's exact test (two-sided). ⁱp=3.12x10⁻⁵, Fisher's exact test (two-sided). ^jp=0.017, Fisher's exact test (two-sided). ^kp=0.027, Fisher's exact test (two-sided). ^lp=0.0045, Fisher's exact test (two-sided). SPP, serrated polyposis; CRC, colorectal cancer; SPP-FHP/CRC, SPP associated with a family history of SPP and/or polyps/CRC (multiple or diagnosed at a young age) in first-degree relatives; MMR, mismatch repair; HCM, hyperplastic colonic mucosa; HP, hyperplastic polyp; TSA, traditional serrated adenoma; TVA, tubulovillous adenoma; TA, tubular adenoma; AD, adenomatous; Ca, carcinoma; SCa, serrated carcinoma; SE, serrated; LOH, loss of heterozygosity.




Figure 1. Proposed pathways for colorectal tumorigenesis in proximal/whole-colon (upper panel) and distal serrated polyposis (SPP) associated with a family history of SPP and/or polyps/colorectal cancer (SPP-FHP/CRC) (bottom panel). Both of these forms may follow a KRAS (alternate) or a BRAF (serrated) pathway, (A and B), respectively, in the upper and lower panels. In addition, in distal SPP, an adenomatous polyposis coli (APC) (traditional) pathway may also occur [(C) in bottom panel]. Each lesion or molecular alteration in these proposed pathways is hypothesized based on the results obtained in the present study and we do not exclude that, in some cases, some of these molecular alterations may not occur, or even that other molecular alterations may also be found. The steps involving lesions that were not analyzed in this study and about which we have previously published information are represented by broken arrows. In some pathways a broken line was used to represent the increase in microsatellite instability (MSI) with tumor progression, to suggest a lower frequency in those cases. Ca, carcinoma; HP, hyperplastic polyp; MMR, mismatch repair; SSA, sessile serrated adenoma; TA, tubular adenoma; TSA, traditional serrated adenoma; TVA, tubulovillous adenoma.

leads to the translocation of the MutS α complex (MSH2 and MSH6) into the nucleus, thus increasing GT mismatch binding activity (47). This may lead to a selective pressure for molecular changes that impair MSH2 or MSH6 function such as promoter hypermethylation or LOH, thus explaining the association between the latter and *MGMT* methylation in the same early SPP-FHP/CRC lesions. Thus, we suggest a primary role for *MGMT* methylation, O6-methylguanine errors and MMR alterations in tumor initiation of proximal/whole-colon SPP-FHP/CRC. We further suggest that this molecular signature may indicate that a germline defect in the mechanisms regulating the response to genotoxic stress underlies the genetic susceptibility in this form of SPP-FHP/CRC.

In the present study, the higher frequency of Wnt gene mutations in proximal SPP-FHP/CRC, particularly in SE lesions, when compared with the AD lesions, suggests that this pathway also plays an important role in this form of SPP-FHP/CRC,

especially in the transition to SE adenoma since this frequency was significantly higher in TSA and SSA than in HP and HCM ($p=8.4 \times 10^{-4}$). In agreement, *CTNNB1* or *AXIN2* mutations, that are selected almost exclusively in proximal colorectal tumors with MSI (33), were detected in 4/14 (28%) TSAs and SSAs from proximal/whole-colon SPP-FHP/CRC. Moreover, among *APC* nonsense and missense mutations, the majority (7/8; 88%) were of the transition type which is a characteristic feature of cells presenting MMR defects (48-50).

The occurrence of LOH of D2S123 and/or D17S250 dinucleotide marker in the present study [17/48 (35%) and 7/51 (14%), respectively], has been previously described in Paneth cell metaplasia, a condition that is commonly observed in the small intestine and the proximal colon of elderly individuals (51,52). Therefore, *MGMT* deficiency may make these cells more exposed to genotoxic stress, thus leading to molecular changes in Wnt genes and consequently to commitment to Paneth cell

lineage [to which Wnt gene mutations largely contribute (53)] and to Paneth cell metaplasia. Therefore, colonic mucosa with Paneth cell metaplasia may be one of the pre-neoplastic lesions in the development of proximal/whole-colon SPP-FHP/CRC.

BRAF and *KRAS* mutations play different roles in proximal and distal SPP-FHP/CRC. Our finding that patients with proximal/whole-colon or distal SPP-FHP/CRC may carry, preferentially, either *KRAS* or *BRAF* mutations, supports previous observations suggesting that different forms of SPP exist, depending on whether lesions follow a *KRAS* or a *BRAF* pathway (22).

In addition, in the present study, we noted distinct roles for these mutations between proximal and distal SPP-FHP/CRC. *BRAF* or *KRAS* mutations were detected in the majority of distal SPP-FHP/CRC lesions, mostly SE early lesions, thus underlining their importance in early stages, whereas in proximal/whole-colon SPP-FHP/CRC these mutations (mainly of *KRAS*) appear to be more important in tumor progression (mostly detected in TSAs, TVAs and Cas). Accordingly, a *KRAS* or alternate pathway has been proposed to be involved in the transition from TSA or TVA to CRC in SPP (10). Indeed, in patients with proximal/whole-colon SPP-FHP/CRC, *KRAS* mutations were found only in TSAs, TVAs or Cas (PH4, PH5 and PH7) which is in accordance with their association with the development of a villous architecture and hence with malignant transformation (10,54-56). Therefore, as TSAs and TVAs shared early somatic events with HPs and TAs, respectively, from the same patients, and based on the model of SPP tumorigenesis previously presented by Leggett and Whitehall (10), we propose that proximal/whole-colon SPP-FHP/CRC tumorigenesis may follow an alternate or *KRAS* pathway, where TVA and TSA may develop from TA and HP, respectively, finally leading to CRC, that may or may not present with MSI (Fig. 1A, upper panel). Alternatively, a serrated or *BRAF* pathway (families PH3 and PH8) where SSA or TSA will probably lead to MSI-H cancer carrying *BRAF* mutations may also occur (Fig. 1B, upper panel). In this model, a deficient DNA repair pathway characterized by *MGMT* and MMR gene methylation and/or LOH followed by Wnt gene mutations, appears to be predominant in proximal SPP-FHP/CRC (Fig. 1, upper panel).

In distal SPP-FHP/CRC, either *KRAS* (PH6 and PH14) or *BRAF* (PH6, PH12, PH19 and PH33) mutations play a major role in tumor initiation, either through an alternate or a serrated pathway (Fig. 1A and B, bottom panel, respectively). Wnt gene mutations and MMR defects were detected in the only carcinoma presented by these patients and thus are likely involved in tumor progression. As PH6 and PH33 also presented TAs and did not present TVAs, we hypothesize that, in distal SPP-FHP/CRC, some AD lesions may also develop through a traditional pathway initiated by *APC* mutations (Fig. 1C, bottom panel).

CRC is more frequent in patients with proximal/whole-colon SPP-FHP/CRC with TSAs, TVAs and KRAS mutations. The association of *KRAS* mutations in TSA or TVA with the development of CRC in proximal/whole-colon SPP-FHP/CRC suggests a higher contribution of the alternate pathway in the development of CRC in patients with SPP-FHP/CRC. Supporting our

hypothesis, *KRAS* mutations have been previously described as more prevalent than *BRAF* mutations in a series of SE carcinomas occurring in the sporadic context, mainly in those presenting adjacent serrated adenomas (51%) (57). Moreover, in the same study, SE carcinomas were frequently MSS and presented a higher frequency of *MGMT* loss compared with traditional carcinomas, which is in agreement with our findings demonstrating that *MGMT* deficiency plays a prominent role in SPP-FHP/CRC, mainly in the proximal colon.

The findings that *KRAS* and *BRAF* mutations promote serrated and hyperplastic features, despite being incapable of initiating colonic adenoma development by themselves (58,59), may contribute to the apparent lower incidence of CRC in distal SPP-FHP/CRC, as according to the results of our present study, *KRAS* or *BRAF* mutations appear to be the initial molecular events in this form. However, additional studies involving more families are warranted.

In conclusion, SPP-FHP/CRC appears to be a distinct clinical and histological entity differing from sporadic SPP. However, we suggest that two forms of SPP-FHP/CRC appear to exist, proximal/whole-colon and distal, which differ mainly in the molecular alterations detected in early lesions. We further propose that a germline defect in the mechanisms regulating the response to genotoxic damage may underlie the genetic susceptibility in the former. In addition, our results suggest that CRC appears to develop more frequently in proximal/whole-colon SPP-FHP/CRC following an alternate *KRAS* pathway, thus underlining the importance of a complete clinical, histological and molecular characterization for CRC risk evaluation in further studies involving families with SPP. The results of these studies may be used to design appropriate guidelines for the clinical management of proximal and distal colonic presentations of SPP that assumes major relevance considering the increased risk of CRC and/or polyps in first-degree relatives.

Acknowledgements

The authors thank all the patients who participated in this study, Dr Inês Francisco for the DNA extraction and Dr Branca Cavaco for the primers for the *NRAS* gene. This study was sponsored and financed by a grant from Núcleo Regional do Sul, Liga Portuguesa Contra o Cancro, Terry Fox (NRS/LPCC-Terry Fox), and Fundação para a Ciência e Tecnologia (FCT), POCI/SAU-OBS/56921/2004.

References

1. Snover DC, Ahnen DJ, Burt RW and Odze RD: Serrated polyps of the colon and rectum and serrated polyposis. In: WHO Classification of Tumours of the Digestive System. Bosman FT, Carneiro F, Hruban RH and Theise ND (eds). IARC, Lyon, pp160-165, 2010.
2. Kalady MF, Jarrar A, Leach B, LaGuardia L, O'Malley M, Eng C and Church JM: Defining phenotypes and cancer risk in hyperplastic polyposis syndrome. *Dis Colon Rectum* 54: 164-170, 2011.
3. Rosty C, Parry S and Young JP: Serrated polyposis: an enigmatic model of colorectal cancer predisposition. *Pathol Res Int* 2011: 157073, 2011.
4. Snover DC, Jass JR, Fenoglio-Preiser C and Batts KP: Serrated polyps of the large intestine: a morphologic and molecular review of an evolving concept. *Am J Clin Pathol* 124: 380-391, 2005.

5. Aust DE and Baretton GB; Members of the Working Group GI-Pathology of the German Society of Pathology: Serrated polyps of the colon and rectum (hyperplastic polyps, sessile serrated adenomas, traditional serrated adenomas, and mixed polyps)-proposal for diagnostic criteria. *Virchows Arch* 457: 291-297, 2010.
6. Rosty C, Hewett DG, Brown IS, Leggett BA and Whitehall VL: Serrated polyps of the large intestine: current understanding of diagnosis, pathogenesis, and clinical management. *J Gastroenterol* 48: 287-302, 2013.
7. Young J and Jass JR: The case for a genetic predisposition to serrated neoplasia in the colorectum: hypothesis and review of the literature. *Cancer Epidemiol Biomarkers Prev* 15: 1778-1784, 2006.
8. Lindor NM: Hereditary colorectal cancer: MYH-associated polyposis and other newly identified disorders. *Best Pract Res Clin Gastroenterol* 23: 75-87, 2009.
9. Roberts A, Nancarrow D, Clendenning M, Buchanan DD, Jenkins MA, Duggan D, Taverna D, McKeone D, Walters R, Walsh MD, *et al*: Linkage to chromosome 2q32.2-q33.3 in familial serrated neoplasia (Jass syndrome). *Fam Cancer* 10: 245-254, 2011.
10. Leggett B and Whitehall V: Role of the serrated pathway in colorectal cancer pathogenesis. *Gastroenterology* 138: 2088-2100, 2010.
11. Jass JR, Iino H, Ruszkiewicz A, Painter D, Solomon MJ, Koorey DJ, Cohn D, Furlong KL, Walsh MD, Palazzo J, *et al*: Neoplastic progression occurs through mutator pathways in hyperplastic polyposis of the colorectum. *Gut* 47: 43-49, 2000.
12. Jass JR, Young J and Leggett BA: Hyperplastic polyps and DNA microsatellite unstable cancers of the colorectum. *Histopathology* 37: 295-301, 2000.
13. Snover DC: Update on the serrated pathway to colorectal carcinoma. *Hum Pathol* 42: 1-10, 2011.
14. Chan AO, Issa JP, Morris JS, Hamilton SR and Rashid A: Concordant CpG island methylation in hyperplastic polyposis. *Am J Pathol* 160: 529-536, 2002.
15. Kambara T, Simms LA, Whitehall VL, Spring KJ, Wynter CV, Walsh MD, Barker MA, Arnold S, McGivern A, Matsubara N, *et al*: BRAF mutation is associated with DNA methylation in serrated polyps and cancers of the colorectum. *Gut* 53: 1137-1144, 2004.
16. Weisenberger DJ, Siegmund KD, Campan M, Young J, Long TI, Faasse NA, Kang GH, Widschwendter M, Weener D, Buchanan D, *et al*: CpG island methylator phenotype underlies sporadic microsatellite instability and is tightly associated with BRAF mutation in colorectal cancer. *Nat Genet* 38: 787-793, 2006.
17. Chow E, Lipton L, Lynch E, D'Souza R, Aragona C, Hodgkin L, Brown G, Winship I, Barker M, Buchanan D, *et al*: Hyperplastic polyposis syndrome: phenotypic presentations and the role of MBD4 and MYH. *Gastroenterology* 131: 30-39, 2006.
18. Yeoman A, Young J, Arnold J, Jass J and Parry S: Hyperplastic polyposis in the New Zealand population: a condition associated with increased colorectal cancer risk and European ancestry. *N Z Med J* 120: U2827, 2007.
19. Boparai KS, Mathus-Vliegen EM, Koornstra JJ, Nagengast FM, van Leerdam M, van Noesel CJ, Houben M, Cats A, van Hest LP, Fockens P and Dekker E: Increased colorectal cancer risk during follow-up in patients with hyperplastic polyposis syndrome: a multicentre cohort study. *Gut* 59: 1094-1100, 2010.
20. Minoo P, Baker K, Goswami R, Chong G, Foulkes WD, Ruszkiewicz AR, Barker M, Buchanan D, Young J and Jass JR: Extensive DNA methylation in normal colorectal mucosa in hyperplastic polyposis. *Gut* 55: 1467-1474, 2006.
21. Wynter CV, Walsh MD, Higuchi T, Leggett BA, Young J and Jass JR: Methylation patterns define two types of hyperplastic polyp associated with colorectal cancer. *Gut* 53: 573-580, 2004.
22. Carvajal-Carmona LG, Howarth KM, Lockett M, Polanco-Echeverry GM, Volikos E, Gorman M, Barclay E, Martin L, Jones AM, Saunders B, *et al*: Molecular classification and genetic pathways in hyperplastic polyposis syndrome. *J Pathol* 212: 378-385, 2007.
23. Guarinos C, Sánchez-Fortún C, Rodríguez-Soler M, Alenda C, Payá A and Jover R: Serrated polyposis syndrome: molecular, pathological and clinical aspects. *World J Gastroenterol* 18: 2452-2461, 2012.
24. Boparai KS, Reitsma JB, Lemmens V, van Os TA, Mathus-Vliegen EM, Koornstra JJ, Nagengast FM, van Hest LP, Keller JJ and Dekker E: Increased colorectal cancer risk in first-degree relatives of patients with hyperplastic polyposis syndrome. *Gut* 59: 1222-1225, 2010.
25. Win AK, Walters RJ, Buchanan DD, Jenkins MA, Sweet K, Frankel WL, de la Chapelle A, McKeone DM, Walsh MD, Clendenning M, *et al*: Cancer risks for relatives of patients with serrated polyposis. *Am J Gastroenterol* 107: 770-778, 2012.
26. Caetano AC, Ferreira H, Soares J, Ferreira A, Gonçalves R and Rolanda C: Phenotypic characterization and familial risk in hyperplastic polyposis syndrome. *Scand J Gastroenterol* 48: 1166-1172, 2013.
27. Hazewinkel Y, Koornstra JJ, Boparai KS, van Os TA, Tytgat KM, van Eeden S, Fockens P and Dekker E: Yield of screening colonoscopy in first-degree relatives of patients with serrated polyposis syndrome. *J Clin Gastroenterol*, 2014.
28. Jaspersion KW, Kanth P, Kirchoff AC, Huismann D, Gammon A, Kohlmann W, Burt RW and Samadder NJ: Serrated polyposis: colonic phenotype, extracolonic features, and familial risk in a large cohort. *Dis Colon Rectum* 56: 1211-1216, 2013.
29. Lanspa SJ, Ahnen DJ and Lynch HT: Serrated polyposis: the last (or only the latest?) frontier of familial polyposis? *Am J Gastroenterol* 107: 779-781, 2012.
30. Albuquerque C, Breukel C, van der Luijt R, Fidalgo P, Lage P, Slors FJ, Leitão CN, Fodde R and Smits R: The 'just-right' signaling model: APC somatic mutations are selected based on a specific level of activation of the beta-catenin signaling cascade. *Hum Mol Genet* 11: 1549-1560, 2002.
31. Miller SA, Dykes DD and Polesky HF: A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 16: 1215, 1988.
32. Francisco I, Albuquerque C, Lage P, Belo H, Vitoriano I, Filipe B, Claro I, Ferreira S, Rodrigues P, Chaves P, *et al*: Familial colorectal cancer type X syndrome: two distinct molecular entities? *Fam Cancer* 10: 623-631, 2011.
33. Albuquerque C, Baltazar C, Filipe B, Penha F, Pereira T, Smits R, Cravo M, Lage P, Fidalgo P, Claro I, *et al*: Colorectal cancers show distinct mutation spectra in members of the canonical WNT signaling pathway according to their anatomical location and type of genetic instability. *Genes Chromosomes Cancer* 49: 746-759, 2010.
34. Liu W, Dong X, Mai M, Seelan RS, Taniguchi K, Krishnadath KK, Halling KC, Cunningham JM, Boardman LA, Qian C, *et al*: Mutations in AXIN2 cause colorectal cancer with defective mismatch repair by activating beta-catenin/TCF signalling. *Nat Genet* 26:146-147, 2000.
35. Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S, Teague J, Woffendin H, Garnett MJ, Bottomley W, *et al*: Mutations of the BRAF gene in human cancer. *Nature* 417: 949-954, 2002.
36. Boland CR, Thibodeau SN, Hamilton SR, Sidransky D, Eshleman JR, Burt RW, Meltzer SJ, Rodriguez-Bigas MA, Fodde R, Ranzani GN and Srivastava S: A National Cancer Institute Workshop on Microsatellite Instability for cancer detection and familial predisposition: development of international criteria for the determination of microsatellite instability in colorectal cancer. *Cancer Res* 58:5248-5257, 1998.
37. Umar A: Lynch syndrome (HNPCC) and microsatellite instability. *Dis Markers* 20:179-180, 2004.
38. Nygren AO, Ameziane N, Duarte HM, Vijzelaar RN, Waisfisz Q, Hess CJ, Schouten JP and Errami A: Methylation-specific MLPA (MS-MLPA): simultaneous detection of CpG methylation and copy number changes of up to 40 sequences. *Nucleic Acids Res* 33: e128, 2005.
39. Gylling A, Ridanpää M, Vierimaa O, Aittomäki K, Avela K, Kääriäinen H, Laivuori H, Pöyhönen M, Sallinen SL, Wallgren-Pettersson C, *et al*: Large genomic rearrangements and germline epimutations in Lynch syndrome. *Int J Cancer* 124: 2333-2340, 2009.
40. Yamada A, Minamiguchi S, Sakai Y, Horimatsu T, Muto M, Chiba T, Boland CR and Goel A: Colorectal advanced neoplasms occur through dual carcinogenesis pathways in individuals with coexisting serrated polyps. *PLoS One* 9: e98059, 2014.
41. Filipe B, Baltazar C, Albuquerque C, Fragoso S, Lage P, Vitoriano I, Mão de Ferro S, Claro I, Rodrigues P, Fidalgo P, *et al*: APC or MUTYH mutations account for the majority of clinically well-characterized families with FAP and AFAP phenotype and patients with more than 30 adenomas. *Clin Genet* 76:242-255, 2009.
42. Albuquerque C, Bakker ER, van Veelen W and Smits R: Colorectal cancers choosing sides. *Biochim Biophys Acta* 1816: 219-231, 2011.
43. Christie M, Jorissen RN, Mouradov D, Sakthianandeswaren A, Li S, Day F, Tsui C, Lipton L, Desai J, Jones IT, *et al*: Different APC genotypes in proximal and distal sporadic colorectal cancers suggest distinct WNT/β-catenin signalling thresholds for tumourigenesis. *Oncogene* 32: 4675-4682, 2013.

44. Leedham SJ, Rodenas-Cuadrado P, Howarth K, Lewis A, Mallappa S, Segditsas S, Davis H, Jeffery R, Rodriguez-Justo M, Keshav S, *et al*: A basal gradient of Wnt and stem-cell number influences regional tumour distribution in human and mouse intestinal tracts. *Gut* 62: 83-93, 2013.
45. Plaschke J, Preussler M, Ziegler A and Schackert HK: Aberrant protein expression and frequent allelic loss of MSH3 in colorectal cancer with low-level microsatellite instability. *Int J Colorectal Dis* 27: 911-919, 2012.
46. Hatch SB, Lightfoot HM Jr, Garwacki CP, Moore DT, Calvo BF, Woosley JT, Sciarrotta J, Funkhouser WK and Farber RA: Microsatellite instability testing in colorectal carcinoma: choice of markers affects sensitivity of detection of mismatch repair-deficient tumors. *Clin Cancer Res* 11: 2180-2187, 2005.
47. Christmann M and Kaina B: Nuclear translocation of mismatch repair proteins MSH2 and MSH6 as a response of cells to alkylating agents. *J Biol Chem* 275: 36256-36262, 2000.
48. Sohn KJ, Choi M, Song J, Chan S, Medline A, Gallinger S and Kim YI: Msh2 deficiency enhances somatic Apc and p53 mutations in Apc^{+/+}Msh2^{-/-} mice. *Carcinogenesis* 24: 217-224, 2003.
49. Oliveira C, Westra JL, Arango D, Ollikainen M, Domingo E, Ferreira A, Velho S, Niessen R, Lagerstedt K, Alhopuro P, *et al*: Distinct patterns of KRAS mutations in colorectal carcinomas according to germline mismatch repair defects and hMLH1 methylation status. *Hum Mol Genet* 13: 2303-2311, 2004.
50. Mark SC, Sandercock LE, Luchman HA, Baross A, Edelmann W and Jirik FR: Elevated mutant frequencies and predominance of G:C to A:T transition mutations in Msh6^(-/-) small intestinal epithelium. *Oncogene* 21: 7126-7130, 2002.
51. Wada R, Yamaguchi T and Tadokoro K: Colonic Paneth cell metaplasia is pre-neoplastic condition of colonic cancer or not? *J Carcinog* 4: 5, 2005.
52. Wada R: Proposal of a new hypothesis on the development of colorectal epithelial neoplasia: nonspecific inflammation - colorectal Paneth cell metaplasia - colorectal epithelial neoplasia. *Digestion* 79 (Suppl 1): 9-12, 2009.
53. Andreu P, Peignon G, Slomianny C, Taketo MM, Colnot S, Robine S, Lamarque D, Laurent-Puig P, Perret C and Romagnolo B: A genetic study of the role of the Wnt/beta-catenin signalling in Paneth cell differentiation. *Dev Biol* 324: 288-296, 2008.
54. Sada M, Mitomi H, Igarashi M, Katsumata T, Saigenji K and Okayasu I: Cell kinetics, p53 and bcl-2 expression, and c-Ki-ras mutations in flat-elevated tubulovillous adenomas and adenocarcinomas of the colorectum: comparison with polypoid lesions. *Scand J Gastroenterol* 34: 798-807, 1999.
55. Maltzman T, Knoll K, Martinez ME, Byers T, Stevens BR, Marshall JR, Reid ME, Einspahr J, Hart N, Bhattacharyya AK, *et al*: Ki-ras proto-oncogene mutations in sporadic colorectal adenomas: relationship to histologic and clinical characteristics. *Gastroenterology* 121: 302-309, 2001.
56. Jass JR, Baker K, Zlobec I, Higuchi T, Barker M, Buchanan D and Young J: Advanced colorectal polyps with the molecular and morphological features of serrated polyps and adenomas: concept of a 'fusion' pathway to colorectal cancer. *Histopathology* 49: 121-131, 2006.
57. García-Solano J, Conesa-Zamora P, Carbonell P, Trujillo-Santos J, Torres-Moreno D D, Pagán-Gómez I, Rodríguez-Braun E and Pérez-Guillermo M: Colorectal serrated adenocarcinoma shows a different profile of oncogene mutations, MSI status and DNA repair protein expression compared to conventional and sporadic MSI-H carcinomas. *Int J Cancer* 131: 1790-1799, 2012.
58. Feng Y, Bommer GT, Zhao J, Green M, Sands E, Zhai Y, Brown K, Burberry A, Cho KR and Fearon ER: Mutant KRAS promotes hyperplasia and alters differentiation in the colon epithelium but does not expand the presumptive stem cell pool. *Gastroenterology* 141: 1003-1013.e1-10, 2011.
59. Carragher LA, Snell KR, Giblett SM, Aldridge VS, Patel B, Cook SJ, Winton DJ, Marais R and Pritchard CA: V600EBraf induces gastrointestinal crypt senescence and promotes tumour progression through enhanced CpG methylation of p16INK4a. *EMBO Mol Med* 2: 458-471, 2010.