

# Effects of all-trans retinoic acid on VEGF and HIF-1 $\alpha$ expression in glioma cells under normoxia and hypoxia and its anti-angiogenic effect in an intracerebral glioma model

CHEN LIANG<sup>1</sup>, SHIWEN GUO<sup>1</sup> and LING YANG<sup>2</sup>

<sup>1</sup>Department of Neurosurgery, First Affiliated Hospital, Medical College of Xi'an Jiaotong University, Xi'an, Shaanxi 710061; <sup>2</sup>Department of Aeromedical Physical Examination, Xi'an Civil Aviation Hospital, Xi'an, Shaanxi 710082, P.R. China

Received June 19, 2013; Accepted May 21, 2014

DOI: 10.3892/mmr.2014.2543

**Abstract.** All-trans retinoic acid (ATRA) is one of the most potent inducers of differentiation and is capable of inducing differentiation and apoptosis in glioma cells. However, the effect of ATRA on glioma angiogenesis is yet to be elucidated. The present study investigated the effects of ATRA on the expression of vascular endothelial growth factor (VEGF) and hypoxia-inducible factor-1 $\alpha$  (HIF-1 $\alpha$ ) in various glioma cell lines under normoxia and hypoxia. The effect of ATRA on angiogenesis in a rat intracerebral glioma model was also investigated, with the aim of revealing the effect of ATRA on glioma angiogenesis. In the present study, U-87 MG and SHG44 glioma cells were treated with ATRA at various concentrations (0, 5, 10, 20 and 40  $\mu$ mol/l) under normoxia or hypoxia. Quantitative polymerase chain reaction and western blot analysis were used to investigate VEGF and HIF-1 $\alpha$  mRNA and protein expression, respectively. An intracerebral glioma model was generated using intracerebral implantation of C6 glioma cells into rats. Tumor-bearing rats were treated with ATRA at different doses (0, 5 and 10 mg/kg/day) for two weeks, and immunohistochemical assays were performed to detect the cluster of differentiation 34-positive cells in order to evaluate the microvessel density (MVD) in each group. Following ATRA treatment, the expression of VEGF and HIF-1 $\alpha$  was found to vary among the different concentration groups. In the glioma cells in the lower concentration groups (5 and 10  $\mu$ mol/l ATRA), a significant increase in VEGF and HIF-1 $\alpha$  expression was observed. Conversely, a significant decrease in VEGF and HIF-1 $\alpha$  expression was found

in the glioma cells in the high ATRA concentration group (40  $\mu$ mol/l), compared with that in the cells in the control group. Furthermore, in the rat intracerebral glioma model, ATRA decreased glioma MVD, particularly in the high-dose group (10 mg/kg/day), compared with the control group. These results suggest that ATRA may exhibit a dose-dependent effect on glioma angiogenesis and may inhibit glioma angiogenesis *in vivo*.

## Introduction

Glioma is one of the most common types of brain tumor. Despite the development of current standard therapy, the prognosis of patients with malignant glioma is poor. All-trans retinoic acid (ATRA), a physiologically active derivative of vitamin A, is one of the most potent inducers of differentiation and has been extensively studied. The biological effects of ATRA are mediated through retinoic acid receptors (RARs) and retinoic X receptors. Several studies have shown that ATRA can induce differentiation and apoptosis in a variety of glioma cells (1,2), including glioma stem cells (3,4), a highly tumorigenic and therapy-resistant tumor subpopulation. ATRA can also induce growth arrest in glioma cells (5). Furthermore, studies have revealed that ATRA exhibits certain additional anti-glioma effects (6-11). Li *et al* (6) reported that ATRA enhanced the tumoricidal effect of suicide-gene therapy against medulloblastoma. Other studies have found that ATRA can significantly enhance the anti-tumor effect of certain chemoimmunotherapy and cytotoxic chemotherapy drugs, including interferon- $\gamma$  and taxol, respectively, on glioma (7-11). These findings indicate that ATRA may have therapeutic potential in patients with glioma.

Angiogenesis is a complex biological process, which involves the degradation of the basement membrane and extracellular matrix, endothelial cell (EC) proliferation, migration and tube formation (12). Angiogenesis is mediated by various regulatory factors, including vascular endothelial growth factor (VEGF), basic fibroblast growth factor, hepatocyte growth factor (HGF) and angiopoietin-1 and -2. These regulatory factors are well-established angiogenic factors, and their biological effects are mediated through interaction

---

*Correspondence to:* Professor Shiwen Guo, Department of Neurosurgery, First Affiliated Hospital, Medical College of Xi'an Jiaotong University, 277 Yanta West Road, Xi'an, Shaanxi 710061, P.R. China  
E-mail: gsw1962@126.com

**Key words:** all-trans retinoic acid, vascular endothelial growth factor, glioma, hypoxia-inducible factor-1 $\alpha$ , angiogenesis

with their membrane receptors. Angiogenesis is important in numerous physiological processes, including embryo development, wound healing and normal growth (13). However, angiogenesis also has a key role in various pathological processes, such as inflammation (14), diabetic retinopathy (15) and the formation, development and recurrence of glioma (16). Therefore, anti-angiogenesis therapy has been developed as a novel therapeutic strategy for patients with glioma. Numerous studies have shown that anti-angiogenesis therapy can significantly inhibit glioma growth (17,18) and improve the outcome of patients with glioma (19,20). However, the effect of ATRA on glioma angiogenesis is yet to be elucidated.

VEGF induces EC proliferation, promotes cell migration and inhibits apoptosis. VEGF is well established as a key regulator of angiogenesis and plays an important role in the formation, development and recurrence of glioma (21-24). Hypoxia-inducible factor-1 $\alpha$  (HIF-1 $\alpha$ ) is a transcription factor that is responsible for the induction of genes that regulate numerous biological processes, including angiogenesis (25). Furthermore, HIF-1 $\alpha$  regulates various genes involved in the different stages of angiogenesis, including VEGF. VEGF is the most potent endothelial-specific mitogen and directly participates in angiogenesis by recruiting ECs and stimulating their proliferation (26).

The present study investigated the effect of ATRA on angiogenesis by detecting VEGF and HIF-1 $\alpha$  expression in two glioma cell lines, U-87 MG (U87) and SHG44, under normoxia and hypoxia. The anti-angiogenic effect of ATRA in a rat intracerebral glioma model was also investigated. The results of this study are likely to provide an enhanced understanding of the mechanisms underlying the therapeutic effect of ATRA in malignant glioma.

## Materials and methods

**Materials.** The U87 and SHG44 human glioma cell lines and the C6 rat glioma cell line were purchased from the Cell Resource Center, Chinese Academy of Sciences (Shanghai, China). A total of 30 male, specific pathogen-free, Sprague Dawley rats, weighing between 280 and 320 g, were obtained from the Laboratory Animal Center, Medical College of Xi'an Jiaotong University (Xi'an, China). All animal procedures were performed in accordance with the Guidance by the Research Ethics Committee of the Medical College of Xi'an Jiaotong University.

**Cell culture.** U87 cells were cultured in Dulbecco's Modified Eagle's Medium (Hyclone, Beijing, China), supplemented with 10% fetal bovine serum (FBS; Hyclone) in 5% CO<sub>2</sub> at 37°C. SHG44 and C6 cells were cultured in RPMI-1640 (Hyclone), supplemented with 10% FBS in 5% CO<sub>2</sub> at 37°C.

**Quantitative polymerase chain reaction (qPCR) analysis.** U87 and SHG44 cells were seeded onto six-well plates (Corning Inc., Lowell, MA, USA) at a density of 3x10<sup>5</sup> cells/well. ATRA (Sigma-Aldrich, St. Louis, MO, USA) was dissolved in dimethyl sulfoxide (DMSO; Sigma-Aldrich) and stored in light-protected vials at -20°C as a stock solution. The stock solution was diluted to the desired concentration prior to use. All experiments were performed under low-light conditions to

minimize ATRA photoisomerization. The cells were incubated in media containing various concentrations of ATRA (5, 10, 20 and 40  $\mu$ mol/l) for 24 h under normoxia. The control group was treated with an equal volume of the solvent (DMSO) in the culture media.

CoCl<sub>2</sub>•6H<sub>2</sub>O (Sigma-Aldrich) was used to simulate hypoxia in order to investigate the effect of ATRA on VEGF and HIF-1 $\alpha$  expression under hypoxic conditions. U87 and SHG44 cells were seeded in six-well plates at a density of 3x10<sup>5</sup> cells/well and were incubated in media containing various concentrations of ATRA (5, 10, 20 and 40  $\mu$ mol/l) and 100  $\mu$ mol/l CoCl<sub>2</sub> for 24 h. The hypoxia control group was treated with an equal volume of the solvent (DMSO) and 100  $\mu$ mol/l CoCl<sub>2</sub> in the culture media. The normoxia control group was solely treated with an equal volume of the solvent (DMSO).

The cells were lysed and the total RNA was isolated using the RNeasy200 kit (Shanghai Fastagen Biotechnology Co., Ltd., Shanghai, China) according to the manufacturer's instructions. The RNA was reverse transcribed using PrimeScript™ RT Master Mix (Takara Bio Inc., Dalian, China). qPCR was performed using SYBR® Premix Ex Taq™ II (Takara Bio Inc.) with a Bio-Rad IQ5 thermal cycler (Bio-Rad Laboratories, Inc., Hercules, CA, USA) and analyzed with Bio-Rad iQ5 software version 2.0. Gene expression was compared using the  $\Delta$ cycle threshold ( $\Delta$ Ct) method ( $\Delta$ Ct=Ct<sub>target</sub>-Ct <sub>$\beta$ -actin</sub>), where  $\beta$ -actin expression was used as an endogenous reference gene. The changes in target gene expression were evaluated using the 2<sup>- $\Delta$ Ct</sup> method (27). All the primers were designed and synthesized by Takara Bio Inc. (Table I).

**Western blot analysis.** U87 and SHG44 cells were cultured in 25-ml culture flasks (Corning Inc.) with ATRA at various concentrations under normoxia or hypoxia for 24 h, in accordance with the aforementioned methods. The cells were then harvested for the subsequent assays. The cells were washed twice with phosphate-buffered saline (PBS), then scraped on ice in 300  $\mu$ l radioimmunoprecipitation assay lysis buffer (Beyotime Institute of Biotechnology, Haimen, China) with 1 mmol/l phenylmethylsulfonyl fluoride. The lysates were cleared of insoluble material using centrifugation and protein concentration was determined using a Bradford protein assay kit (Beyotime Institute of Biotechnology). The samples were boiled in 1X SDS-PAGE sample loading buffer, resolved using SDS-PAGE and transferred to polyvinylidene fluoride membranes (Bio-Rad Laboratories, Inc.). The membranes were probed with anti-VEGF (Epitomics, Burlingame, CA, USA) and anti-GAPDH (Bioworld Technology Inc., St. Louis Park, MN, USA) antibodies diluted 1:1,000 and 1:5,000, respectively. Membranes from the hypoxia groups were also probed with anti-HIF-1 $\alpha$  antibodies (Cell Signaling Technology, Inc., Beverly, MA, USA) diluted 1:1,000. Subsequent to washing in Tris-buffered saline containing 0.02% Tween 20, the membranes were incubated with a secondary polyclonal anti-rabbit immunoglobulin G (IgG) antibody conjugated to horseradish peroxidase (Thermo Fisher Scientific, Inc., Waltham, MA, USA) diluted 1:2,000. Membranes were developed using Super Signal™ West Pico chemiluminescent reagent (Pierce Biotechnology, Inc., Rockford, IL, USA).

Table I. Primer sequences for the quantitative polymerase chain reaction.

| Gene           | Sequence   | Product length (bp) |
|----------------|--|---------------------|
| HIF-1 $\alpha$ | Forward, 5'-TCTGGGTTGAAACTCAAGCAACTG-3'<br>Reverse, 5'-CAACCGGTTTAAGGACACATTCTG-3' | 150                 |
| VEGF           | Forward, 5'-TCACAGGTACAGGGATGAGGACAC-3'<br>Reverse, 5'-CAAAGCACAGCAATGTCCTGAAG-3'  | 72                  |
| $\beta$ -actin | Forward, 5'-TGGCACCCAGCACAATGAA-3'<br>Reverse, 5'-CTAAGTCATAGTCCGCCTAGAAGCA-3'     | 186                 |

HIF-1 $\alpha$ , hypoxia-inducible factor-1 $\alpha$ ; VEGF, vascular endothelial growth factor.

**Animal studies.** A total of 10  $\mu$ l cell suspension containing  $1 \times 10^6$  C6 cells was injected at the coronal suture 3 mm away from the midline and at a depth of 5 mm into the right frontal lobe of the rats. Seven days after C6 cell injection, 30 tumor-bearing rats were randomly divided into three groups (10 rats/group). The low- and high-dose groups were treated with different doses of ATRA (5 and 10 mg/kg/day, respectively) diluted in corn oil. ATRA was administered using intraperitoneal injection between days 8 and 21. The control group was treated with an equal volume of solvent.

On day 22, after two weeks of treatment, all tumor-bearing rats were sacrificed. Brain and tumor samples were obtained, and frozen sections were prepared. Immunohistochemistry was performed to detect the cluster of differentiation (CD) 34-positive cells in order to evaluate the microvessel density (MVD) in each group using an immunohistochemical assay kit (Boster Biological Technology Co., Ltd., Wuhan, China). All procedures were performed according to the manufacturer's instructions. Briefly, the sections were fixed in 4% paraformaldehyde for 20 min and 3% hydrogen peroxide in methanol was then used to quench any endogenous peroxidase activity. Non-specific protein binding was blocked using 5% bovine serum albumin for 10 min at room temperature. The sections were incubated overnight with rabbit anti-CD34 antibody (Boster Biological Technology Co., Ltd.) diluted 1:100 at 4°C. The negative control sections were incubated with PBS instead of the antibody. Subsequent to three washes in PBS, the sections were incubated in biotinylated anti-rabbit IgG antibody at 3°C for 30 min. Following incubation, the sections were washed three times for 2 min in PBS and antibody location was determined using a 3,3'-diaminobenzidine substrate kit (Tiangen Biotech Co., Ltd., Beijing, China) for 5 min. Normal vascular endothelium was used as a positive control. MVD was determined as previously described by Weidner *et al* (28). Briefly, the tumor sections were scanned at low magnifications (x40 or x100) to determine the areas of most intense tumor angiogenesis, termed the 'hot spots'. Following 'hot spot' identification, the MVD was calculated by averaging the number of individual microvessels in five fields at high magnification (x400).

**Statistical analysis.** Values are presented as the mean  $\pm$  standard deviation and data were analyzed using SPSS 17.0 software (SPSS, Inc., Chicago, IL, USA). One-way analysis of variance was used to compare the groups and least significant difference tests were performed for further inter-group

comparisons. A value of  $P < 0.05$  was considered to indicate a statistically significant difference.

## Results

**Effects of ATRA on the VEGF and HIF-1 $\alpha$  expression in glioma cells under normoxia.** Following incubation with various concentrations of ATRA for 24 h, VEGF mRNA levels were observed to vary among the different groups. As shown in Fig. 1A, after 24 h the VEGF mRNA expression was observed to be significantly upregulated in the U87 and SHG44 glioma cells in the lower ATRA concentration groups (5 and 10  $\mu$ mol/l), compared with that in the control group ( $P < 0.01$ ). Treatment with 5  $\mu$ mol/l ATRA was found to increase VEGF mRNA expression ~2.4-fold in the U87 and 1.3-fold in the SHG44 glioma cells. Treatment with 10  $\mu$ mol/l ATRA increased VEGF mRNA expression ~4.42-fold in the U87 and 2.24-fold in the SHG44 glioma cells. Conversely, following treatment with 40  $\mu$ mol/l ATRA, the VEGF mRNA expression was observed to be significantly downregulated to 0.80- and 0.77-fold that in the corresponding control groups, in the U87 ( $P < 0.01$ ) and SHG44 ( $P < 0.05$ ) glioma cells, respectively. Furthermore, following treatment with 20  $\mu$ mol/l ATRA, VEGF mRNA levels in the U87 glioma cells increased 1.53-fold relative to the levels in the control group ( $P < 0.01$ ); however, no significant difference was observed in the SHG44 glioma cells.

VEGF protein expression was analyzed using western blot analysis, as shown in Fig. 1B and C. In accordance with the changes in VEGF mRNA expression, treatment with lower concentrations of ATRA (5 and 10  $\mu$ mol/l) was observed to upregulate VEGF protein expression in the U87 and SHG44 glioma cells. Furthermore, treatment with high ATRA concentrations (40  $\mu$ mol/l) significantly inhibited VEGF expression in the glioma cell lines.

HIF-1 $\alpha$  mRNA expression was also analyzed in the U87 and SHG44 glioma cells following ATRA treatment. As shown in Fig. 1D, qPCR revealed that, following treatment with lower concentrations of ATRA (5 and 10  $\mu$ mol/l), HIF-1 $\alpha$  mRNA expression was upregulated in the glioma cell lines. Treatment with 5  $\mu$ mol/l ATRA was observed to upregulate HIF-1 $\alpha$  mRNA expression ~1.41- and 1.24-fold in the U87 and SHG44 glioma cells, respectively. Furthermore, treatment with 10  $\mu$ mol/l ATRA was found to upregulate HIF-1 $\alpha$  mRNA expression ~2.2- and 2.5-fold in the U87 and SHG44 glioma






Figure 1. Effects of ATRA on VEGF and HIF-1 $\alpha$  expression under normoxia. (A and D) Effects of ATRA on (A) VEGF and (D) HIF-1 $\alpha$  mRNA expression. U87 and SHG44 glioma cells were treated with various concentrations of ATRA (5, 10, 20 and 40  $\mu$ mol/l) for 24 h under normoxia. The control group was treated with an equal volume of solvent (dimethyl sulfoxide) in the culture media. VEGF and HIF-1 $\alpha$  mRNA expression increased following treatment with 5 and 10  $\mu$ mol/l ATRA in the two glioma cell lines ( $P < 0.01$  versus the control group). Following treatment with 40  $\mu$ mol/l ATRA, the VEGF and HIF-1 $\alpha$  mRNA expression was significantly downregulated ( $P < 0.01$  versus the control group). (B and C) Western blot analysis of VEGF protein expression in (B) U87 and (C) SHG44 glioma cells treated with various concentrations of ATRA. (a) Control; (b) 5, (c) 10, (d) 20 and (e) 40  $\mu$ mol/l ATRA. \* $P < 0.05$  and \*\* $P < 0.01$  versus the control group. ATRA, all-trans retinoic acid; VEGF, vascular endothelial growth factor; HIF-1 $\alpha$ , hypoxia-inducible factor-1 $\alpha$ .

cells, respectively ( $P < 0.01$  versus the control group). However, treatment with 40  $\mu$ mol/l ATRA was observed to significantly decrease HIF-1 $\alpha$  mRNA expression in glioma cells to 0.71- and 0.55-fold that in the control group in the U87 and SHG44 glioma cells, respectively ( $P < 0.01$ ). HIF-1 $\alpha$  protein is rapidly degraded under normoxic conditions; therefore, HIF-1 $\alpha$  protein expression was not detected under normoxia.

**Effects of ATRA on VEGF and HIF-1 $\alpha$  expression in glioma cells under hypoxia.** Hypoxia is one of the primary regulators of angiogenesis; therefore, the effects of ATRA on VEGF and HIF-1 $\alpha$  expression was analyzed under hypoxia in the U87 and

SHG44 cells. As shown in Fig. 2, hypoxia was observed to significantly upregulate VEGF protein and mRNA expression ( $P < 0.01$  versus the normoxia control group). However, hypoxia only increased HIF-1 $\alpha$  protein expression, with HIF-1 $\alpha$  mRNA expression unaffected by the hypoxia mimic ( $P > 0.05$ ).

Under hypoxia, the U87 and SHG44 glioma cells treated with various concentrations of ATRA exhibited a similar reaction to those under normoxia. VEGF and HIF-1 $\alpha$  mRNA and protein expression was significantly upregulated by lower concentrations of ATRA (5 and 10  $\mu$ mol/l), compared with the hypoxia control group ( $P < 0.01$ ). However, treatment with high concentrations of ATRA (40  $\mu$ mol/l) was observed to decrease the expression of VEGF and HIF-1 $\alpha$  ( $P < 0.01$ ).

**Effect of ATRA on MVD in the rat intracerebral glioma model.** In order to investigate the anti-angiogenic effect of ATRA on glioma *in vivo*, CD34 was examined using immunohistochemistry to assess glioma MVD in each group of tumor-bearing rats. As shown in Fig. 3, ATRA was found to significantly decrease the number of CD34-stained glioma microvessels, particularly at higher doses (10 mg/kg/day) ( $P < 0.01$  versus the control group). The average MVDs were  $106.56 \pm 17.80$ ,  $68.04 \pm 11.95$  and  $33.66 \pm 12.05$  in the control, low- (5 mg/kg/day) and high- (10 mg/kg/day) dose groups, respectively.

## Discussion

It is well established that VEGF is a key regulator of angiogenesis and has an important role in the formation, development and recurrence of glioma. Several studies have reported that ATRA affects VEGF expression; however, the results are controversial (29-34). Reports have demonstrated that ATRA is capable of inhibiting VEGF expression in several cell types, including human gastric cancer (29), esophageal squamous cell carcinoma (30) and leukemia (31) cells. However, ATRA has been reported to have a stimulatory effect on VEGF expression in various other cell types, such as human umbilical vein ECs (32), human retinal pigment epithelial cells (33) and retinoblastoma Y78 cells (34). The only consensus regarding the effect of ATRA on VEGF expression is that the anti-leukemia effect of ATRA is exerted through the downregulation of VEGF (31,35-37). The contradictory results produced by these studies may be a consequence of differences in cell type, methodology, experimental conditions and the concentrations of ATRA. However, little is known concerning the effect of ATRA on VEGF expression in glioma.

In the present study, ATRA was observed to have a dose-dependent effect on VEGF expression in glioma cells. Lower concentrations of ATRA were found to significantly increase VEGF expression at the transcriptional and translational level in glioma cells. However, high concentrations of ATRA were observed to significantly decrease the expression of VEGF mRNA and protein. These converse effects have also been reported in ECs. Saito *et al* (32) reported that, at concentrations between 1 nmol/l and 1  $\mu$ mol/l, ATRA induced the expression of VEGF and its receptor, VEGFR-2, in ECs. However, it has also been reported that 5  $\mu$ mol/l ATRA decreases VEGF and VEGFR-2 expression, as well as Akt phosphorylation in ECs (38). In addition to the differences in




Figure 2. Effects of ATRA on VEGF and HIF-1 $\alpha$  expression under hypoxia. (A and C) Effect of hypoxia mimic on (A) VEGF and (C) HIF-1 $\alpha$  mRNA expression. Hypoxia was simulated using 100  $\mu$ mol/l CoCl<sub>2</sub>. The mRNA expression of VEGF, but not of HIF-1 $\alpha$ , was induced by the hypoxia mimic. (B and D) Effects of ATRA on (B) VEGF and (D) HIF-1 $\alpha$  mRNA expression under hypoxia. U87 and SHG44 glioma cells were treated with various concentrations of ATRA (5, 10, 20 and 40  $\mu$ mol/l) for 24 h under hypoxia. The hypoxia control group was treated with an equal volume of solvent (dimethyl sulfoxide) and 100  $\mu$ mol/l CoCl<sub>2</sub> in the culture media. The VEGF and HIF-1 $\alpha$  mRNA expression increased following treatment with 5 and 10  $\mu$ mol/l ATRA in the two glioma cell lines ( $P < 0.01$  versus the hypoxia control group). Following treatment with 40  $\mu$ mol/l ATRA, the VEGF and HIF-1 $\alpha$  mRNA expression was significantly downregulated ( $P < 0.01$  versus the hypoxia control group). (E and F) Western blot analysis of VEGF and HIF-1 $\alpha$  protein expression in (E) U87 and (F) SHG44 glioma cells treated with various concentrations of ATRA. (a) Normoxia control; (b) hypoxia control; (c) 5, (d) 10, (e) 20 and (f) 40  $\mu$ mol/l ATRA. \*\* $P < 0.01$  versus the hypoxia control group; ## $P < 0.01$  versus the normoxia control group. ATRA, all-trans retinoic acid; VEGF, vascular endothelial growth factor; HIF-1 $\alpha$ , hypoxia-inducible factor-1 $\alpha$ .

methodology and experimental conditions in these two studies, the different concentrations of ATRA may have contributed to the contradictory results. The present study revealed that ATRA has a concentration-dependent effect on VEGF mRNA and protein expression in glioma cells.

The mechanism underlying the ATRA-induced regulation of VEGF expression is yet to be elucidated. Evidence suggests that ATRA increases the protein expression of specificity protein (Sp) 1 through post-transcriptional mechanisms, and that elevated Sp1 protein levels induce VEGF expression (34). It has also been shown that ATRA is capable of directly inducing VEGF expression through the RAR signaling pathway (32). However, it has additionally been suggested that ATRA may decrease VEGF gene transcription through a direct repeat 1 element located at the transcription initiation site (31). Furthermore, HIF-1 $\alpha$  is well established as a key regulator of VEGF expression. Studies have indicated that ATRA regulates VEGF expression through the regulation of HIF-1 $\alpha$  (39,40). Therefore, in the present study, the effect of ATRA on HIF-1 $\alpha$  expression in glioma cells was investigated.

The effect of ATRA on HIF-1 $\alpha$  mRNA expression was observed to be similar to that on VEGF expression. Lower concentrations of ATRA significantly increased HIF-1 $\alpha$

expression and high concentrations of ATRA significantly decreased HIF-1 $\alpha$  expression. HIF-1 $\alpha$  protein is rapidly degraded under normoxic conditions and is undetectable; therefore, hypoxia was mimicked using CoCl<sub>2</sub>, and the effect of ATRA on VEGF and HIF-1 $\alpha$  expression, particularly HIF-1 $\alpha$  protein expression, was investigated in glioma cells under hypoxia.

Hypoxia can induce angiogenesis in numerous physiological and pathological processes, primarily through the HIF pathway. In the present study, mimicking hypoxia was found to significantly increase HIF-1 $\alpha$  protein, but not mRNA, expression. This phenomenon is associated with the mechanism underlying CoCl<sub>2</sub>-mediated hypoxia mimicry (41). Briefly, cobalt displaces iron from the iron-binding site of the enzyme, thus interrupting with the degradation of the HIF-1 $\alpha$  protein under normoxia. In the present study, following treatment with different concentrations of ATRA under hypoxia, similar dose-dependent changes in HIF-1 $\alpha$  and VEGF protein expression were observed. Thus, there is reason to speculate that the regulation of VEGF expression by ATRA in glioma cells, may partially be through the regulation of HIF-1 $\alpha$  expression.

The mechanism underlying the regulation of HIF-1 $\alpha$  expression by ATRA remains unclear. It has been suggested




Figure 3. Effect of ATRA on MVD in a rat intracerebral glioma model. (A) Rat intracerebral glioma model (black arrow shows the tumor). (B) Immunohistochemical staining of cluster of differentiation 34 in the (a) control group, (b) low-dose group treated with 5 mg/kg/day ATRA and (c) high-dose group treated with 10 mg/kg/day ATRA (magnification, x100). (C) MVD in each group. Following treatment with ATRA, the glioma MVD was significantly decreased, particularly in the high-dose group. \*\* $P < 0.01$  versus the control group. ATRA, all-trans retinoic acid; MVD, microvessel density.

that the ATRA-induced upregulation of HIF-1 $\alpha$  expression may involve the stabilization of HIF-1 $\alpha$  mRNA (39). It has also been reported that ATRA can induce HIF-1 $\alpha$  expression through intracrine prostaglandin E<sub>2</sub> signaling (40). However, the mechanisms underlying the inhibitory effect of high concentrations of ATRA on HIF-1 $\alpha$  expression are yet to be elucidated, and further research is required.

In the present study, a rat intracerebral glioma model was generated in order to investigate the effect of ATRA on the inhibition of glioma angiogenesis *in vivo*. ATRA has been reported to inhibit angiogenesis in solid tumors, including esophageal squamous cell carcinoma (30). In the present study, ATRA was found to significantly decrease glioma MVD, particularly in the high-dose group.

The results of the present study suggest that, despite the increased expression of VEGF induced by lower concentrations of ATRA in glioma cells *in vitro*, ATRA has strong anti-angiogenic effects on glioma *in vivo*. However, the mechanisms underlying the anti-angiogenic effects of ATRA on glioma *in vivo* are yet to be fully elucidated. The findings of the present study suggest that the inhibitory effects of high

concentrations of ATRA on VEGF and HIF-1 $\alpha$  expression are likely to be associated with the anti-angiogenic mechanisms. Furthermore, ATRA has been reported to significantly inhibit the secretion and expression of HGF in glioma cells (42). HGF is an important pro-angiogenic cytokine; therefore, the inhibition of HGF may contribute to these anti-angiogenic effects *in vivo*. However, ATRA-mediated differentiation has also been shown to reduce angiogenesis in stem-like glioma cells *in vivo* (4). Therefore, the mechanisms underlying the anti-angiogenic effect of ATRA *in vivo* are likely to be complex and require further investigation.

VEGF has an important role in angiogenesis. The upregulation of VEGF expression in glioma cells induced by lower concentrations of ATRA may reduce its therapeutic effects on glioma *in vivo*. Therefore, as a therapeutic strategy, ATRA has certain limitation; the combination of ATRA with therapies targeting VEGF may have greater beneficial effects for the treatment of glioma.

In conclusion, in the present study, ATRA was found to have a concentration-dependent effect on the expression of VEGF and HIF-1 $\alpha$ . Furthermore, ATRA was observed to



inhibit glioma angiogenesis *in vivo*; however, further research is required in order to reveal its mechanisms.

## References

1. Zang CB, Wächter M, Liu H, *et al*: Ligands for PPARgamma and RAR cause induction of growth inhibition and apoptosis in human glioblastomas. *J Neurooncol* 65: 107-118, 2003.
2. Tang K, Cao L, Fan SQ, *et al*: Effect of all-trans-retinoic acid on C6 glioma cell proliferation and differentiation. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 33: 892-897, 2008 (In Chinese).
3. Karsy M, Albert L, Tobias ME, Murali R and Jhanwar-Uniyal M: All-trans retinoic acid modulates cancer stem cells of glioblastoma multiforme in an MAPK-dependent manner. *Anticancer Res* 30: 4915-4920, 2010.
4. Campos B, Wan F, Farhadi M, *et al*: Differentiation therapy exerts antitumor effects on stem-like glioma cells. *Clin Cancer Res* 16: 2715-2728, 2010.
5. Chang Q, Chen Z, You J, *et al*: All-trans-retinoic acid induces cell growth arrest in a human medulloblastoma cell line. *J Neurooncol* 84: 263-267, 2007.
6. Li S, Gao Y, Pu K, Ma L, Song X and Liu Y: All-trans retinoic acid enhances bystander effect of suicide-gene therapy against medulloblastomas. *Neurosci Lett* 503: 115-119, 2011.
7. Das A, Banik NL and Ray SK: Molecular mechanisms of the combination of retinoid and interferon-gamma for inducing differentiation and increasing apoptosis in human glioblastoma T98G and U87MG Cells. *Neurochem Res* 34: 87-101, 2009.
8. Haque A, Banik NL and Ray SK: Emerging role of combination of all-trans retinoic acid and interferon-gamma as chemotherapeutic in the management of human glioblastoma. *Neurochem Res* 32: 2203-2209, 2007.
9. Zhang R, Banik NL and Ray SK: Combination of all-trans retinoic acid and interferon-gamma upregulated p27(kip1) and down regulated CDK2 to cause cell cycle arrest leading to differentiation and apoptosis in human glioblastoma LN18 (PTEN-proficient) and U87MG (PTEN-deficient) cells. *Cancer Chemother Pharmacol* 62: 407-416, 2008.
10. Karmakar S, Banik NL, Patel SJ and Ray SK: Combination of all-trans retinoic acid and taxol regressed glioblastoma T98G xenografts in nude mice. *Apoptosis* 12: 2077-2087, 2007.
11. Karmakar S, Banik NL and Ray SK: Combination of all-trans retinoic acid and paclitaxel-induced differentiation and apoptosis in human glioblastoma U87MG xenografts in nude mice. *Cancer* 112: 596-607, 2008.
12. Distler JH, Hirth A, Kurowska-Stolarska M, Gay RE, Gay S and Distler O: Angiogenic and angiostatic factors in the molecular control of angiogenesis. *Q J Nucl Med* 47: 149-161, 2003.
13. Folkman J and Klagsbrun M: Angiogenic factors. *Science* 235: 442-447, 1987.
14. Muramatsu R, Takahashi C, Miyake S, Fujimura H, Mochizuki H and Yamashita T: Angiogenesis induced by CNS inflammation promotes neuronal remodeling through vessel-derived prostacyclin. *Nat Med* 18: 1658-1664, 2012.
15. Crawford TN, Alfaro DV III, Kerrison JB and Jablon EP: Diabetic retinopathy and angiogenesis. *Curr Diabetes Rev* 5: 8-13, 2009.
16. Würdinger T and Tannous BA: Glioma angiogenesis: Towards novel RNA therapeutics. *Cell Adh Migr* 3: 230-235, 2009.
17. Chen H, Fan K, Wang S, Liu Z and Zheng Z: Dual targeting of glioma U251 cells with nanoparticles prevents tumor angiogenesis and inhibits tumor growth. *Curr Neurovasc Res* 9: 133-138, 2012.
18. Guo SW, Che HM and Li WZ: Anti-tumor effect of lentivirus-mediated gene transfer of alaphostatin on human glioma. *Cancer Sci* 102: 1038-1044, 2011.
19. Vredenburgh JJ, Desjardins A, Herndon JE II, *et al*: Phase II trial of bevacizumab and irinotecan in recurrent malignant glioma. *Clin Cancer Res* 13: 1253-1259, 2007.
20. Kreisl TN, Kim L, Moore K, *et al*: Phase II trial of single-agent bevacizumab followed by bevacizumab plus irinotecan at tumor progression in recurrent glioblastoma. *J Clin Oncol* 27: 740-745, 2009.
21. Plate KH, Breier G, Weich HA, Mennel HD and Risau W: Vascular endothelial growth factor and glioma angiogenesis: coordinate induction of VEGF receptors, distribution of VEGF protein and possible *in vivo* regulatory mechanisms. *Int J Cancer* 59: 520-529, 1994.
22. Bodey B, Siegel SE and Kaiser HE: Up-regulation of VEGF expression and related neo-angiogenesis in childhood high-grade gliomas: implications for anti-angiogenic anti-neoplastic therapy. *In Vivo* 20: 511-518, 2006.
23. Zagzag D, Lukyanov Y, Lan L, *et al*: Hypoxia-inducible factor 1 and VEGF upregulate CXCR4 in glioblastoma: implications for angiogenesis and glioma cell invasion. *Lab Invest* 86: 1221-1232, 2006.
24. Li D, Li XP, Wang HX, *et al*: VEGF induces angiogenesis in a zebrafish embryo glioma model established by transplantation of human glioma cells. *Oncol Rep* 28: 937-942, 2012.
25. Carmeliet P, Dor Y, Herbert JM, *et al*: Role of HIF-1alpha in hypoxia-mediated apoptosis, cell proliferation and tumour angiogenesis. *Nature* 394: 485-490, 1998.
26. Conway EM, Collen D and Carmeliet P: Molecular mechanisms of blood vessel growth. *Cardiovasc Res* 49: 507-521, 2001.
27. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25: 402-408, 2001.
28. Weidner N, Semple JP, Welch WR and Folkman J: Tumor angiogenesis and metastasis - correlation in invasive breast carcinoma. *N Engl J Med* 324: 1-8, 1991.
29. Zhang JP, Chen XY and Li JS: Effects of all-trans-retinoic acid on human gastric cancer cells BGC-823. *J Dig Dis* 8: 29-34, 2007.
30. Lu TY, Li WC, Chen RY, *et al*: Inhibition effects of all trans-retinoic acid on the growth and angiogenesis of esophageal squamous cell carcinoma in nude mice. *Chin Med J (Engl)* 124: 2708-2714, 2011.
31. Tee MK, Vigne JL and Taylor RN: All-trans retinoic acid inhibits vascular endothelial growth factor expression in a cell model of neutrophil activation. *Endocrinology* 147: 1264-1270, 2006.
32. Saito A, Sugawara A, Uruno A, *et al*: All-trans retinoic acid induces *in vitro* angiogenesis via retinoic acid receptor: possible involvement of paracrine effects of endogenous vascular endothelial growth factor signaling. *Endocrinology* 148: 1412-1423, 2007.
33. Chen JT, Liang JB, Chou CL, Shyu RC and Lu DW: Retinoic acid induces VEGF gene expression in human retinal pigment epithelial cells (ARPE-19). *J Ocul Pharmacol Ther* 21: 413-419, 2005.
34. Akiyama H, Tanaka T, Maeno T, *et al*: Induction of VEGF gene expression by retinoic acid through Sp1-binding sites in retinoblastoma Y79 cells. *Invest Ophthalmol Vis Sci* 43: 1367-1374, 2002.
35. Wang C, Chen FY, Gu CH, *et al*: Effect of ATRA and DNR on the expression and secretion of VEGF in leukemic cells. *Zhonghua Xue Ye Xue Za Zhi* 25: 171-174, 2004 (In Chinese).
36. Ye J, Liu FQ and Wu YP: The effect of all-trans retinoid acid and sodium selenite (Na<sub>2</sub>SeO<sub>3</sub>) on VEGF and its receptor expression in HL-60 cells. *Zhongguo Shi Yan Xue Ye Xue Za Zhi* 12: 142-146, 2004 (In Chinese).
37. Bai X, Fu JX, Ding KY, Cen JN, Wang W and Ruan CG: Quantitative analysis of gene expression for vascular endothelial growth factor and its application. *Zhongguo Shi Yan Xue Ye Xue Za Zhi* 13: 548-552, 2005 (In Chinese).
38. Cho DH, Choi YJ, Jo SA, Nam JH, Jung SC and Jo I: Retinoic acid decreases nitric oxide production in endothelial cells: a role of phosphorylation of endothelial nitric oxide synthase at Ser(1179). *Biochem Biophys Res Commun* 326: 703-710, 2005.
39. Fernández-Martínez AB, Jiménez MI, Hernández IS, *et al*: Mutual regulation of hypoxic and retinoic acid related signalling in tubular proximal cells. *Int J Biochem Cell Biol* 43: 1198-1207, 2011.
40. Fernández-Martínez AB, Arenas Jiménez MI and Lucio Cazaña FJ: Retinoic acid increases hypoxia-inducible factor- $\alpha$  through intracrine prostaglandin E(2) signaling in human renal proximal tubular cells HK-2. *Biochim Biophys Acta* 1821: 672-683, 2012.
41. Ardyanto TD, Osaki M, Tokuyasu N, Nagahama Y and Ito H: CoCl<sub>2</sub>-induced HIF-1  $\alpha$  expression correlates with proliferation and apoptosis in MKN-1 cells: A possible role for the PI3K/Akt pathway. *Int J Oncol* 29: 549-555, 2006.
42. Chattopadhyay N, Butters RR and Brown EM: Agonists of the retinoic acid- and retinoid X-receptors inhibit hepatocyte growth factor secretion and expression in U87 human astrocytoma cells. *Brain Res Mol Brain Res* 87: 100-108, 2001.