GABA_A receptor overexpression in the lateral hypothalamic area attenuates gastric ischemia-reperfusion injury in rats

LIN GAO^{1*}, TAO ZHU^{2,3*}, GUILIN XIE^{4*}, XIANGXIN LOU⁵, SHIBAO LI⁶, YAN ZHOU⁶, ZHENXU DENG³, DECHANG CHU³, JIYU LOU¹ and DONGSHU DU^{2,7}

¹Department of Neurology, The Affiliated Second Hospital of Zhengzhou University, Zhengzhou, Henan 450014;
²Laboratory of Neuropharmacology and Neurotoxicology, Shanghai University, Shanghai 200444; ³Department of Life Science, Heze University, Heze, Shandong 274500; ⁴Life Science College of Northeast Agricultural University, Harbin, Heilongjiang 150030; ⁵Department of Bioengineering, College of Chemistry, Chemical Engineering and Biotechnology, Donghua University, Shanghai 201620; ⁶Department of Laboratory Medicine, Lianyungang Hospital Affiliated Bengbu Medical College, Lianyungang, Jiangsu 222006;
⁷Shanghai Key Laboratory of Bio-Energy Crops, Shanghai University,

Shanghai 200444, P.R. China

Received January 27, 2014; Accepted August 22, 2014

DOI: 10.3892/mmr.2014.2816

Abstract. Excessive activation of the greater splanchnic nerve (GSN) has previously been determined to contribute to the progression of gastric ischemia-reperfusion (GI-R) injury. The present study was designed to estimate the protective effects of GABA_A receptor (GABA_AR) overexpression in the lateral hypothalamic area (LHA) against GI-R injury. The GI-R injury model was induced in rats by clamping the celiac artery for 30 min and then reperfusing for 1 h. Microinjection of recombinant adenoviral vectors overexpressing GABA_AR (Ad-GABA_AR) or control adenoviral vectors (Ad-Con) into the LHA was conducted in GI-R and normal control rats. Significant protective effects were observed on day 2 after Ad-GABA_AR treatment in the GI-R injury rats. Ad-GABA_AR treatment reduced plasma norepinephrine levels, plasma angiotensin II levels and peripheral GSN activity, but increased the gastric mucosal blood flow, as compared with Ad-Con treatment. These results indicate that adenoviral vector-induced GABA_AR overexpression in

Correspondence to: Dr Dongshu Du, Laboratory of Neuropharmacology and Neurotoxicology, Shanghai University, 99 Shangda Road, Shanghai 200444, P.R. China E-mail: sdhzdds@163.com

Professor Jiyu Lou, Department of Neurology, The Affiliated Second Hospital of Zhengzhou University, 2 Jingba Road, Zhengzhou, Henan 450014, P.R. China E-mail: loujiyu@yahoo.com.cn

*Contributed equally

Key words: lateral hypothalamic area, gastric ischemia reperfusion injury, GABA_A receptor

the LHA blunts GSN activity and subsequently alleviates the effects of gastric injury in GI-R rats.

Introduction

Major surgery-evoked ischemia has been demonstrated to induce gastric mucosal injury and gastrointestinal dysmotility (1,2). Previous studies (3-5) have shown that the hypothalamic paraventricular nucleus and the lateral hypothalamic area (LHA) are two specific hypothalamic nuclei that modulate gastric activity and gastric mucosal injury. The microinjection of GABAA receptor blocker in the LHA enhances GI-R injury. However, little is known regarding GABAA receptor expression and the protective effects of GABA_AR overexpression in the LHA against gastric ischemia-reperfusion (GI-R) injury in rats. As determined by a previous study (6), cerebellar-hypothalamic circuits regulate the gastric mucosal injury induced by ischemia-reperfusion. However, the detailed GABA_A receptor (GABA_AR)-mediated regulative mechanism in the LHA upon GI-R injury is not clear. In the present study, the effects of GABA AR overexpression induced by recombinant adenovirus vectors in the LHA following GI-R injury in rats were investigated. The aim of this study was to investigate the potential mechanisms of the GABA_A receptor in GI-R injury.

Materials and methods

Animals. Adult male Sprague Dawley (SD) rats were obtained from the Animal Resource Centre (Fudan University, Shanghai, China). All experimental procedures used in this study were performed in accordance with the Experimental Animal Care and Use Committee of Fudan University and conformed to the guidelines set out by the National Health and Medical Research Council of China. All rats were housed under controlled conditions (12 h light initiated at 20:00; 22-24°C) and provided access to water *ad libitum* for the duration of the study. The animals were fasted for 24 h prior to the experiment and were then allocated to the different groups. Following the experiments, the animals were deeply anesthetized with 10% chloral hydrate (solvent, 0.9% normal saline) and euthanized by cervical dislocation followed by decapitation.

Viral microinjection. Rats were anesthetized and placed in a stereotaxic frame (51600; Stoelting, Chicago, IL, USA). The recombinant adenoviral vectors overexpressing GABA_AR (Ad-GABA_AR; Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA) or control adenoviral vectors (Ad-Con) were bilaterally microinjected into the LHA (15 μ l for each side). The stereotaxic coordinates of LHA:LHA:AP 2.8 mm, LR: 1.5 mm, H: 8.3-8.5 mm, which was in accordance with Paxinos & Watson's rat atlas (http://www.callisto-science.org/NSI/Neuroscience_Image_Database/Rat_Brain_Atlas.html).

GI-R injury model. GI-R was performed following the microinjection of recombinant adenoviral vectors into the LHA according to previously reported methods (7). In brief, the abdominal cavity was cut open and the celiac artery was carefully isolated from the adjacent tissues. The celiac artery was clamped with a small vascular clip for 30 min and then reperfusion was established by removal of the clip for 1 h.

Control animals underwent an identical surgical procedure with the exception of clamping the celiac artery. At the end of the experiments, the rats were sacrificed as described. The stomachs were rapidly removed and were cut open along the greater curvature, then the gastric mucosa was carefully assessed for ulcers.

Assessment of gastric mucosal injury index (GMII). Gastric mucosal injury was measured as previously described (6). The stomach was incised along the greater curvature and washed with phosphate-buffered saline. The GMII was determined by a cumulative-length scale, in which an individual lesion limited by the mucosal epithelium, including the pinpoint erosions, ulcers and hemorrhagic spots, was scored according to length. The scores were calculated using the following criteria: 1, lesions ≤ 1 mm; 2, lesions ≥ 1 mm and ≤ 2 mm; 3, lesions ≥ 2 mm and ≤ 3 mm. For lesions of width ≥ 1 mm, the lesion score was doubled. The sum of the scores indicated the GMII. To prevent researcher bias, the GMII was measured by a researcher who was blind to the treatments.

Measurement of greater splanchnic nerve (GSN) activity. The rat was anesthetized with chloral hydrate (400 mg/kg intraperitoneally) and mounted onto a stereotaxic apparatus. The LHA coordinates were as described. Ad-GABA_AR (15 μ l for each side) was microinjected via a cannula connected to a microsyringe with a polyethylene tube. The injection lasted 2 min and the injection cannula was left for an additional 10 min to prevent backflow. The central end of the GSN was placed on thin bipolar platinum electrodes. The nerve-electrode junction was fixed and was electrically insulated from the surrounding tissues with a Vinyl Polysioxane Impression Material-auto-mixture (Sigma-Aldrich, St. Louis, MO, USA). The rectified output from the amplifier was monitored using the PowerLab[®] system (ADInstruments, Dunedin, New

Zealand) to record the raw nerve discharge. The basal nerve activity (baseline) was determined by analysis of the efferent GSN activity at the beginning of the experiment and background noise was measured by the nerve activity recorded at the end of the experiment. The nerve activity during the experiment was calculated by subtracting the background noise from the recorded value. The GSN activity response to the Ad-GABA_AR treatment was expressed as the percentage change from the basal value.

Gastric mucosal blood flow (GMBF) measurement. The GMBF was analyzed with a laser-Doppler flowmeter (LDF-2; Nankai University, Tianjin, China). In brief, the rats were anesthetized with chloral hydrate (400 mg/kg intraperitone-ally), the abdomen was opened, the stomach was exposed and transected, and the gastric contents were marginally evacuated to the exterior through the 5 mm incision in the stomach. Subsequently, the laser probe was placed 0.5 mm above and perpendicular to the mucosal surface to monitor GMBF, with measurements expressed in mV (value of Doppler signal voltage) on the digital panel of the flowmeter. When the GMBF was stable, four points for measurement were selected (one point for 1 min), and the average value was calculated and expressed as U/mV.

Western blot analysis of GABA_AR expression levels. Sample brain tissues were collected and homogenized in RIPA lysis buffer. The concentration of total protein was detected by bicinchoninic acid assay (Nanjing Jiancheng Bioengineering Institute, Nanjing, China). The proteins (40 μ g) were separated by 10% SDS-PAGE and transferred to nitrocellulose membranes (Pall Corporation, Pensacola, FL, USA). The membranes were probed with primary antibody to GABA_AR (1:200; Santa Cruz Biotechnology, Inc.) for 2 h. Next, the membranes were washed three times with phosphate-buffered saline and incubated with goat anti-rabbit IgG (1:1,000) secondary antibody for 2 h. Following incubation with enhanced chemiluminescence solution (Pierce Biotechnology, Inc., Rockford, IL, USA) and visualization by exposure to BioMax films (Kodak, Rochester, NY, USA), the membranes were stripped and probed with mouse monoclonal anti-β-actin primary antibody (1:300; Santa Cruz Biotechnology, Inc.) and rabbit anti-mouse secondary antibody for 2 h. The results are expressed as the optical density of the experimental band divided by that of the β -actin band of four replicate experiments. The optical density was measured by a gel-pro analyzer (Shanghai Furi Science & Technology Co., Ltd., Shanghai, China).

Measurement of plasma norepinephrine (NE). Blood samples were obtained from the carotid artery, through a tube that contained ETDA. The sample was centrifuged and mixed in an antioxidizing stabilizer sodium metabisulphite solution (5.2 mM). The plasma NE level was determined by high-performance liquid chromatography (HPLC) using a YWG-C18 column (250 mm x 4.6 mm x 5 μ m) and electrochemical detection (Waters 2465; Waters Corporation; Milford, MA, USA) as previously reported (8).

Measurement of plasma angiotensin (Ang) II. Plasma Ang II levels were analyzed using commercial ELISA kits (R&D



Figure 1. Effects of Ad-GABA_AR on GI-R injury in rats. P <0.05, as compared with the SD Ad-Con group. P <0.05, as compared with the GI-R Ad-Con group. n=8 in each group. Ad-GABA_AR, adenoviral vectors overexpressing GABA_A receptor; GI-R, gastric ischemia-reperfusion; SD, Spague Dawley; Ad-Con, control adenoviral vector.



Figure 2. $GABA_AR$ expression and cellular localization were detected by immunohistochemistry. As compared with the control group, $GABA_AR$ was upregulated in the Ad-GABA_AR groups. $GABA_AR$, $GABA_A$ receptor; Ad-GABAAR, adenoviral vectors overexpressing $GABA_AR$; Ad-Con, control group.

systems, Minneapolis, MN, USA) according to the manufacturer's instructions. Briefly, following incubation of 96-well plates with antibody specific for rat Ang II (1:500), the samples and standard diluent buffer were added to the wells, which were subsequently incubated and washed. Horseradish peroxidase-conjugated solution (1:100) was added and then washed out. The reactions were terminated with stop solution and the final solution was read at 450 nm using an ELISA plate reader (Shanghai Utrao Medical Instrument Co., Ltd., Shanghai, China).

HPLC. The amino acids in the microdialysis sample were separated by HPLC (LC-6A UQUID Chromatograph; Shimadzu Biotech Corporation, Kyoto, Japan) using a reverse-phase column (C18; Ultrasphere ODS; 4.6 mmx25 cm; 5 μ m particles), and quantified by o-phthaldialdehyde derivative and

fluorescence detection (RF-10AXL Shimadzu Fluorescence detector; 0.01 relative fluorescence units, 330 nm excitation wave-length and 450 nm emission wave-length). The mobile phase consisted of 0.1 m 63% potassium phosphate (pH 6.00, 6.25), 35% methanol and 2% tetrahydrofuran, and the flow rate was 1 ml/min. The experiments were conducted at 19-23°C (9).

Histology. Following the experiments, the rats were euthanized by an overdose injection of urethane followed by thoracotomy. The brain was removed from the skull, fixed in 10% formalin for four days or used to produce 40 mm coronal frozen sections and stained with GABA_AR antibody for immunohistochemical analysis, as previously reported (6).

Statistical analysis. A student's t-test and one-way analysis of variance was used for data analysis, and the data are presented as the means \pm standard error of the mean. GraphPad 5 software (GraphPad Software, Inc., La Jolla, CA, USA) was used and a P<0.05 was considered to indicate a statistically significant difference.

Results

*Effects of Ad-GABA*_A*R on GI-R injury.* It was first determined whether the overexpression of GABA_A*R* in the LHA exerted a protective effect upon GI-R injury. Microinjection of Ad-GABA_A*R* into the LHA was found to attenuate GI-R injury. The GMII that followed Ad-GABA_A*R* microinjection into the LHA, was significantly reduced as compared with the GMII subsequent to Ad-Con injection (122.7±15.6 versus 57.0±6.67; P<0.05; Fig. 1).

*Effects of Ad-GABA*_A*R microinjection into LHAwon GABA*_A*R expression*. In order to detect the depressive effects of Ad-GABA_A*R* on the expression of GABA_A*R* in the LHA, GABA_A*R* expression and cellular localisation were detected by immunohistochemistry. The results revealed that GABA_A*R* expression was upregulated by microinjection of Ad-GABA_A*R* in the normal SD group and the GI-R group (Fig. 2). The recombinant adenoviral vectors encoding GABA_A*R* in the LHA at two days after the viral microinjection in GI-R and normal SD rats (P<0.05; Fig. 3).

Plasma Ang II and NE levels. Plasma Ang II was significantly increased in the GI-R injury rats, as compared with the SD control rats (P<0.05), but Ad-GABA_AR treatment significantly reduced the plasma Ang II levels in the GI-R injury rats, as compared with Ad-Con treatment (P<0.05; Fig. 4A). The plasma NE levels, an indication of sympathetic activity (10), were significantly increased in the GI-R rats, as compared with the normal SD rats (P<0.05), although this effect was significantly normalized by Ad-GABA_AR treatment (P<0.05; Fig. 4B).

Effects of Ad- $GABA_AR$ expression on amino acid release. To investigate whether GI-R expression may result in amino acid change in the LHA, Ad-GABA_AR was microinjected into the LHA in the GI-R and SD control rats, and amino acid release was assessed. The amino acids examined in



Figure 3. Expression levels of GABA_AR in the LHA. The data were normalized to β -actin expression levels. *P<0.05, as compared with the SD Ad-Con group. *P<0.05, as compared with the GI-R Ad-Con group (n=4 in each group). GABA_AR, GABA_A receptor; LHA, lateral hypothalamic area; SD, Sprague Dawley; Ad-Con, control; GI-R, gastrointestinal ischemia-reperfusion.



Figure 4. Plasma Ang II and norepinephrin (NE) measurement following Ad-GABA_AR microinjection; n=8 for plasma Ang II and NE measurement in each group. (A) Ad-GABAAR treatment significantly reduced the plasma Ang II levels. (B) Recombinant adenoviral vectors encoding GABAAR significantly increased the expression levels of GABAAR in the LHA *P<0.05, as compared with the control SD rats. #P<0.05, as compared with GI-R rats subjected to Ad-Con treatment. Values are mean ± standard error of the mean. Ang II, Angiotensin II; Ad-GABA_AR, adenoviral vectors overexpressing GABA_A receptor; SD, Sprague Dawley; GI-R, gastrointestinal ischemia-reperfusion; Ad-Con, control adenoviral vector.



Figure 5. Effects of microinjection of Ad-GABA_AR into the lateral hypothalamic area on the local release of amino acids. (A) Aspartate (Asp); (B) glutamate; (C) glycine (Gly); (D) taurine (Tau). *P<0.05, as compared with the control SD rats. *P<0.05, as compared with GI-R rats subjected to Ad-Con treatment. n=8 per group. Ad-GABA_AR, adenoviral vectors overexpressing GABA_A receptor; SD, Sprague Dawley; GI-R, gastrointestinal ischemia-reperfusion; Ad-Con, control adenoviral vector.



Figure 6. Effects of GABA_AR on GSN activity and gastric mucosal blood flow. *P<0.05, as compared with the control SD rats. *P<0.05, as compared with GI-R rats subjected to Ad-Con treatment. n=8 in each group. GABA_AR, GABA_A receptor; GSN, greater splanchnic nerve; SD, Sprague Dawley; GI-R, gastrointestinal ischemia-reperfusion; Ad-Con, control adenoviral vector.

the LHA included excitatory [glutamate (Glu) and aspartate (Asp)] as well as inhibitory [taurine (Tau) and glycine (Gly)] amino acids. The baseline release of excitatory amino acid neurotransmitters (Glu and Asp) was significantly increased but that of the inhibitory amino acids (Tau and Gly) was significantly reduced in the GI-R injury rats, as compared with the normal SD rats (all P<0.05). Microinjection of Ad-GABA_AR into the LHA significantly increased Tau and Gly release, and significantly reduced Glu and Asp release, as compared with Ad-Con microinjection (P<0.05; Fig. 5).

Effects of GABA_AR on GSN activity and GMBF. In order to determine whether the central GABA_AR mediated GSN activity, the GSN fire frequency was analyzed. The GSN mediates GMBF, which regulates gastric activity. As shown in Fig. 6, microinjection of Ad-GABA_AR into the LHA following GI-R, significantly reduced GSN activity (P<0.05) and increased GMBF (P<0.05), as compared with Ad-Con treatment.

Discussion

The present study provides evidence that $GABA_AR$ overexpression in the LHA results in a profound protection against GI-R injury in rats. In addition to reducing plasma Ang II and NE levels, the LHA-targeted adenovirus reduces the degree of gastric mucosal injury. These results suggest the importance of the GABA_AR in the LHA in the neural gastrointestinal control and support the hypothesis that GABA_AR in the LHA is predominantly involved in the pathophysiological process of GI-R injury (5,11). The present study also demonstrated that an adenovirus, targeting $GABA_AR$ in the LHA in GI-R rats, effectively inhibited the GSN activity that contributes to the elevated GMBF (12).

An important finding from the present study was that GABA_AR gene overexpression in the LHA reduced the plasma NE accompanied with improved GI-R injury. One limitation of the present study is that the complex association between over-enhanced GSN activity and the degree of gastric mucosal injury is difficult to clarify. Nevertheless, GSN overdrive has been well-established as a causative factor of GI-R injury, and a close association between GSN activity and GMBF has been identified. Central enhanced GABA signals have been shown to reduce plasma Ang II levels in exercise-training rats (13). In the present study, the plasma Ang II levels were increased in GI-R rats. The adenovirus-induced GABA_AR overexpression in the LHA normalized plasma Ang II levels in GI-R rats, which may be beneficial for the attenuation of GI-R injury. Furthermore, adenoviruses allow the efficient delivery of relatively large transgenes to the brain, and these viruses infect glial and neuronal cells (14,15). The viruses used in the present study were considered to be specific for $GABA_AR$ and not for $GABA_BR$ or $GABA_CR$.

The data from the present study support the hypothesis that the $GABA_AR$ signaling pathway mediates regulative effects on GI-R injury via an increase in excitatory and a reduction in inhibitory amino acid release. Excitatory amino acids (Glu and Asp) induce a GSN tension effect, whereas inhibitory amino acids (Gly and Tau) cause a GSN relaxed response (16,17). In the present study, the release of excitatory amino acids (Gly and Tau) were lower in the GI-R than in the normal SD control group. Therefore, the GABA_ARs in the LHA may regulate gastric activity via the modulation of amino acid release.

In conclusion, in the present study, a $GABA_AR$ signaling pathway in the LHA during the development of GI-R injury was investigated. Furthermore, the protective effects of $GABA_AR$ overexpression in the LHA against GI-R injury in rats were analyzed and an increased inhibitory and suppressed excitatory amino acid release was identified.

Acknowledgements

This study was sponsored by the National Natural Science Foundation of China (grant nos. 31100838 and 31172147).

References

- 1. Tsukamoto T, Antonic V, El Hajj II, *et al*: Novel model of peripheral tissue trauma-induced inflammation and gastrointestinal dysmotility. Neurogastroenterol Motil 23: 379-386, 2011.
- Lucas CÉ, Sugawa C, Riddle J, et al: Natural history and surgical dilemma of "stress" gastric bleeding. Arch Surg 102: 266-273, 1971.
- Zhang JF, Zhang YM, Yan CD and Zhou XP: Neuroregulative mechanism of hypothalamic paraventricular nucleus on gastric ischemia-reperfusion injury in rats. Life Sci 71: 1501-1510, 2002.
- Zhang YM, Wei EQ, Li L, et al: Extracellular signal-regulated kinase pathways may mediate the protective effect of electrical stimulation of the paraventricular nucleus against ischaemia-reperfusion injury of the gastric mucosa. Clin Exp Pharmacol Physiol 34: 742-752, 2007.
- 5. Zhu SP, Fei SJ, Zhang JF, *et al*: Lateral hypothalamic area mediated the protective effects of microinjection of glutamate into interpositus nucleus on gastric ischemia-reperfusion injury in rats. Neurosci Lett 1: 39-43, 2012.
- 6. Du DS, Zhu T, Ren ST, et al: γ-aminobutyric acid-mediated neurotransmission in cerebellar-hypothalamic circuit attenuates gastric mucosal injury induced by ischemia-reperfusion. Neurogastroenterol Motil 4: 313-e249, 2013.
- 7. Du DŠ, Ma XB, Zhang JF, *et al*: The protective effect of capsaicin receptor-mediated genistein postconditioning on gastric ischemia-reperfusion injury in rats. Dig Dis Sci 55: 3070-3077, 2010.
- Shi Z, Chen AD, Xu Y, *et al*: Long-term administration of tempol attenuates postinfarct ventricular dysfunction and sympathetic activity in rats. Pflugers Arch 458: 247-257, 2009.
- 9. Wang J, Peng YJ, Zhu DN, *et al*: Amino acids mediate the hypotensive effect of angiotensin-(1-7) at the caudal ventrolateral medulla in rats. Regul Pept 129: 1-7, 2005.
- Meng L, Ye XJ, Ding WH, et al: Plasma catecholamine release-inhibitory peptide catestatin in patients with essential hypertension. J Cardiovasc Med (Hagerstown) 12: 643-647, 2011.
- 11. Zhu JZ, Fei SJ, Zhang JF, *et al*: Lateral hypothalamic area mediated the aggravated effect of microinjection of Baclofen into cerebellar fastigial nucleus on stress gastric mucosal damage in rats. Neurosci Lett 509: 125-129, 2012.
- Gao L, Fei S, Qiao W, *et al*: Protective effect of chemical stimulation of cerebellar fastigial nucleus on stress gastric mucosal injury in rats. Life Sci 88: 871-878, 2011.
- Rossi NF, Chen H and Maliszewska-Scislo M: Paraventricular nucleus control of blood pressure in two-kidney, one-clip rats: effects of exercise training and resting blood pressure. Am J Physiol Regul Integr Comp Physiol 305: R1390-R1400, 2013.
- Davidson BL, Allen ED, Kozarsky KF, et al: A model system for in vivo gene transfer into the central nervous system using an adenoviral vector. Nat Genet 3: 219-223, 1993.
- 15. Toscano MG, Romero Z, Muñoz P, *et al*: Physiological and tissue-specific vectors for treatment of inherited diseases. Gene Ther 18: 117-127, 2011.
- 16. Koganezawa T, Shimomura Y and Terui N: The viscerosympathetic response in rabbits is mediated by GABAergic and glutamatergic inputs into the sympathetic premotor neurons of the rostral ventrolateral medulla. Exp Physiol 95: 1061-1070, 2010.
- 17. Kawabe T, Kawabe K and Sapru HN: Tonic γ -aminobutyric acid-ergic activity in the hypothalamic arcuate nucleus is attenuated in the spontaneously hypertensive rat. Hypertension 62: 281-287, 2013.