Antineoplastic effect of a novel chemopreventive agent, neokestose, on the Caco-2 cell line via inhibition of expression of nuclear factor-κB and cyclooxygenase-2

SHUN-MEI LEE¹, JAN-YI CHANG¹, JIANN-SHING WU^2 and DEY-CHYI SHEU¹

¹Department of Bioengineering, Tatung University, Taipei 104; ²Department of Styling and Cosmetology, Hsin Shen Junior College of Medical Care and Management, Taoyuan 325, Taiwan, R.O.C.

Received June 3, 2014; Accepted February 17, 2015

DOI: 10.3892/mmr.2015.3507

Abstract. Neokestose is a ⁶G-fructooligosaccharide (FOS) and an important prebiotic. When FOS are ingested by patients with colorectal cancer, they may come into contact with cancer cells prior to being fermented by bifidobacteria in the colon. In the present study, the effects of neokestose on cell proliferation, cell cycle and apoptosis of the colorectal cancer cell line Caco-2 were investigated to evaluate its anti-cancer effect. An MTT assay showed that neokestose-treated Caco-2 cells exhibited a significant and dose-dependent loss of viability. Flow cytometric analysis indicated that the sub-G1 population of Caco-2 cells was significantly increased following treatment with neokestose, and the percentage of Caco-2 cells in the stage of late apoptosis was also significantly increased in a dose-dependent manner. Western blot analysis showed that the overexpression of nuclear factor-kB, a central molecule responsible for the transition from inflammation to cancer, and cyclooxygenase-2, an important enzyme in colorectal tumorigenesis, in colorectal carcinoma cells was inhibited by neokestose. Accordingly, the present study provided in vitro evidence that neokestose may be used as a dietary chemopreventive agent, whose application is more rational than that of COX-2 inhibitors or aspirin for preventing colorectal cancer.

Correspondence to: Professor Dey-Chyi Sheu, Department of Bioengineering, Tatung University, 40 Section 3 Zhongshan North Road, Taipei 104, Taiwan, R.O.C. E-mail: dcsheu@ttu.edu.tw

Professor Jiann-Shing Wu, Department of Styling and Cosmetology, Hsin Shen Junior College of Medical Care and Management, 418 Section Gaoping, Zhongfeng Road, Lontang District, Taoyuan 325, Taiwan, R.O.C.

E-mail: jswu@hsc.edu.tw

Key words: antineoplastic effect, Caco-2, cyclooxygenase-2, neokestose, nuclear factor-кB

Introduction

Sporadic colorectal cancer accounts for 75% of total cases of colorectal cancer, and is the third most common cancer in the world as well as the second leading cause of cancer mortality (1). Screening tests are able to reduce the mortality rate of colorectal cancer; however, a large proportion of individuals do not undergo colorectal cancer screening due to its inconvenience and high cost (2-5). The increasing frequency of colorectal cancer worldwide makes the prevention of this disease an urgent problem to be solved.

Previous studies suggested that fructooligosaccharides (FOS) can inhibit the process of colon cancer development, mainly by its prebiotic effect and the short-chain fatty acids (SCFA) produced during the fermentation of FOS in the colon (6-9). FOS are important prebiotics, since FOS cannot be digested in the small intestine of humans, but can be fermented in the colon to form short-chain fatty acids (SCFA) and lactic acid (10). FOS have recently attracted attention due to their beneficial physiological effects. These include relief of constipation, improved mineral absorption, as well as reduced blood levels of total cholesterol, triglycerides and phospholipids (8,11,12).

FOS are classified into three categories: ¹F-, ⁶F- and ⁶G-FOS (13-15). Among these FOS, only ¹F-FOS is commercially available. However, ⁶G-FOS, also called neo-FOS, has fructosyl units bound at the β (2 \rightarrow 6) position in the glucose moiety of sucrose. Hence, neo-FOS has superior bifidogenecity as well as better heat and chemical stability than ¹F-FOS (16,17). Neo-FOS comprises neokestose and neo-nystose (18); the structure of neokestose, together with the structures of 1-kestose (a ¹F-FOS) and 6-kestose (a ⁶F-FOS), is shown in Fig. 1.

When FOS are ingested by patients with colorectal cancer, they may come into contact with cancer cells prior to being fermented by bifidobacteria in the colon. Among various FOS, neokestose was selected as the candidate in the present study, as it is the prebiotic with the highest potency, as mentioned above. The effects of neokestose on the colon cancer cell line Caco-2 were investigated, and the potential applications of neokestose for the dietary chemoprevention of colorectal cancer were also evaluated. The present study aimed to determine the biological anti-cancer activity of neokestose towards the colon cancer cell line Caco-2, since neokestose is the most potential prebiotic among the fructooligosaccharides. For this, neokestose was produced using yeast fermentation of sucrose, isolated using high-performance liquid chromatography (HPLC), and its anti-cancerous effect against Caco-2 cells was assessed using MTT and flow cytometric assays as well as western blot analysis. The results indicated that neokestose may be an effective dietary chemopreventive against colorectal cancer.

Materials and methods

Preparation of neokestose. As previously reported, neo-FOS was produced by the culture of *Xanthophyllomyces dendrorhous* (BCRC 22367; Bioresource Collection and Research Center, Hsinchu, Taiwan), using 250 g/l sucrose as the substrate (15). Neokestose was purified by HPLC on a semipreparative YMC-pack ODS-AQ column (20x250 mm; YMC Co., Ltd., Kyoto, Japan) with a Waters 410 differential refractive index detector (Waters Corporation, Milford, MA, USA). Water at a flow rate of 8 ml/min was used as the mobile phase. A 200- μ l sample of neo-FOS (containing 109 g/l neokestose) was injected for the purification of neokestose. The respective peak fractions were collected and concentrated by rotary evaporation.

Cell culture. The Caco-2 cell line (BCRC 60182) was obtained from the Bioresource Collection and Research Center. Cells were routinely maintained and subcultured in 10 cm²-dishes at 37°C in a humidified CO_2 incubator (95% air and 5% CO_2). The medium for cell maintenance consisted of 10% heat-inactivated fetal bovine serum and 100 U/ml penicillin and 100 g/ml streptomycin in Dulbecco's modified Eagle's medium (DMEM) solution, which are all from HyClone Laboratories, Inc. (Logan, UT, USA). When cells were 80% confluent, they were subcultured using 0.25% trypsin and 0.02% EDTA in D-Hanks' balanced salt solution (HyClone Laboratories, Inc.). The medium was replaced every 48 h.

MTT assay. MTT (Sigma, St Louis, MO) is a tetrazolium salt that can be cleaved by mitochondrial dehydrogenase in living cells. The effects of neokestose (0, 0.5, 1.0 or 2.5 mg/ml) on cell growth were tested using the MTT assay. Caco-2 cells were cultured in 96-well plates at a density of 1.0×10^4 cells/well for two days, then the medium was discarded, $20 \ \mu$ l MTT solution [0.5 mg/ml in phosphate-buffered saline (PBS)] was added to all wells, and the cells were incubated for 2 h. At the end of the incubation, $100 \ \mu$ l dimethylsulfoxide was added to a microplate reader (SynergyTM HT; BioTek, Winooski, VT, USA). Absorbance was recorded at 570 nm. All experiments were performed at least four times with four wells of each concentration.

Cell cycle analysis. To analyze the cell cycle distribution, cells were washed twice with PBS, collected by centrifugation at 725 x g for 5 min at 4°C, and fixed in 70% (v/v) ethanol at 4°C for 30 min. After fixation, cells were resuspended in PBS and stained with propidium iodide (PI; Sigma-Aldrich, St. Louis, MO, USA) solution (containing 48 μ g/ml propidium





Figure 1. Chemical structures of 1-kestose, 6-kestose and neokestose.

iodide and 48 μ g/ml RNase A; Invitrogen Life Technologies, Carlsbad, CA, USA) for 20 min. The DNA content of the cells was examined by flow cytometry (FACSCalibur; BD Biosciences, Franklin Lakes, NJ).

Apoptosis analysis. Apoptosis was measured by using a fluorsecein isothiocyanate (FITC) Annexin V Apoptosis Detection Kit I (BD Biosciences) according to the manufacturer's instructions. Briefly, cells were washed twice with PBS and collected by centrifugation at 725 x g for 5 min at 25°C. Cells were resuspended in 100 μ l of binding buffer and labeled with 5 μ l FITC Annexin V and 5 μ l PI solution for 15 min in the dark. After labeling, cells were resuspended in 400 μ l binding buffer and detected by flow cytometry (FACSCalibur).

Western blot analysis. Caco-2 cells were lysed with Cell Extraction Buffer (M-PER® Mammalian Protein Extraction Reagent, Thermo Fisher Scientific, Waltham, MA, USA). The bicinchoninic acid protein assay kit (cat. no. 23225; Thermo Fisher Scientific) was used to measure the protein concentration. The cell extracts were subjected to 10% SDS-PAGE. Following electrophoresis, the proteins were transferred onto a nitrocellulose membrane (Bio-Rad Laboratories, Inc., Hercules, CA, USA). After blocking with 3% bovine serum albumin (Sigma-Aldrich) in Tris-buffered saline (20 mM Tris, pH 7.5, 150 mM NaCl) containing 0.1% Tween 20 (T-TBS) for 1 h, nitrocellulose membranes were incubated with the following primary antibodies (all diluted 1:1,000): Rabbit monoclonal anti-COX-2 (cat. no. 12282; Cell Signaling Technology, Danvers, MA, USA) mouse monoclonal anti-NF-kB (cat. no. MA5-16160; Thermo Fisher Scientific) and mouse monoclonal anti-actin (cat. no. MA1-744; Thermo Fisher Scientific), for 8 h at 4°C and then washed in T-TBS.

The secondary antibody, horseradish peroxidase-conjugated rabbit anti-mouse IgG(H+L) (cat. no. 315-035-003; Jackson ImmunoResearch Laboratories, Inc., West Grove, MA, USA) or mouse anti-rabbit IgG(H) (cat. no. GTX628140-01; GeneTex, Inc., Irvine, CA, USA), was incubated with the membranes at a dilution of 1:2,000 in T-TBS. After washing, the antibody complexes were detected by chemiluminescence using ECL reagents (Clarity[™] Western ECL Substrate; Bio-Rad Laboratories, Inc.) and ImageQuant LAS 4000 (GE Healthcare Bio-Sciences, Pittsburgh, PA, USA).

Statistical analysis. The results of multiple experiments are expressed as the mean \pm standard error. Differences between the treatment and control groups were analyzed using Student's *t*-test. A P<0.05 was considered to indicate a statistically significant difference, which was denoted by asterisks in the figures.

Results

Preparation of neokestose. Neokestose was isolated from neo-FOS, which is produced from sucrose by the culture of *Xanthophyllomyces dendrorhous* BCRC 22367, which has ⁶G-fructofuranosidase (⁶G-FFase) activity (15). Fig. 2 shows the HPLC chromatogram of the neo-FOS mixture. Peak separation for the neo-FOS components was achieved. In Fig. 2, peak 6 indicates the neokestose fraction, which was the major component of the neo-FOS mixture. Approximately 20 mg pure neokestose was obtained from every HPLC run. The neo-FOS product consisted of 77.0% neokestose, 10.4% neo-nystose, 2.8% 1-kestose, 4.5% monosaccharides (glucose and fructose) and 5.3% sucrose on a dry weight basis. This product is adequate for large-scale preparation of neokestose.

Effect of neokestose on cell viability. The antiproliferative effect of neokestose on Caco-2 cells was measured by an MTT assay. Caco-2 cells were treated with different concentrations (0-2.5 mg/ml) of neokestose for 48 h, and the obtained results are shown in Fig. 3. In Fig. 3, the cell viability of the Caco-2 cells was decreased from 100% to ~59% as the concentration of neokestose was increased from 0 to 2.5 mg/ml. The results showed that Caco-2 cells treated with neokestose exhibited a significant and dose-dependent loss of viability.

Effect of neokestose on the sub-G1 population and tumor cell apoptosis. Caco-2 cells were treated with neokestose for 48 h, and the cell cycle distribution and apoptosis were measured by flow cytometric analysis. Neokestose treatment significantly increased the sub-G1 phase population (Fig. 4). Similarly, neokestose treatment also dramatically increased the percentage of the late apoptotic cells, according to the PI-Annexin V staining assay (Fig. 5).

Fig. 4 clearly indicates that the sub-G1 phase population increased from 8, 22, 28 and 45% when the Caco-2 cells were treated by an increasing concentrations of neokestose from 0 (control), 0.5, 1.0 and 2.5 mg/ml. In the flow cytometric cell cycle distribution curve, the sub-G1 area represents a population of cells with a reduced DNA content, which is a marker for apoptosis. Therefore, the significantly increased sub-G1 phase population indicates increasing apoptosis in the Caco-2 cells.



Figure 2. High-performance liquid chromatogram of neo-fructooligosaccharides on a semipreparative ODS-AQ column (Peaks: S, salts; 1, fructose and glucose; 2, glycerol; 3, sucrose; 4, ethanol; 5, 1-kestose; 6, neokestose; 7, neonystose).



Figure 3. Anti-proliferative effect of neokestose on Caco-2 cells. Caco-2 cells were treated with 0.5, 1.0 or 2.5 mg/ml of neokestose for 48 h, and Caco-2 cells without neokestose treatment served as a control. The cell viability of Caco-2 cells was then determined using the MTT assay. Values are expressed as the mean \pm standard error of four independent experiments (*P<0.05 vs. control).

In addition, as mentioned above, apoptosis was further verified by annexin V-FITC/PI double staining using a flow cytometry. Therefore, neokestose-treated Caco-2 cells were stained with PI and annexin V-FITC to determine the apoptotic rate of the cells. Cells were distiguished as viable (annexin V-FITC, PI double negative), early apoptotic (annexin V-FITC positive, PI negative) and late apoptotic (annexin V-FITC, PI double positive) cells. In Fig. 6, it can be clearly seen that when the Caco-2 cells were treated with neokestose at concentrations of 0 (control), 0.5, 1.0 and 2.5 mg/ml, the percentage of early apoptotic cells was 8.3, 10.9, 9.0 and 8.7%, respectively, along with percentages of late apoptotic cells of 15.3, 25.3, 27.2 and 31.2%, respectively. This indicated that neokestose efficiently induced apoptosis of Caco-2 cells.

From the results, it was evident that neokestose affected cell cycle progression and apoptosis in colorectal cancer cells.



Figure 4. Effect of neokestose on cell cycle distribution of Caco-2 cells. Cells were treated with 0.5, 1.0 or 2.5 mg/ml neokestose, and untreated cells served as a control. (A) Caco-2 cells were stained with PI and their DNA content was analyzed by flow cytometric analysis. (B) Quantified values obtained by integration of the cell cycle distribution curves are expressed as the mean ± standard error of four independent experiments. PI, propidium iodide.



Annexin V-FITC

Figure 5. Effect of neokestose on cell apoptosis of Caco-2 cells. Caco-2 cells were treated with 0.5, 1.0 and 2.5 mg/ml neokestose for 48 h, and Caco-2 cells without neokestose treatment served as a control. Apoptosis of Caco-2 cells was then determined by a PI and annexin V-FITC double-staining assay and quantified by flow cytometric analysis. Annexin V-FITC '/PI', annexin V-FITC '/PI' and annexin V-FITC '/PI' cells were considered to be viable, early-apoptotic, and late-apoptotic cells, respectively. PI, propidium iodide; FITC, fluorescein isothiocyanate.



Figure 6. Effect of neokestose on the protein levels of (A) NF- κ B and (B) COX-2 in Caco-2 cells. Caco-2 cells were treated with neokestose at 0.5, 1.0 or 2.5 mg/ml for 48 h, and Caco-2 cells without neokestose treatment served as a control. NF- κ B and COX-2 proteins were extracted and analyzed by western blotting. Protein expressions were normalized to the β -actin control. NF- κ B, nuclear factor kappa B; COX, cyclooxygenase.

Effect of neokestose on expression of NF-кВ and COX-2. To evaluate potential molecular mechanisms causing apoptosis, the expression of NF-KB and COX-2 was assessed by western blot analysis (19,20). Caco-2 cells were treated with neokestose at concentrations of 0 (control), 0.5, 1.0 or 2.5 mg/ml for 48 h. Protein was then extracted and analyzed by western blotting. Fig. 6 shows that NF-KB and COX-2 were overexpressed in the Caco-2 cell line; however, following treatment with neokestose, the expression of NF-kB and COX-2 protein was significantly reduced in a dose-dependent manner.

Discussion

In the present study, neokestose produced by the conversion of sucrose by Xanthophyllomyces dendrorhous BCRC 22367 induced apoptosis in the colorectal cancer cell line Caco-2. Flow cytometric analysis showed that neokestose increased the population of Caco-2 cells in sub-G1 phase, and increase the percentage of annexin V-FITC-positive cells. These results indicated that neokestose induced apoptosis of Caco-2 cells in a dose-dependent manner, which is consistent with the results of the cell viability test.

NF-kB and COX-2 are commonly used markers of cell apoptosis. Therefore, the protein expression of NF-kB and COX-2 in Caco-2 cells treated with various concentrations of neokestose was evaluated. The results showed that the expression of NF-κB and COX-2 was decreased in Caco-2 cells treated with neokestose. NF-kB is a central molecule responsible for the transition from inflammation to cancer (19), and COX-2 has an important role in colorectal tumorigenesis (4,19). Accordingly, the results of the present study demonstrated that neokestose has a potent bifidogenic effect directed against colorectal cancer.

In conclusion, the results of the present study demonstrated that neokestose significantly induces apoptotic cell death in a dose-dependent manner via inhibition of NF-KB and COX-2 expression in colorectal carcinoma cells. Accordingly, neokestose may be used as a chemopreventive agent for colorectal cancer. Ideally, an anti-cancer agent or chemopreventive should inhibit the growth of cancer cells without affecting normal cell growth. To the best of our knowledge, the present study was the first to provide in vitro evidence that neokestose may serve as a potential dietary chemopreventive agent for colorectal cancer.

Acknowledgements

The authors are grateful for the financial supports provided by Tatung University and Taipei Medical University.

References

- 1. Pisani P, Bray F and Parkin DM: Estimates of the world-wide prevalence of cancer for 25 sites in the adult population. Int J Cancer 97: 72-81, 2002
- 2. Arber N and Levin B: Chemoprevention of colorectal cancer: ready for routine use? Curr Top Med Chem 5: 517-525, 2005.
- 3. Dvory-Sobol H and Arber N: Cyclooxygenase-2 as target for chemopreventive interventions: new approaches. Cancer Biomark 3: 153-161, 2007.
- 4. Manzano A and Pérez-Segura P: Colorectal cancer chemoprevention: is this the future of colorectal cancer prevention? Scientific World Journal 2012: 327-341, 2012
- 5. Temraz S, Mukherji D and Shamseddine A: Potential targets for colorectal cancer prevention. Int J Mol Sci 14: 17279-17303, 2013.
- 6. Pool-Zobel B, van Loo J, Rowland I and Roberfroid MB: Experimental evidences on the potential of prebiotic fructans to reduce the risk of colon cancer. Brit J Nutr 87 (Suppl 2): 273-281, 2002
- 7. Swennen K, Courtin CM and Delcour JA: Non-digestible oligosaccharides with prebiotic properties. Crit Rev Food Sci Nutr 46: 459-471, 2006.
- 8. Sabater-Molina M, Larque E, Torrella F and Zamora S: Dietary fructooligosaccharides and potential benefits on health. J Physiol Biochem 65: 315-328, 2009. 9. Vargas AJ and Thompson PA: Diet and nutrient factors in
- colorectal cancer risk. Nutr Clin Pract 27: 613-623, 2012.
- 10. Oku T, Tokunaga T and Hosoya N: Nondigestibility of a new sweetener, 'Neosugar' in the rat. J Nutr 114: 1574-1581, 1984.
- 11. Hidaka H, Eida T, Takizawa T, Tokunaga T and Tashiro Y: Effects of fructooligosaccharides on intestinal flora and human health. Bifidobacteria Microflora 5: 37-50, 1986.
- 12. Ooi LG and Liong MT: Cholesterol-lowering effects of probiotics and prebiotics: a review of in vivo and in vitro findings. Int J Mol Sci 11: 2499-2522, 2010.
- 13. Maiorano AE, Piccoli RM, da Silva ES and de Andrade Rodrigues MF: Microbial production of fructosyltransferases for synthesis of pre-biotics. Biotechnol Lett 30: 1867-1877, 2008.
- 14. de Abreu M, Alvaro-Benito M, Sanz-Aparicio J, Plou FJ, Fernandez-Lobato M and Alcalde M: Synthesis of 6-kestose using an efficient β -fructofuranosidase engineered by directed evolution. Adv Synth Catal 355: 1698-1702, 2013. 15. Sheu DC, Chang JY, Chen YJ and Lee CW: Production of high-purity
- neofructooligosaccharides by culture of Xanthophyllomyces dendrorhous. Bioresour Technol 132: 432-435, 2013.
- 16. Kilian S, Kritzinger S, Rycroft C, Gibson G and du Preez J: The effects of the novel bifidogenic trisaccharide, neokestose, on the human colonic microbiota. World J Microbiol Biotechnol 18: 637-644, 2002
- 17. Lim JS, Lee JH, Kang SW, Park SW and Kim SW: Studies on production and physical properties of neo-FOS produced by co-immobilized Penicillium citrinum and neo-fructosyltransferase. Eur Food Res Technol 225: 457-462, 2007.
- 18. Linde D, Rodríguez-Colinas B, Estévez M, Poveda A, Plou FJ and Fernández-Lobato M: Analysis of neofructooligosaccharides production mediated by the extracellular β -fructofuranosidase from Xanthophyllomyces dendrorhous. Biores Technol 109: 123-130, 2012
- 19. Karin M, Cao Y, Greten FR and Li ZW: NF-kappaB in cancer: from innocent bystander to major culprit. Nat Rev Cancer 2: 301-310, 2002.
- 20. Oshima M, Dinchuk JE, Kargman SL, Oshima H, Hancock B, Kwong E, Trzaskos JM, Evans JF and Taketo MM: Suppression of intestinal polyposis in Apc delta716 knockout mice by inhibition of cyclooxygenase 2 (COX-2). Cell 87: 803-809, 1996.