

1,25(OH)₂D₃ inhibits the progression of hepatocellular carcinoma via downregulating HDAC2 and upregulating P21(WAFI/CIP1)

JIAN HUANG¹, GUOZHEN YANG², YUNZHU HUANG³, WEIYING KONG¹ and SHU ZHANG²

¹Biochemistry Department, Affiliated Hospital of Guiyang Medical College; ²Medical Laboratory, Guizhou Medical University; ³Biochemistry Department, Affiliated Hospital of Guizhou Medical University, Guiyang, Guizhou 550004, P.R. China

Received January 19, 2015; Accepted October 28, 2015

DOI: 10.3892/mmr.2015.4676

Abstract. Vitamin D, termed 1,25(OH)₂D₃ in it's active form, activity is associated with a reduced risk of hepatocellular carcinoma (HCC) and is an important immune regulator. However, the detail molecular mechanisms underlying the effects of 1,25(OH)₂D₃ on the progression of HCC are widely unknown. Histone deacetwylase 2 (HDAC2) is usually expressed at high levels in tumors, and its downregulation leads to high expression levels of cell cycle components, including p21(WAF1/Cip1), a well-characterized modulator, which is critical in cell senescence and apoptosis. The present study investigated whether vitamin D inhibits HCC via the regulation of HDAC2 and p21(WAF1/Cip1). Firstly, the toxic concentrations of 1,25(OH)₂D₃ were determined, according to trypan blue and [³H]thymidine incorporation assays. Secondly, HCC cells lines were treated with different concentrations of 1,25(OH)₂D₃. The expression of HDAC2 was either silenced via short hairpin (sh)RNA or induced by transfection of plasmids expressing the HDAC2 gene in certain HCC cells. Finally the mRNA and protein levels of HDAC2 and p21(WAF1/Cip1) were measured using reverse transcription-quantitative polymerase chain reaction and western blot analyses. The results revealed that $1,25(OH)_2D_3$ treatment reduced the expression of HDAC2 and increased the expression of p21(WAF1/Cip1), in a dose-dependent manner, resulting in the reduction of HCC growth. Elevated levels of HDAC2 reduced the expression of p21(WAF1/Cip1), resulting in an increase in HCC growth. HDAC2 shRNA increased the expression of p21(WAF1/Cip1), resulting in reduction in HCC growth. Thus, 1,25(OH)₂D₃ exerted antitumorigenic effects through decreasing the expression levels of HDAC2 and increasing the expression of p21(WAF1/Cip1), which inhibited the development of HCC and may indicate the possible underlying mechanism. These results suggest that vitamin D_3 may be developed as a potential drug for effective therapy in the treatment of HCC.

Introduction

Hepatocellular carcinoma (HCC) accounts for 90% of all primary liver malignancies and is resistant to chemotherapy (1,2). The survival rates of patients with HCC is usually <1 year following diagnosis (3). Only 10-20% of patients with HCC are suitable for resective surgery, and the five-year relapse-free survival rate is no more than 20-30% (4). Developments in the techniques used to treat HCC have improved the survival rates of patients with HCC (5), however, this remains far from satisfactory. Therefore, novel effective therapeutic strategies are in high demand.

Epidemiological and clinical studies have shown that the downregulation of vitamin D is associated with an increased risk of various types of cancer; whereas a high level of $1,25(OH)_2D_3$ decreases the risk of cancer (6). Other investigations have also demonstrated that 1,25(OH)₂D₃ can modulate calcium and skeletal homeostasis, and exert a marked effect on the growth and differentiation of various tissues (7). Further evidence indicates that 1,25(OH)₂D₃ modulates the bioactivity of various immune cells (8). Of note, protein modification, and the acetylation of histones and non-histone proteins mediates gene expression levels and cell signaling. The removal of acetyl groups from acetylated histones by histone deacetylase 2 (HDAC2) reverses the bioactivity of histone acetyltransferases and returns histones to a basal state, which is followed by suppression of the activities of gene transcription. There is evidence indicating that HDAC2 controls the functions of key proteins associated with the cell cycle (9). Furthermore, the overexpression of HDAC2 is found in various types of cancer, resulting in the deregulation of uncontrolled proliferation (10). HDAC inhibitors have been reported to exert anticancer activities in multiple types of solid and hematologic malignancies (11,12). The expression levels of HDAC2 are enhanced between non-tumor tissues and histopathological grades of HCC (13). Thus, 1,25(OH)₂D₃

Correspondence to: Dr Yunzhu Huang, Biochemistry Department, Affiliated Hospital of Guizhou Medical University, 28 Guiyi Street, Guiyang, Guizhou 550004, P.R. China E-mail: yunzhugy@163.com

Professor Guozhen Yang, Medical Laboratory, Guizhou Medical University, 9 Beijing Road, Guiyang, Guizhou 550004, P.R. China E-mail: guozheng1@163.com

Key words: $1,25(OH)_2D_3$, histone deacetylase 2, hepatocellular carcinoma, p21(WAF1/Cip1)

may exert antitumor activities in HCC via mediating the levels of HDAC2. Aberrant regulation of HDAC2 deletion leads to the upregulation of p21(WAF1/Cip1) (14), which is one of the well-characterized modulators critical in cell senescence and apoptosis (15). Therefore, the present study investigated the effects of $1,25(OH)_2D_3$ on the expression levels of HDAC2 and p21(WAF1/Cip1). It is possible that $1,25(OH)_2D_3$ exhibits antitumor properties against HCC and may be a potential therapeutic agent for the treatment of HCC.

Materials and methods

Cell lines and cell culture. The HpG2 hepatocellular carcinoma cell line was obtained from China Center for Typical Culture Collection (Wuhan, China) and cultured in Dulbecco's modified Eagle's medium (DMEM; Beijing Solarbio, Beijing, China) supplemented with 10% heat-inactivated fetal bovine serum (GE Healthcare Life Sciences, Logan, UT, USA; cat. no. F4135), 100 U/ml penicillin (Macgene Co., Ltd., Beijing, China), and 100 U/ml streptomycin (Macgene Co., Ltd.). During the experiments, 3x10⁵ cells were plated in culture flasks and grown in a humid atmosphere (37°C; 5% CO₂). The cells were cultured with different concentrations (0, 0.001, 0.01, 0.1, 1 and 10 nM) of 1,25(OH)₂D₃ (Biofriendship, Inc., Beijing, China) or vehicle (dimethyl-sulfoxide).

Cell viability. The viability of the HpG2 cells was measured using a trypan blue uptake assay (Beyotime Institute of Biotechnology, Beijing, China) and a [³H]thymidine incorporation assay (Atom HighTech, Co., Ltd., Beijing, China) following treatment of the cells with different concentrations of 1,25(OH)₂D₃ (0, 0.001, 0.01, 0.1, 1, 10, 100 and 1,000 nM) (Beyotime, Beijing, China) for 24 h. The numbers of trypan blue-positive cells were calculated among 400 cells in two separate microscopic fields using an optical microscope (CX41; Olympus Corporation, Tokyo, Japan). To avoid bias caused by the disappearance of nonviable cells, the total cell number was counted at different times, with the results presented as the percentage of viable cells, reflecting data collected from experiments, compared with the fixed number of total cells. For the [³H]thymidine uptake assay, the cells were counted following exposure to different concentrations of $1,25(OH)_2D_3$, and a 200 μ 1 cell suspension, containing 1.0x10⁵ cells/ml for each condition, were added to each well of 96-well plates (Falcon Labware, Oxnard, CA, USA). Following culture for 24 h, the cells were labeled with 1 Ci/well of [3H]thymidine (67 Ci/mmol) for 4 h, following which the samples were collected using an automated sample harvester (MASH II; Microbiological Associates, Inc., Rockville, MD, USA), dried and measured in 5 ml Aquassure (New England Nuclear; PerkinElmer, Inc., Waltham, MA, USA), and were measured using a Scintillation counter (4810TR; PerkinElmer, Waltham, MA, USA)

Construction of the pcDNA3.1-HDAC2 plasmid. The HDAC2 gene (Accession no. CR541717; http://www.ncbi.nlm. nih.gov/nuccore/CR541717) was amplified using the following primers (Takara Biotechnology Co., Ltd., Dalian, China): Forward 5'-AGTC<u>CATATG</u>GCGCAGACGCAGGGCACC CGGAG-3' and reverse 5'-CTAC<u>GAATTC</u>TCAGGCCAACT

TGACCTCCTTCG-3', generating a 1,449 bp product. The underline highlights the NdeI and EcoRI, respectively The polymerase chain reaction (PCR; 100 µl) contained 20 µl 5X Primer star PCR buffer, 8 µl 25 mM dNTP, 0.5 µl 100 µM fo rward primer, 0.5 μ l 100 μ M reverse primer, 1 μ l template plasmid, 0.5 µl Primer star DNA polymerase (Takara Biotechnology Co., Ltd.) and 34.5 µl distilled water. The following PCR thermocycling steps were performed on Eppendorf 5331 MasterCycler Gradient Thermal Cycler (Hamburg, Germany): 94°C for 5 min; followed by 30 cycles at 94°C for 30 sec, 55°C for 30 sec and 68°C for 2 min; and a final step at 68°C for 10 min. The PCR product was linked to the NheI-EcoRI sites (Takara Biotechnology Co., Ltd.) of a pcDNA3.1 vector (Applied Biosystems; Thermo Fisher Scientific, Inc., Waltham, MA, USA), which was termed pcDNA3.1-HDAC2. The pcDNA3.1-HDAC2 plasmid was amplified in E. coli TOP10 (Takara Biotechnology Co., Ltd.) and isolated using a QIAprep Miniprep kit (Qiagen, Beijing, China), then confirmed using DNA sequencing (Takara Biotechnology Co., Ltd.).

Short hairpin (sh)RNA constructs for HDAC2 gene silencing. The pTZU6+1 plasmids were provided by Chongqing Medical University (Chongqing, China). According to the principles of shRNA design (16), and using the HDAC2 cDNA sequence, 19-21 nt DNA fragments were designed, which spanned the TTGG insertion sequence, with an identical sequence to that of the HDAC2 gene. The HDAC2 coding sequence and the reverse complementary sequence were designed as follows: siHDAC2, forward 5'-GACTGTCGACTC GACCCTCCTTGACTGTACGCCATGTTGGCATGGCGT ACAGTCAAGGAGGTTTTTT-3' and reverse 5'-AGTCTC **TAGACTAGAAAAAACCTCCTTGACTGTACGCCATGC** CAACATGGCGTACAGTCAAGGAGGG-3'. The SalI and XbaI (underlined; Takara Biotechnology Co., Ltd.) restriction sites were added to either end of the oligos for constructing the pTZU 6+1-HDAC2 vectors.

shRNA expression levels of HDAC2 and HDAC2. The cells were transfected with 1 μ g of the HDAC2 or HDAC2-shRNA expression vector. Following 48 h transfection, Geneticin G418 (Invitrogen; Thermo Fisher Scientific, Inc.) was added to the DMEM (500 μ g/ml), and the G418-resistant colonies were selected. The medium was replaced every 2 days. After 3 weeks, the G418-resistant cells were selected and cultured separately.

Reverse transcription-quantitative (RT-qPCR). Total RNA was isolated from the cells using a Takara MiniBEST Universal RNA Extraction kit (Takara Biotechnology Co., Ltd.), according to the manufacturer's protocol. cDNA was amplified using a Transcriptor First Strand cDNA Synthesis kit (Takara Biotechnology Co., Ltd). RT-qPCR was performed on a qTOWER2 (Analytik Jena AG, Jena, Germany), according to the manufacturer's protocol. The following PCR primers were used: HDAC2 (NCBI Reference Sequence: NM_001527.3), forward 5'-GGT GATGGTGTTGAAGAAGC-3' and reverse 5'-GCACTA GGTTGATACATCTC-3'. The glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene (NCBI Reference Sequence:



NM_001289746.1), forward 5'-GATCCCTCCAAA ATCAAGTG-3' and reverse 5'-ATGATCTTGAGGCTG TTGTC-3') was used as a loading control for normalizing data.

Two master mixes (HDAC2 and GAPDH) were used for qPCR analysis. Each mix contained 10 µl SYBR Green (Bioline, Ltd., London, UK), 0.5 µl of each of primer (100 pM), and 8 µl Aqua Dest distilled water/per well of a 96-well plate (Frame Star Fast Plate 96; 4titude, Surrey, UK). A total of 19 µl master mix and 1 μ l cDNA were used, and two replicates for the expression levels of HDAC2 and GAPDH were used for each individual. Consequently, 21 animals were measured per plate, together with 12 non-template controls. These controls contained RNAse-free water instead of cDNA. The plates were sealed using q-stick adhesive for qPCR (4titude) and were centrifuged for a few seconds to ensure all liquid was inside the well (MPS 1000; Labnet, Edison, NJ, USA). Subsequently, the plates were incubated at 95°C for 2 min and run on an Applied Biosystems 7500 Real-Time PCR system (Invitrogen; Thermo Fisher Scientific, Inc.) in 45 cycles of 5 sec at 95°C, follwed by 30 sec at 60°C and 30 sec at 72°C. The mRNA expression levels were normalized against the levels of GAPDH.

Western blot analysis. The cells were lysed using radioimmunoprecipitation assay buffer (Shanghai Sigma High-Tech Co., Ltd., Shanghai, China), containing 20 mM TrisHCl (pH 8.0), 100 mM NaCl, 1% NP-40, 1% sodium deoxycholate, 0.1 % SDS, 0.1 mg/ml phenylmethylsulfonyl fluoride, 1 µg/ml aprotinin and 0.5% sodium orthovanadate. Polyacrylamide gels (10%) for SDS-PAGE (Bio-Rad Laboratories, Inc., Hercules, CA, USA) were used for the separation of protein samples. Following SDS-PAGE, the proteins were transferred onto polyvinylidene difluoride membranes (EMD Millipore, Billerica, MA, USA). Following the blotting, the membranes were blocked with 5% skim milk in Tris-buffered saline with Tween 20 (TBS-T) for 1 h at room temperature. The membranes were incubated with rabbit anti-human anti-HDAC2 antibody (1:1,000; cat. no. SC-7872; Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA) overnight at 4°C. Following three washes with TBS-T, the membranes were incubated with horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG antibody (1:10,000; cat. no. G21234; Thermo Fisher Scientific, Inc.) for 1 h at room temperature. Signals were measured using chemiluminescent HRP substrate (EMD Millipore). The relative expression of HDAC2 was measured using Image J software (Windows version 1.49; National Institutes of Health, Maryland, MD, USA).

Cell growth assay. The HpG2 cells (1×10^4) were cultured in 24-well plates with DMEM, with or without $1,25-(OH)_2D_3$. The number of viable cells were examined by adding 2 mg/ml 3-(4, 5-dimethylthiazol-2-yl)-2,5-diphenyltetrazoliumbromide (Sigma-Aldrich, St. Louis, MO, USA) in each well for 3 days consecutively. The medium was discarded after 2, and the formazan crystals were dissolved in 100 μ l dimethyl-sulfoxide/per well. The absorbance values were read at 590 nm using a microplate reader (WD-2102A; Beijing Liuyi Instrument Factory, Beijing, China).



Figure 1. Effects of $1,25(OH)_2D_3$ toxicity on the viability of HpG2 cells following culture for 24 h. (A) Cell viability of the HpG2 cells was measured using a trypan blue assay. (B) Cell viability of the HpG2 cells was measured using [³H]thymidine incorporation analysis. The cell viabilities shown represent the results of four independent experiments, and data are expressed as the mean ± standard deviation. *P<0.05, vs. 0 nM.

Statistical analysis. All experiments were performed three times and all samples were measured in triplicate. The results are presented as the mean \pm standard deviation). The statistical difference between groups was examined using Student's t-test. Statistical comparisons were made using a two-way analysis of variance. P<0.05 was considered to indicate a statistically significant difference. All data were analyzed using the SPSS 20.0 software package (IBM SPSS, Armonk, NY, USA).

Results

Effects of $1,25-(OH)_2D_3$ toxicity in HpG2 cells. To avoid the presence of cytotoxic effects of $1,25-(OH)_2D_3$ on the HCC cells, the concentration of 1,25-(OH)₂D₃ was investigated prior to performing experiments to assess its effects. Exposure of the HpG2 cells to $1,25-(OH)_2D_3$, in the presence of >1,000 nM, resulted in loss of cell viability, as observed by the uptake of trypan blue (P<0.05). At concentrations <100 nM, the cell viability was >90% (Fig. 1). These effects of $1,25-(OH)_2D_3$ were not noted immediately following addition. The percentage viabilities of the cells in the 1,25-(OH)₂D₃ groups were 100, 98 and 96%, compared with the controls, after 6, 12 and 24 h culture, respectively (Fig. 1). Toxicity was also evaluated by examining the ability of cells to incorporate [³H]thymidine into DNA following exposure to different concentrations of 1,25-(OH)₂D₃. As shown in Fig. 1B, this method was a more sensitive index of cell injury. A significant reduction in [3H]thymidine uptake



Figure 2. Reverse transcription-quantitative polymerase chain reaction analysis determine the mRNA levels of HDAC2 and p21(WAF1/Cip1) in HpG2 cells. The mRNA expression of HDAC2 was normalized using GAPDH. The expression of HDAC2 was induced by pCDNA3.1-HDAC2 transfection or silenced by HDAC2 shRNA in the HpG2 cells. The mRNA levels were measured in four independent experiments, and data are expressed as the mean \pm standard deviation. HDAC2, histone deacetylase 2; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; shRNA, short hairpin RNA.

was measured at concentrations >100 nM (P<0.05). Based on these results, the lower concentrations (0.001, 0.01, 0.1, 1 and 10 nM) of 1,25-(OH)₂D₃ were used for the subsequent culture of HpG2 cells.

Treatment with $1,25(OH)_2D_3$ reduces the mRNA levels of HDAC2 and increase the levels of p21(WAF1/Cip1). To

understand the mechanism underlying the antitumor activity of $1,25(OH)_2D_3$ *in vitro*, the mRNA levels of HDAC2 and p21(WAF1/Cip1) were investigated following treatment with $1,25(OH)_2D_3$. The results showed that $1,25(OH)_2D_3$ treatment reduced the mRNA levels of HDAC2 and increased the levels of p21(WAF1/Cip1) (Fig. 2). The mRNA level of HDAC2 was significantly enhanced when the HpG2 cells were transfected





Figure 3. Western blot analysis determination of the protein levels of HDAC2 and p21(WAF1/Cip1) in HpG2 cells. The protein level of HDAC2 was normalized using GAPDH. The expression of HDAC2 was induced by transfection with pCDNA3.1-HDAC2 or silenced by HDAC2 shRNA in the HpG2 cells. The protein levels were measured in four independent experiments, and the data are expressed as the mean \pm standard deviation. R² values <0.4 were considered to indicate no association between the concentration of 1,25(OH)₂D₃ and the levels of HDAC2 or p21(WAF1/Cip1). HDAC2, histone deacetylase 2; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; shRNA, short hairpin RNA.

with pCDNA3.1-HDAC2. In addition, the mRNA level of p21(WAF1/Cip1) decreased significantly, compared with the cells without HDAC2 transfection. The changes in the levels of

HDAC2 and p21(WAF1/Cip1) were also marginally affected by $1,25(OH)_2D_3$ in a dose-dependent manner, according to the logistic regression (Fig. 2). By contrast, the mRNA expression



Figure 4. Effects of $1,25(OH)_2D_3$ on (A) the growth rate of HpG2 cells. The expression of HDAC2 was (B) induced by transfection with pCDNA3.1-HDAC2 or (C) silenced by HDAC2 shRNA in the HpG2 cells. The growth rate were measured following culture for 24, 48 and 72 h, in four independent experiments. Data are expressed as the mean \pm standard deviation. R² values <0.4 were considered to indicate no association between the concentration of $1,25(OH)_2D_3$ and the levels of HDAC2 or p21(WAF1/Cip1). HDAC2, histone deacetylase 2; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; shRNA, short hairpin RNA.

levels of HDAC2 were almost completely absent when the HpG2 cells were transfected with HDAC2 shRNA, whereas the mRNA expression levels of p21(WAF1/Cip1) were significantly increased, compared with the cells without HDAC2 shRNA transfection. In addition, the change in the expression levels of HDAC2 and p21(WAF1/Cip1) were affected by $1,25(OH)_2D_3$ in a dose-dependent manner, according to the logistic regression (Fig. 2). These results suggested that $1,25(OH)_2D_3$ affected the development of HCC, predominantly by regulating the mRNA levels of HDAC2, which mediated the mRNA level of p21(WAF1/Cip1).

Treatment with $1,25(OH)_2D_3$ reduces the protein levels of HDAC2 and increases the protein levels of p21(WAF1/Cip1). To further investigate the mechanism underlying the antitumor activity of $1,25(OH)_2D_3$ in vitro, the protein expression levels of HDAC2 and p21(WAF1/Cip1) were investigated following treatment with $1,25(OH)_2D_3$. The results showed that $1,25(OH)_2D_3$ treatment reduced the protein levels of HDAC2 and increased the protein levels of p21(WAF1/Cip1); Fig. 3). The protein expression of HDAC2 was significantly enhanced when the HpG2 cells were transfected with pCDNA3.1-HDAC2,

whereas the protein expression of p21(WAF1/Cip1) was decreased significantly, compared with that in the cells without HDAC2 transfection. In addition, the changes in the levels of HDAC2 and p21(WAF1/Cip1) were also marginally affected by $1,25(OH)_2D_3$ in a dose-dependent manner, according to the logistic regression (Fig. 3). By contrast, the protein expression of HDAC2 was almost completely eliminated when the HpG2 cells were transfected with HDAC2 shRNA. The protein levels of p21(WAF1/Cip1) were also significantly increased, compared with those in the cells without HDAC2 shRNA transfection, and the changes in the levels of HDAC2 and p21(WAF1/Cip1) were affected by $1,25(OH)_2D_3$ in a dose-dependent manner, according to the logistic regression (Fig. 3). These results suggested that 1,25(OH)₂D₃ affected the development of HCC, predominantly by regulating the protein levels of HDAC2, which mediated the expression of p21(WAF1/Cip1).

 $1,25(OH)_2D_3$ treatment significantly reduces the growth rate of HCC. To examine the antitumor effects of $1,25(OH)_2D_3$ in HCC, the present study examined its antiproliferative effects in the human HpG2 HCC line. The cells were incubated with different concentrations of $1,25(OH)_2D_3$ for 3 days and



the status of cell growth was measured. Consistent with previous evidence (17), cell growth in the $1,25(OH)_2D_3$ group was decreased in a dose-dependent manner, compared with the control group. The growth curve line (Fig. 4A) showed a significant proliferation inhibitory effect of $1,25(OH)_2D_3$, which occurred in a dose-dependent manner.

The average growth rate of the pCDNA3.1-HDAC2-transfected HpG2 cells was increased, compared with the untransfected cells (Fig. 4). Additionally, expression of HDAC2 in the HpG2 cells reversed the inhibitory effect of $1,25(OH)_2D_3$ (Fig. 4). The average growth rate was also enhanced in the cells treated with a high concentration of $1,25(OH)_2D_3$ (10 nM/day). Of note, there was difference in growth rate between the cells treated with $1,25(OH)_2D_3$ and those without treatment when the expression of HDAC2 was induced by transfection (Fig. 4), although the inhibitory effects of HDAC2 remained, according to the logistic regression. Following shRNA transfection, the average growth rate of the HpG2 cells was reduced. Additionally, HDAC2 shRNA transfection of the HpG2 cells had a similar function as the inhibitory effect of $1,25(OH)_2D_3$ (Fig. 4). The average growth rate was also decreased, although the cells was not treated with high concentration of 1,25(OH)₂D₃. Of note, there was a difference in growth rate between the cells treated with $1,25(OH)_2D_3$ and the untreated cells when HDAC2 was silenced (Fig. 4), although the inhibitory effects of HDAC2 remained, according to the logistic regression.

Discussion

At present, epidemiological evidence indicates that decreased concentrations of vitamin D are associated with an enhanced risk of various types of cancer (18). Although previous studies have indicated that $1,25(OH)_2D_3$ inhibits the proliferation of HCC in a dose dependent manner (19), the detailed molecular mechanism underlying the function of $1,25(OH)_2D_3$ in preventing the progression of HCC remains to be fully elucidated.

Prior to commencement of the present study, the potential toxicity of high concentrations of $1,25(OH)_2D_3$ on HCC cells were examined. As observed inhibitory effects may be caused by toxicity, rather than by other molecular mechanisms, the toxicity of $1,25(OH)_2D_3$ was first measured using trypan blue and [³H]thymidine incorporation assays. As expected, the viability of the HCC cells was significantly affected when the concentration of $1,25(OH)_2D_3$ was >100 nM (Fig. 1). Therefore, the concentration used in the subsequent experiments was ≤10 nM. This ensured that any inhibitory effects were not caused by toxicity, but by other molecular mechanisms.

To investigate the molecular mechanism underlying the inhibitory effect of $1,25(OH)_2D_3$ in HCC, the present study transfected HCC cells with HDAC2. The resulting high expression levels of HDAC2 led to a significant reduction in the inhibitory effects of $1,25(OH)_2D_3$ on the HCC cells (Fig. 4). In addition, the expression of p21(WAF1/Cip1) was reduced. By contrast, when the HCC cells were transfected with HDAC2 shRNA, leading to silencing of the HDAC2 gene, similar inhibitory effects on HCC cell growth were observed in the cells, which were not treated with $1,25(OH)_2D_3$.

Furthermore, exogenous $1,25(OH)_2D_3$ treatment decreased the expression levels of HDAC2 and enhanced the expression levels of p21(WAF1/Cip1) in a dose-dependent manner. Therefore, $1,25(OH)_2D_3$ inhibited the development of HCC via the downregulation of HDAC2, which mediated the levels of p21(WAF1/Cip1).

There is increasing evidence reinforcing the hypothesis that excessively and chronically expressed HDAC2 contributes to the development of HCC (13,14). HDAC2 is a major deacetylase, and its expression is inversely associated with changes in the acetylation state of histone H4K5. RNA interference-mediated gene silencing of HDAC2 leads to hyperacetylation of histone H4 and a developmental delay, although the levels of HDAC3 remain high (20,21). Increased expression levels of p21(WAF1/Cip1) may contribute to the observed developmental delay. HDAC2 regulates a number of biological processes in cells. Although HDAC2 inhibits the growth of the liver in aged mice, the levels of HDAC2 are also increased in HCC (10). Increased levels of HDAC2 lead to its interaction with p21(WAF1/Cip1) via the Sp1 binding sites of the p21(WAF1/Cip1) gene promoter, and inhibition of acetylated histone H3 on these sites (22). Histone H3 is a conserved protein in the nucleus, which can be readily modified post-translationally, and the state of acetylation shows clinical diagnostic significance in HepG2 cells. The levels of acetylated peptides are associated with malignant transformation in the liver (23). The results of the present study indicated that 1,25(OH)₂D₃ inhibited HDAC2, and its downregulation was involved in the upregulation of p21(WAF1/Cip1), which resulted in the inhibition of HCC proliferation.

Although $1,25(OH)_2D_3$ has potent inhibitory effects in several liver diseases (24,25), the molecular mechanism underlying the inhibitory effect of $1,25(OH)_2D_3$ in HCC development remains to be fully elucidated. The present study demonstrated the novel finding that downregulating the expression of HDAC2 resulted in an increase in the expression of p21(WAF1/Cip1), which eventually inhibited HCC development. HDAC2 ablation suppressed HCC development and enhanced the expression of p21(WAF1/Cip1), which was similar to the inhibitory effects of $1,25(OH)_2D_3$ on HCC (Figs. 2-4). In addition, elevated levels of HDAC2 promoted the development of HCC and decreased the activity of p21(WAF1/Cip1), which reversed the inhibitory effects of $1,25(OH)_2D_3$ on the development of HCC, even when the cells were treated with a high dose of $1, 25(OH)_2D_3$. Treatment with $1,25(OH)_2D_3$ led to the downregulation in the expression of HDAC2, which mediated the level of p21(WAF1/Cip1). Further investigations of the effects of 1,25(OH)₂D₃ on hepatocarcinogenesis may provide further insights into the detailed molecular mechanism underlying vitamin D in antitumor activity. The results of the present study support the potential development of $1,25(OH)_2D_3$ as an effective drug for the treatment of HCC.

Acknowledgements

The present study was supported by the joint fund of the Science and Technology Department of Guizhou Province and Guiyang Medical College (grant no. LH(2014)7114).

References

- Tang B, Zhang Y, Liang R, Gao Z, Sun D and Wang L: RNAi-mediated EZH2 depletion decreases MDR1 expression and sensitizes multidrug-resistant hepatocellular carcinoma cells to chemotherapy. Oncol Rep 29: 1037-1042, 2013.
- Kirikoshi H, Yoneda M, Mawatari H, Fujita K, Imajo K, Kato S, Suzuki K, Kobayashi N, Kubota K, Maeda S, *et al*: Is hepatic arterial infusion chemotherapy effective treatment for advanced hepatocellular carcinoma resistant to transarterial chemoembolization? World J Gastroenterol 18: 1933-1939, 2012.
- Llovet JM, Bustamante J, Castells A, Vilana R, Ayuso Mdel C, Sala M, Brú C, Rodés J and Bruix J: Natural history of untreated nonsurgical hepatocellular carcinoma: rationale for the design and evaluation of therapeutic trials. Hepatology 29: 62-67, 1999.
- Portolani N, Coniglio A, Ghidoni S, Giovanelli M, Benetti A, Tiberio GA and Giulini SM: Early and late recurrence after liver resection for hepatocellular carcinoma: Prognostic and therapeutic implications. Ann Surg 243: 229-235, 2006.
 Lu L, Sun HC, Zhang W, Chai ZT, Zhu XD, Kong LQ, Wang WQ, Zhang KZ, Zhang YY, Zhang QB, *et al*: Aspirin minimized
- 5. Lu L, Sun HC, Zhang W, Chai ZT, Zhu XD, Kong LQ, Wang WQ, Zhang KZ, Zhang YY, Zhang QB, *et al*: Aspirin minimized the pro-metastasis effect of sorafenib and improved survival by up-regulating HTATIP2 in hepatocellular carcinoma. PLoS One 8: e65023, 2013.
- 6. Deeb KK, Trump DL and Johnson CS: Vitamin D signalling pathways in cancer: Potential for anticancer therapeutics. Nat Rev Cancer 7: 684-700, 2007.
- 7. Gocek E and Studzinski GP: Vitamin D and differentiation in cancer. Crit Rev Clin Lab Sci 46: 190-209, 2009.
- Bhalla AK, Amento EP, Clemens TL, Holick MF and Krane SM: Specific high-affinity receptors for 1,25-dihydroxyvitamin D3 in human peripheral blood mononuclear cells: Presence in monocytes and induction in T lymphocytes following activation. J Clin Endocrinol Metab 57: 1308-1310, 1983.
- Jung KH, Noh JH, Kim JK, Eun JW, Bae HJ, Xie HJ, Chang YG, Kim MG, Park H, Lee JY and Nam SW: HDAC2 overexpression confers oncogenic potential to human lung cancer cells by deregulating expression of apoptosis and cell cycle proteins. J Cell Biochem 113: 2167-2177, 2012.
- 10. Kim HS, Chang YG, Bae HJ, Eun JW, Shen Q, Park SJ, Shin WC, Lee EK, Park S, Ahn YM, *et al*: Oncogenic potential of CK2α and its regulatory role in EGF-induced HDAC2 expression in human liver cancer. FEBS J 281: 851-861, 2014.
- 11. Shimizu T, LoRusso PM, Papadopoulos KP, Patnaik A, Beeram M, Smith LS, Rasco DW, Mays TA, Chambers G, Ma A, *et al*: Phase I first-in-human study of CUDC-101, a multitargeted inhibitor of HDACs, EGFR and HER2 in patients with advanced solid tumors. Clin Cancer Res 20: 5032-5040, 2014.
- 12. Wang L, Zou X, Berger AD, Twiss C, Peng Y, Li Y, Chiu J, Guo H, Satagopan J, Wilton A, *et al*: Increased expression of histone deacetylaces (HDACs) and inhibition of prostate cancer growth and invasion by HDAC inhibitor SAHA. Am J Transl Res 1: 62-71, 2009.

- 13. Noh JH, Bae HJ, Eun JW, Shen Q, Park SJ, Kim HS, Nam B, Shin WC, Lee EK, Lee K, *et al*: HDAC2 provides a critical support to malignant progression of hepatocellular carcinoma through feedback control of mTORC1 and AKT. Cancer Res 74: 1728-1738, 2014.
- 14. Noh JH, Jung KH, Kim JK, Eun JW, Bae HJ, Xie HJ, Chang YG, Kim MG, Park WS, Lee JY and Nam SW: Aberrant regulation of HDAC2 mediates proliferation of hepatocellular carcinoma cells by deregulating expression of G1/S cell cycle proteins. PLoS One 6: e28103, 2011.
- Parra E, Gutiérrez L and Ferreira J: Increased expression of p21Waf1/Cip1 and JNK with costimulation of prostate cancer cell activation by an siRNA Egr-1 inhibitor. Oncol Rep 30: 911-916, 2013.
- Sachidanandam R: RNAi: design and analysis. Current Protocols in Bioinformatics: 13: 10-11, 2004.
- Abe J, Nakano T, Nishii Y, Matsumoto T, Ogata E and Ikeda K: A Novel Vitamin-D3 Analog, 22-Oxa-1,25-Dihydroxyvitamin-D3, Inhibits the Growth of Human Breast-Cancer Invitro and Invivo without Causing Hypercalcemia. Endocrinology 129: 832-837, 1991.
- 18. Giovannucci E: Vitamin D and cancer incidence in the harvard cohorts. Ann Epidemiol 19: 84-88, 2009.
- Pourgholami MH, Akhter J, Lu Y and Morris DL: In vitro and in vivo inhibition of liver cancer cells by 1,25-dihydroxyvitamin D3. Cancer Lett 151: 97-102, 2000.
- Methot JL, Hoffman DM, Witter DJ, Stanton MG, Harrington P, Hamblett C, Siliphaivanh P, Wilson K, Hubbs J, Heidebrecht R, *et al*: Delayed and prolonged histone hyperacetylation with a selective HDAC1/HDAC2 inhibitor. ACS Med Chem Lett 5: 340-345, 2014.
- 21. Ma P and Schultz RM: Histone deacetylase 2 (HDAC2) regulates chromosome segregation and kinetochore function via H4K16 deacetylation during oocyte maturation in mouse. PLoS Genet 9: e1003377, 2013.
- 22. Di Padova M, Bruno T, De Nicola F, Iezzi S, D'Angelo C, Gallo R, Nicosia D, Corbi N, Biroccio A, Floridi A, *et al*: Che-1 arrests human colon carcinoma cell proliferation by displacing HDAC1 from the p21WAF1/CIP1 promoter. J Biol Chem 278: 36496-36504, 2003.
- 23. Wang W, Xu L, Kong J, Fan H and Yang P: Quantitative research of histone H3 acetylation levels of human hepatocellular carcinoma cells. Bioanalysis 5: 327-339, 2013.
- 24. Küçükazman M, Ata N, Dal KA, Yeniova AÖ, Kefeli A, Basyigit S, Aktas B, Akin KO, Ağladıoğlu K, Üre ÖS, *et al*: The association of vitamin D deficiency with non-alcoholic fatty liver disease. Clinics (Sao Paulo) 69: 542-546, 2014.
- 25. Mandorfer M, Payer BA, Schwabl P, Steiner S, Ferlitsch A, Aichelburg MC, Stättermayer AF, Ferenci P, Obermayer-Pietsch B, Grabmeier-Pfistershammer K, *et al*: Revisiting liver disease progression in HIV/HCV-coinfected patients: The influence of vitamin D, insulin resistance, immune status, IL28B and PNPLA3. Liver Int 35: 876-885, 2014.