

Tanshinone IIA may inhibit the growth of small cell lung cancer H146 cells by up-regulating the Bax/Bcl-2 ratio and decreasing mitochondrial membrane potential

CHUN-YUAN CHENG^{1,2} and CHIN-CHENG SU^{3,4}

¹Institute of Medicine of Chung Shan Medical University; ²Changhua Christian Hospital; ³Tzu-Chi University, Hualien 97004; ⁴Division of General Surgery, Buddhist Tzu-Chi General Hospital, Hualien 97004, Taiwan, R.O.C.

Received January 4, 2010; Accepted April 30, 2010

DOI: 10.3892/mmr_00000310

Abstract. Tanshinone IIA (Tan-IIA) may inhibit the growth of human non-small cell lung cancer A549 cells. However, the molecular mechanisms behind this malignancy have yet to be established. In the present study, we examined the effects of Tan-IIA on human small cell lung cancer H146 cells *in vitro*. The cytotoxicity of Tan-IIA in H146 cells was measured using the MTT assay. Mitochondrial membrane potential (MMP), reactive oxygen species (ROS) and Ca²⁺ in H146 cells were detected by flow cytometry, and the protein expression of Bax, Bcl-2, Caspase-3, NF-κBp65, GADD153 and β-actin in H146 cells was measured by Western blotting. H146 cells were inhibited in a dose-dependent manner. The protein expression of GADD153 and Caspase-3 was increased, but the proto-oncogene bcl-2 was notably decreased in H146 cells treated with Tan-IIA (5 μg/ml) for 24 h. FACS showed that Tan-IIA may increase the production of ROS and Ca²⁺, but decreases MMP. The results indicate that Tan-IIA is capable of inhibiting the proliferation of H146 cells. One of the molecular mechanisms behind this effect may be the induction of ROS release and the decrease in MMP caused by an increase in the Bax/Bcl-2 ratio. Another may involve endoplasmic reticulum stress caused by the release of Ca²⁺ and an increase in GADD153 expression followed by a decrease in Bcl-2 expression, which induces a higher ratio of Bax/Bcl-2, in turn causing a decrease in MMP and leading to an increase in Caspase-3 expression and the inhibition of H146 cells. Thus, Tan-IIA may be a promising novel chemotherapeutic agent for the treatment of human small cell lung cancer H146 cells.

Introduction

Danshen (*Salviae miltiorrhizae Radix*) was first recorded in the classical treatise Shen Nong Pen Tsao Ching (circa A.D. 100) (1), and is clinically administered to many cancer patients following chemotherapy or radiotherapy in Taiwan. However, its functions are controversial and have not been adequately explored. Lung cancer is the leading cause of cancer related death in Taiwan and in Western countries (2). However, the efficacy of chemotherapy for lung cancer remains unsatisfactory. Tanshinone IIA (Tan-IIA) (C19H18O3) is extracted from Danshen, *Salviae miltiorrhizae Radix* (3,4). There have been extensive investigations into the potential mechanisms of Tan-IIA as an anti-tumor agent in various tumor cells, including leukemia (5,6), breast cancer (7,8), colon cancer (9) and hepatocellular carcinoma cells (10,11). Tan-IIA may inhibit the proliferation of lung cancer NCI-H460 cells by decreasing Bcl-2 expression and inducing apoptosis (12). Our previous study also revealed that Tan-IIA inhibits the growth of non small cell lung cancer A549 cells by decreasing mitochondrial membrane potential (MMP) and increasing the ratio of Bax/Bcl-2 (13). However, the effects of Tan-IIA in human small cell lung cancer have yet to be established. In the present study, we investigated the molecular mechanisms of Tan-IIA in human small cell lung cancer H146 cells.

Materials and methods

Chemicals and reagents. Tan-IIA (Molecular Formula C19H18O3, >96% HPLC) was purchased from Herbasin Co. (Shenyang, China). Aprotinin, antipain, sodium deoxycholate, leupeptin, propidium iodide, sodium orthovanadate, Triton X-100, Tris-HCl, ribonuclease-A and MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Dimethyl sulfoxide (DMSO), potassium phosphates and TE buffer were purchased from Merck Co. (Darmstadt, Germany). RPMI 1640 medium, fetal bovine serum (FBS), penicillin-streptomycin, trypsin-EDTA, and glutamine were obtained from Gibco BRL (Grand Island, NY, USA). 10X SDS-PAGE running buffer, Tris, Tween 20, SDS, 5X TBE buffer were obtained from Amresco. (St. Louis, MO, USA);

Correspondence to: Dr Chin-Cheng Su, Division of General Surgery, Breast-medicine center, Buddhist Tzu Chi General Hospital, 707, S.3, Chung-Yang Road, Hualien City 97004, Taiwan, R.O.C.
E-mail: succ.maeva@msa.hinet.net

Key words: tanshinone IIA, H146 cells, reactive oxygen species, Bax/Bcl-2 ratio, mitochondrial membrane potential

BioMax Film was obtained from Kodak. DioC6 (84715) and DCFH-DA (C6827) were obtained from Biocompare (CA, USA). Anti- β -actin (MAB1501), anti-GADD153 and goat anti-mouse IgG horseradish peroxidase conjugated antibody (AP124P) were obtained from Chemicon (MA, USA). Anti-Bcl-2 (N-19-sc-492) and anti-Bax (N-19-sc-492) were from Santa Cruz Biotechnology, Inc.

Cell line and culture. Human lung cancer H146 cells were obtained from the Food Industry Research and Development Institute (Hsinchu, Taiwan). The H146 cells were placed in 25-cm² tissue culture flasks and maintained in RPMI 1640 containing 10% heat-inactivated FBS (Gibco BRL, Grand Island, NY, USA), 100 U/ml penicillin and 100 μ g/ml streptomycin. Cells were grown at 37°C in a humidified atmosphere of 95% air and 5% CO₂. All data presented are from at least three independent experiments.

Determination of cell viability. The H146 cells were plated at a density of 1x10⁵ cells/well and grown for 24 h. Various concentrations of Tan-IIA were added and the cells were grown for 24, 48 and 72 h. For the control regimen, DMSO 0.2% (solvent) was used. After 24, 48 and 72 h of culture, cell viability was evaluated by the MTT assay (14), conducted in triplicate. Briefly, H146 cells were seeded in a 6-well plate at a density of 1x10⁵ cells/well and allowed to adhere overnight. After removing the medium, 2,000 μ l of fresh medium per well, containing 10 mM HEPES and 200 μ l of MTT, was added to the wells. The plate was incubated for 30 min at 37°C in the dark, then the medium was removed and 1,000 μ l of DMSO was added to the wells. Absorbance was measured using an ELISA plate reader at 590 nm.

Detection of reactive oxygen species. The levels of reactive oxygen species (ROS) in the H146 cells were examined and determined by flow cytometry (Becton Dickinson FACSCalibur) as previously described (13). Briefly, ~2x10⁵ cells/well of H146 cells in 12-well plates were incubated with or without Tan-IIA for different time periods (0, 6, 12 and 24 h) to detect changes in ROS. The cells were harvested and washed twice, re-suspended in 500 μ l of 2,7-dichlorodihydrofluorescein diacetate (DCFH-DA) (10 μ M) and incubated at 37°C for 30 min, then analyzed by flow cytometry.

Detection of mitochondrial membrane potential. The mitochondrial membrane potential (MMP) levels in H146 cells were determined by flow cytometry (Becton Dickinson FACSCalibur) as previously described (13). Briefly, ~2x10⁵ cells/well of H146 cells in 12-well plates were incubated with or without Tan-IIA for different time periods (0, 6, 12 and 24 h) to determine changes in MMP. Cells were harvested and washed twice, re-suspended in 500 μ l of DiOC6 (4 μ mol/l) and incubated at 37°C for 30 min, then analyzed by flow cytometry.

Determination of Ca²⁺ concentrations. Ca²⁺ levels in the H146 cells were determined by flow cytometry (Becton Dickinson FACSCalibur) as previously described (13). Briefly, ~2x10⁵ cells/well of H146 cells in 12-well plates were incubated




Figure 1. H146 cells were treated with increasing Tan-IIA concentrations of 0, 0.5, 1, 2 and 4 μ g/ml for different time periods (24, 48 and 72 h). The cytotoxicity of Tan-IIA in H146 cells was determined using the MTT assay. Tan-IIA significantly inhibited H146 cell growth. Each point is the mean \pm SD of three experiments. *P<0.05.

with or without Tan-IIA for different time periods (0, 6, 12 and 24 h) to detect changes in Ca²⁺ levels. Cells were harvested and washed twice, re-suspended in Indo 1/AM (3 μ g/ml), incubated at 37°C for 30 min, then analyzed by flow cytometry.

Western blotting. To determine the effect of Tanshinone IIA on P53, Bcl-2, Bax, Caspase-3, GADD153, NF- κ Bp65 and β -actin levels in H146 cells, Western blotting was used. After treatment with or without various concentrations of Tan-IIA (0, 1, 2 and 5 μ g/ml) for 24 h, total proteins were collected from H146 cells and GADD153 and β -actin levels were examined by sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE) and Western blotting, as described previously (15,16). After treatment with or without various concentrations of Tan-IIA (0, 1, 2.5 and 5 μ g/ml) for 24 h, total proteins were collected from H146 cells and P53, Bcl-2, Bax, Caspase-3, NF- κ Bp65 and β -actin levels were examined by SDS-PAGE and Western blotting, as described previously (15,16).

Statistical analysis. Values are presented as the percentage (mean \pm SD) of the control. The Student's t-test was used to analyze the statistical significance between the Tan-IIA-treated and control groups. A P-value of <0.05 was considered significant for all tests.

Results

Cytotoxic effects of Tanshinone IIA in H146 cells. H146 cells were cultured with various concentrations of Tan-IIA (control, 0.5, 1, 2 and 4 μ g/ml) for different time periods (24, 48 and 72 h), then the cytotoxicity of Tan-IIA in H146 cells was determined using the MTT assay. The percentage of viable cells (relative to the control) was 58.97 \pm 0.53, 41.59 \pm 0.25, 33.57 \pm 0.2 and 23.64 \pm 0.1%, respectively, at 24 h, 81.22 \pm 1.05, 64.45 \pm 0.4, 42.56 \pm 1.12 and 29.11 \pm 0.1%, respectively, at 48 h and 60.27 \pm 0.61, 22.32 \pm 0.46, 14.39 \pm 0.06 and 8.67 \pm 0.17%, respectively, at 72 h. The IC₅₀ was 0.82, 0.96 and 0.69 μ g/ml, respectively, for Tan-IIA to affect H146 cells at different time periods of treatment (24, 48 and 72 h). The results indicate that Tan-IIA inhibited the proliferation of H146 cells in a dose-dependent manner (Fig. 1).




Figure 2. Flow cytometric analysis of reactive oxygen species (ROS) in H146 cells treated with Tan-IIA. Approximately 2×10^5 cells/well of H146 cells in 12-well plates were incubated with Tan-IIA for different time periods (0, 6, 12, and 24 h) to detect changes in ROS. The percentage of cells stained by DCFH-DA dye was determined by flow cytometry. Each point is the mean \pm SD of three experiments. * $P < 0.05$.




Figure 4. Flow cytometric analysis of Ca^{2+} concentration in H146 cells treated with Tan-IIA. Approximately 2×10^5 cells/well of H146 cells in 12-well plates were incubated with Tan-IIA for different durations (0, 6, 12, and 24 h) to detect changes in Ca^{2+} levels. The percentage of cells that were stained by Indo-1/AM dye was determined by flow cytometry. Each point is the mean \pm SD of three experiments. * $P < 0.05$.




Figure 3. Flow cytometric analysis of mitochondrial membrane potential (MMP) in H146 cells treated with Tan-IIA. Approximately 2×10^5 cells/well of H146 cells in 12-well plates were incubated with Tan-IIA for different time periods (0, 6, 12, and 24 h) to determine changes in MMP. The percentage of cells stained by DiOC6 dye was determined by flow cytometry. Each point is the mean \pm SD of three experiments. * $P < 0.05$.

Tanshinone IIA induced the production of reactive oxygen species in H146 cells. Following treatment of the H146 cells with or without $1 \mu\text{g/ml}$ of Tan-IIA for various time periods (0, 6, 12 and 24 h), the increasing percentages of ROS were 5.32 ± 2.11 , 18.32 ± 3.25 , 58.62 ± 4.78 and $87.66 \pm 2.14\%$, respectively. Increasing the co-treated time periods resulted in an increase in ROS release (Fig. 2A and B). These results showed that Tan-IIA promoted ROS production in a time-dependent manner.

Tanshinone-IIA suppressed mitochondrial membrane potential in H146 cells. H146 cells were treated with or without Tan-IIA ($1 \mu\text{g/ml}$) for various time periods (0, 6, 12 and 24 h), then the MMP was analyzed by flow cytometry. The decreasing percentages of the MMP were 2.38 ± 0.19 , 2.98 ± 3.14 , 23.32 ± 4.87 and $44.33 \pm 5.10\%$, respectively. Increasing the duration of incubation led to the decrease of MMP in the H146 cells (Fig. 3A and B).

Tanshinone IIA induced Ca^{2+} production in H146 cells. H146 cells were treated with or without $1 \mu\text{g/ml}$ Tan-IIA for various time periods (0, 6, 12 and 24 h). The increasing percentages of Ca^{2+} concentrations were 1.09 ± 0.98 , 2.09 ± 1.00 , 2.87 ± 2.69 and $18.32 \pm 5.12\%$, respectively. Increasing the duration of incubation led to an increase in the concentration of Ca^{2+} in H146 cells (Fig. 4A and B).

Effect of Tanshinone IIA on NF- κ Bp65, Bcl-2, Bax, Caspase-3, GADD153 and β -actin levels by Western blotting. H146 cells ($5 \times 10^6/\text{ml}$) were treated with various concentrations of Tan-IIA (1, 2 and $5 \mu\text{g/ml}$) for 24 h. Once the total proteins




Figure 5. Protein expression of GADD153, NF- κ Bp65 and β -actin in H146 cells. H146 cells ($5 \times 10^6/\text{ml}$) were treated with various concentrations of Tan-IIA (0, 1, 2 and 5 $\mu\text{g/ml}$) for 24 h before being harvested by centrifugation. Total protein was prepared and determined, then levels of GADD153 and β -actin expression were estimated by Western blotting. Protein expression of GADD153 was significantly increased after treatment with Tan-IIA (Fig. 5A). Tan-IIA decreased the protein expression of NF- κ Bp65 in H146 cells at the concentrations of 1, 2 and 5 $\mu\text{g/ml}$ 24 h (Fig. 5B). Each point is the mean \pm SD of three experiments. * $P < 0.05$.

were prepared and determined, the levels of GADD153 and β -actin expression were estimated by Western blotting. Tan-IIA increased the protein expression of GADD153 in the H146 cells at concentrations of 1, 2 and 5 $\mu\text{g/ml}$ at 24 h (Fig. 5A). Tan-IIA decreased the protein expression of NF- κ Bp65 in the H146 cells at concentrations of 1, 2 and 5 $\mu\text{g/ml}$ at 24 h (Fig. 5B). The H146 cells ($5 \times 10^6/\text{ml}$) were treated with various concentrations of Tan-IIA (1, 2.5 and 5 $\mu\text{g/ml}$) for 24 h. After the total proteins were prepared and determined, the levels of Bax and Bcl-2 expression were estimated by Western blotting. The protein expression of Bax (Fig. 6A) was mildly decreased




Figure 6. Protein expression of Bax, Bcl-2 and β -actin in H146 cells. H146 cells ($5 \times 10^6/\text{ml}$) were treated with various concentrations of Tan-IIA (1, 2.5 and 5 $\mu\text{g/ml}$) for 24 h. Total protein was prepared and determined, then levels of Bax and Bcl-2 expression were estimated. The protein expression of Bax was mildly decreased after treatment with Tan-IIA for 24 h (Fig. 6A). The protein expression of Bcl-2 was significantly decreased after treatment with Tan-IIA for 24 h (Fig. 6B), while the ratio of Bax/Bcl-2 was significantly increased (Fig. 6C). Each point is the mean \pm SD of three experiments. * $P < 0.05$.




Figure 7. Protein expression of Caspase-3 and β -actin in H146 cells. The H146 cells ($5 \times 10^6/ml$) were treated with various concentrations of Tan-IIA (1, 2.5 and 5 $\mu g/ml$) for 24 h. Total protein was prepared and determined, then levels of Caspase-3 expression were estimated by Western blotting. The protein expression of Caspase-3 was significantly increased after treatment with Tan-IIA for 24 h. Each point is the mean \pm SD of three experiments. * $P < 0.05$.

after treatment with Tan-IIA for 24 h. The protein expression of Bcl-2 (Fig. 6B) was significantly decreased after treatment with Tan-IIA for 24 h, but the ratio of Bax/Bcl-2 was significantly increased (Fig. 6C). The protein expression of Caspase-3 was significantly increased after treatment with Tan-IIA for 24 h (Fig. 7).

Discussion

The results of the present study indicate that Tan-IIA may inhibit the proliferation of small cell lung cancer H146 cells in a dose-dependent manner. The mitochondria have been shown to be the central control point of apoptosis (17). It is well documented that decreases in MMP induce apoptosis (18). The overexpression of Bax may decrease MMP; however, it is well known that the overexpression of Bcl-2 prevents the decrease of MMP (19). Tan-IIA may increase Caspase-3 protein expression in H146 cells. This is in agreement with a previous study (12). Our results show that Tan-IIA decreased the protein expression of Bax and Bcl-2. This is not in agreement with a previous study that showed that Tan-IIA up-regulated Bax but down-regulated Bcl-2 protein expression in SPC-A-1 cells (20). In the present study, the results revealed that Tan-IIA increased the ratio of Bax/Bcl-2 and decreased MMP significantly. Tan-IIA increased ROS generation significantly after 6 h, but decreased MMP significantly after 12 h. On the basis of these results, we suggest that Tan-IIA may act through the induction of ROS and by increasing the ratio of Bax/Bcl-2, leading to a decrease in MMP, which in turn induces apoptosis in H146 cells. Endoplasmic reticulum (ER) stress is one apoptotic cell death pathway. It is well documented that the overexpression of GADD153 represents a new and important strategy in anticancer therapy (21). GADD153 is indicative of DNA damage and ER stress, and




Figure 8. Proposed model for the molecular mechanisms of Tan-IIA inhibition of the proliferation of H146 cells. One of the molecular mechanisms may be through the induction of ROS release and the decrease in MMP due to an increase in the Bax/Bcl-2 ratio. Another may be through ER stress caused by the release of Ca^{2+} and an increase in GADD153 expression followed by a decrease in Bcl-2 expression, leading to the induction of a higher ratio of Bax/Bcl-2 and then a decrease in MMP, resulting in increased Caspase-3 expression and the inhibition of H146 cells.

was promoted in previous studies, where it induced apoptosis (22). Several studies have shown that GADD153 suppresses the activation of Bcl-2 (23,24). Our results also reveal that Tan-IIA is capable of increasing the protein expression of GADD153 in H146 cells. On the other hand, the marked release of Ca^{2+} was observed in the H146 cells at 24 h. One of the molecular mechanisms for the Tan-IIA inhibition of H146 cells may be through ER stress caused by the release of Ca^{2+} and an increase in the protein expression of GADD153 followed by a decrease in Bcl-2 protein expression, which induces a higher ratio of Bax/Bcl-2 then a decrease in MMP, leading to an increase in the protein expression of Caspase-3 and the inhibition of the proliferation of H146 cells. This proposed model is the first report regarding the action of Tan-IIA in small cell lung cancer H146 cells (Fig. 8). Tan-IIA appears to be a promising new chemotherapeutic agent for small cell lung cancer.

Acknowledgements

This study was supported by grants CCMP95-RD-206, CCMP97-RD-011 and CCMP98-RD-102 from the Committee on Chinese Medicine and Pharmacy, Department of Health, Executive Yuan, Taiwan, R.O.C.

References

1. Bensky D and Gamble A: Chinese Herbal Medicine: Materia Medica. Revised edition. Eastland Press, Seattle, WA, 1993.
2. Parkin DM, Pisani P and Ferlay J: Global cancer statistics. CA Cancer J Clin 49: 33-64, 1999.
3. Che AJ, Zhang JY, Li CH, Chen XF, Hu ZD and Chen XG: Separation and determination of active components in Radix Salviae miltiorrhizae and its medicinal preparations by nonaqueous capillary electrophoresis. J Sep Sci 27: 569-575, 2004.

4. Zhou L, Zuo Z and Chow MS: Danshen: an overview of its chemistry, pharmacology, pharmacokinetics, and clinical use. *J Clin Pharmacol* 45: 1345-1359, 2005.
5. Sung HJ, Choi SM, Yoon Y and An KS: Tanshinone IIA, an ingredient of *Salvia miltiorrhiza* Bunge, induces apoptosis in human leukemia cell lines through the activation of caspase-3. *Exp Mol Med* 31: 174-178, 1999.
6. Liu JJ, Lin DJ, Liu PQ, Huang M, Li XD and Huang RW: Induction of apoptosis and inhibition of cell adhesive and invasive effects by tanshinone IIA in acute promyelocytic leukemia cells in vitro. *J Biomed Sci* 13: 813-823, 2006.
7. Wang X, Wei Y, Yuan S, *et al*: Potential anticancer activity of tanshinone IIA against human breast cancer. *Int J Cancer* 116: 799-807, 2005.
8. Su CC and Lin YH: Tanshinone IIA inhibits human breast cancer cells through increased Bax to Bcl-xL ratios. *Int J Mol Med* 22: 357-361, 2008.
9. Su CC, Chen GW, Kang JC and Chan MH: Growth inhibition and apoptosis induction by tanshinone IIA in human colon adenocarcinoma cells. *Planta Med* 74: 1357-1362, 2008.
10. Yuan SL, Wei YQ, Wang XJ, Xiao F, Li SF and Zhang J: Growth inhibition and apoptosis induction of tanshinone II-A on human hepatocellular carcinoma cells. *World J Gastroenterol* 10: 2024-2028, 2004.
11. Li Q, Wang Y, Feng N, Fan Z, Sun J and Nan Y: Novel polymeric nanoparticles containing tanshinone IIA for the treatment of hepatoma. *J Drug Target* 16: 725-732, 2008.
12. Hu H, Zhang Y, Huang F and Deng H: Inhibition of proliferation and induction of apoptosis by tanshinone II A in NCI-H460 cell. *Zhong Yao Cai* 28: 301-304, 2005.
13. Chiu TL and Su CC: Tanshinone IIA induces apoptosis in human lung cancer A549 cells through the induction of reactive oxygen species and decreasing the mitochondrial membrane potential. *Int J Mol Med* 25: 231-236, 2010.
14. Ossman T: Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 65: 55-63, 1983.
15. Su CC and Lin YH: Tanshinone IIA inhibits human breast cancer cells through increased Bax to Bcl-xL ratios. *Int J Mol Med* 22: 357-361, 2008.
16. Su CC, Chen GW and Lin JG: Growth inhibition and apoptosis induction by tanshinone I in human colon cancer Colo 205 cells. *Int J Mol Med* 22: 613-618, 2008.
17. Desagher S and Martinou JC: Mitochondria as the central control point of apoptosis. *Trends Cell Biol* 10: 369-377, 2000.
18. Ledgerwood EC and Morison IM: Targeting the apoptosome for cancer therapy. *Clin Cancer Res* 15: 420-424, 2009.
19. Gross A, McDonnell JM and Korsmeyer SJ: Bcl-2 family members and the mitochondria in apoptosis. *Genes Dev* 13: 1899-1911, 1999.
20. He JT, Zhou QH, Yuan SL, Wang YP, Chen XO and Qin JJ: Apoptosis-inducing effect of Tanshinone and its molecular mechanism on human lung cancer cells. *Chin J Lung Cancer* 5: 257-259, 2002.
21. Oyadomari S and Mori M: Roles of CHOP/GADD153 in endoplasmic reticulum stress. *Cell Death Differ* 11: 381-389, 2004.
22. Kim SH, Hwang CL, Park WY, Lee JH and Song YS: GADD153 mediates celecoxib-induced apoptosis in cervical cancer cells. *Carcinogenesis* 27: 1961-1969, 2006.
23. Kim R, Emi M, Tanabe K and Murakami S: Role of the unfolded protein response in cell death. *Apoptosis* 11: 5-13, 2006.
24. Rasheva VI and Domingos PM: Cellular responses to endoplasmic reticulum stress and apoptosis. *Apoptosis* 14: 996-1007, 2009.