

Diffusion weighted imaging for the differential diagnosis of benign vs. malignant ovarian neoplasms

XIANG-FU MENG^{1*}, SHI-CAI ZHU^{1*}, SHAO-JUAN SUN¹, JI-CAI GUO² and XUE WANG³

Departments of ¹Radiology, ²Respiratory Medicine and ³Ultrasound,
Linyi Traditional Chinese Medicine Hospital, Linyi, Shandong 276003, P.R. China

Received February 26, 2015; Accepted January 5, 2016

DOI: 10.3892/ol.2016.4445

Abstract. In order to assess the diagnostic accuracy of diffusion weighted imaging (DWI) in differentiating between benign and malignant ovarian neoplasms, a systemic meta-analysis was conducted. Relevant studies were retrieved from scientific literature databases, including the PubMed, Wiley, EBSCO, Ovid, Web of Science, Wanfang, China National Knowledge Infrastructure and VIP databases. Following a multi-step screening and study selection process, the relevant data was extracted for use in the present study. Statistical analyses were performed using Meta-disc software version 1.4 and STATA statistical software version 12.0. A total of 285 articles were retrieved from the database searches. Following a careful screening process, 10 case-control studies were selected for the present meta-analysis. The 10 studies investigated the efficacy of DWI in diagnosing ovarian neoplasms, and included a combined total of 1,159 subjects, of which 559 patients had malignant lesions and 600 had benign lesions. The results showed that the pooled sensitivity, pooled specificity, pooled positive likelihood ratio, pooled negative likelihood ratio, pooled diagnostic odds ratio (DOR) and area under the curve of the summary receiver operating characteristics curve of DWI for differentiating between benign and malignant ovarian neoplasms were 0.93, 0.89, 7.58, 0.10, 85.33 and 0.95, respectively. A subgroup analysis based on ethnicity revealed no significant difference between Asians and Caucasians. Another subgroup analysis by magnetic resonance imaging (MRI) type showed that the DORs for GE Healthcare Life Sciences and Siemens AG machines were 100.76 [95% confidence interval (CI), 65.28-155.53] and 30.85 (95% CI, 10.40-91.53), respectively; this indicates that the diagnostic efficiency of the GE Healthcare Life Sciences MRI is superior compared with

the Siemens AG MRI. The DWI demonstrated an excellent diagnostic performance in discriminating between benign and malignant ovarian neoplasms, and predicted the surgical outcome in ovarian neoplasms.

Introduction

Ovarian neoplasms are the sixth most common type of cancer in terms of affected population and the seventh leading cause of cancer-associated mortality among all gynecological malignancies worldwide (1,2). Advanced ovarian neoplasms may exhibit extensive spread to the surrounding abdominal organs, including the stomach, intestine, colon, liver, lungs and pancreas, and may also promote ascites within the peritoneal cavity (3). Due to the rapid proliferation and spread of ovarian cancer within the abdominal cavity, affected patients usually undergo primary debulking surgery prior to chemotherapy with carboplatin and taxol, which may result in a disease-free survival time of 16-22 months; however, the 5-year survival rate is only 27% (4). Ovarian neoplasms have diverse morphological features and varied genetic and epigenetic alterations; consequently, distinguishing between benign lesions and malignant ovarian cancers is challenging (5). Certain evidence indicates that combination therapies are incapable of improving the clinical outcome of patients with advanced ovarian neoplasms; therefore, the early detection of ovarian malignancy is critical for patient survival (6).

Diagnostic imaging is employed to determine the suitability of a patient for surgery by assessing the primary tumor, disease volume and extent (7,8). Diffusion weighted imaging (DWI) is a non-invasive technique based on molecular water diffusion properties, and may be used to characterize tissue architecture based on microstructure, cellular density, microcirculation and cell organization (9-11). DWI has been demonstrated to be more accurate compared with computed tomography (CT) for characterizing the sites of implants of ovarian cancers, including metastases to the liver, lungs, kidneys, uterus and pancreas (12,13). In addition, the combination of DWI with gadolinium-enhanced magnetic resonance imaging (MRI), which is used for whole-body imaging of advanced ovarian cancers, was shown to increase the accuracy of determining the operability of the tumor and reduce the rate of false diagnosis, particularly for lesions that were obscured by impeded diffusion in the spleen (9).

Correspondence to: Professor Xue Wang, Department of Ultrasound, Linyi Traditional Chinese Medicine Hospital, 211 Jie Fang Road, Linyi, Shandong 276003, P.R. China
E-mail: wangxue0205@126.com

*Contributed equally

Key words: diffusion weighted imaging, ovarian neoplasms, meta-analysis, gynecological malignancy, apparent diffusion coefficients, magnetic resonance imaging, case-control studies

Quantitative analysis of DWI is conducted by employing the apparent diffusion coefficients (ADCs) (14). In general, high ADC values indicate that water molecules may move freely, which suggests a low cellularity and better organization of the tissue structure, a typical feature of normal tissues. By contrast, low ADC values indicate the impeded mobility of water molecules, which suggests high cellularity, a characteristic feature of malignant lesions. Benign lesions, including simple cysts and hemangiomas, also have high ADC values due to the low cell density of the lesion (15-17). Although multiple studies have described the advantages of DWI over other techniques for the differential diagnosis of benign and malignant tumors, other reports have indicated that DWI has a comparable sensitivity and lower specificity compared with CT (18,19). In order to address this issue, a meta-analysis based on high quality published studies was conducted in the present study in order to evaluate the diagnostic value of DWI in discriminating between benign and malignant ovarian neoplasms.

Materials and methods

Data sources and keywords. In order to identify relevant papers that assessed the diagnostic value of DWI in ovarian neoplasms, a comprehensive search of the Ovid (available from, ovidsp.ovid.com/ovidweb.cgi), PubMed (available from, www.ncbi.nlm.nih.gov/pubmed), EBSCO (available from, search.ebscohost.com/), Wiley (available from, onlinelibrary.wiley.com/), Web of Science (available from, www.webofknowledge.com/), China National Knowledge Infrastructure (available from, www.cnki.net), Wanfang (available from, www.wanfangdata.com.cn/) and VIP (available from, www.vip.com) databases (last updated search on September 30, 2014) was conducted, applying selected common keywords associated with ovarian neoplasms and DWI. The search terms for a sensitive search strategy were as follows: 'Diffusion weighted imaging', 'diffusion magnetic resonance imaging', 'diffusion MRI', 'diffusion weighted MRI', 'diffusion weighted imaging' or 'diffusion' in combination with 'ovarian neoplasms', 'ovary neoplasms', 'ovary cancer', 'ovarian cancer', 'cancer of ovary', 'ovarian tumor', 'malignant tumor of ovary', 'ovarian carcinoma', 'ovarian epithelial carcinoma' or 'OCE'. A manual search was also employed to identify other relevant studies.

Selection and exclusion criteria. The published studies that were selected for the current meta-analysis fulfilled the following inclusion criteria: i) Studies are prospective or retrospective and evaluate the accuracy of DWI in discriminating between benign and malignant ovarian neoplasms; ii) studies must use histopathological results as the diagnostic gold standard; iii) studies must provide available data to calculate the sensitivity, specificity, negative likelihood ratio and positive likelihood ratio, and provide the number of ovarian neoplasm lesions; iv) studies must provide information on the type of MRI device used; and v) studies must be Chinese or English articles. The exclusion criteria were as follows: i) Studies do not conform to the inclusion criteria; ii) articles are abstracts, reviews, case reports, letters, meta-analyses or proceedings; iii) studies do not contain the number of benign or malignant neoplasms; iv) the data lacks integrity; v) publications are

	Han (2005)	Low (2009)	Yuan (2011)	Li (2012)	Zhang (2012)	Cui (2012)	Zhang (2012)	Espada (2013)	Kierans (2013)	Michelsen (2014)	
Representative spectrum?	+	+	+	+	+	+	+	+	+	+	+
Acceptable reference standard?	+	+	+	+	+	+	+	+	+	+	+
Acceptable delay between tests?	?	?	?	?	?	?	?	?	?	?	?
Partial verification avoided?	+	+	+	+	+	+	+	+	+	+	+
Differential verification avoided?	+	+	+	+	+	+	+	+	+	+	+
Incorporation avoided?	+	+	+	+	+	+	+	+	+	+	+
Reference standard results blinded?	?	+	?	?	?	?	?	+	?	+	+
Index test results blinded?	+	+	+	+	+	+	+	+	+	+	+
Relevant clinical information?	+	+	+	+	+	+	+	+	+	+	+
Uninterpretable results reported?	+	+	?	?	+	?	?	+	+	+	+
Withdrawals explained?	-	-	-	-	-	-	-	-	-	-	-

Figure 1. Quality assessment of diagnostic accuracy studies score of the studies included in the present meta-analysis. +, yes; -, no; ?, unclear.




Figure 2. Bivariate boxplot of diffusion weighted imaging for discriminating benign vs. malignant ovarian neoplasms in the present meta-analysis. LOGIT SENS, logit sensitivity; LOGIT SPEC, logit specificity.

duplicates or studies contain overlapping data; and vi) studies are conducted in a non-human population. In addition, only the largest or most recently published study was included in instances where the author had published several studies based on one clinical dataset.

Data extraction and quality assessment. A standard data extraction form was used, and descriptive information on a range of factors was independently collected, including first author, year of submission, state, ethnicity, research design, number of lesions, gender, age and pathological types; sensitivity, specificity, accuracy, true positive, false positive, true-negative and false-negative values; and MRI apparatus types (manufacturer) and ADC threshold. Discrepancies during data extraction were resolved by reexamination of all items and discussion between reviewers. In order to determine whether the studies involved were of high quality, a quality assessment of diagnostic accuracy studies (QUADAS) analysis was used to evaluate the studies by two independent

Table I. Baseline characteristics of patients.

Study author	Year	Ethnicity	Number of cases			Age, years			QUADAS score	Refs.
			Total	Benign	Malignant	Benign	Malignant	All patients		
Michielsen <i>et al</i>	2014	Caucasian	475	267	208			61.9±31.5	9	(9)
Kierans <i>et al</i>	2013	Caucasian	37	28	9			54.0±14.0	8	(25)
Espada <i>et al</i>	2013	Caucasian	34	26	8			53.1±11.9	9	(12)
Zhang <i>et al</i>	2012	Asian	75	45	30	48.2±30.0	50.2±28.5		7	(28)
Cui <i>et al</i>	2012	Asian	84	34	50	50.8±13.1	54.5±11.5		7	(27)
Zhang <i>et al</i>	2012	Asian	202	74	128			56.5±15.3	8	(26)
Li <i>et al</i>	2012	Asian	131	46	85	46.2±15.5	59.9±10.6		7	(29)
Yuan <i>et al</i>	2011	Asian	45	22	23			47.8±26.5	7	(30)
Low <i>et al</i>	2009	Caucasian	34	7	27			58.5	9	(11)
Han <i>et al</i>	2005	Asian	42	10	32			-	8	(31)

Age data are expressed as the mean ± standard deviation. Missing data are marked by '-'. QUADAS, quality assessment of diagnostic accuracy studies; Ref., reference number.




Figure 3. (A) Sensitivity and (B) specificity analyses of the included studies on the diagnostic capabilities of diffusion weighted imaging for discriminating between benign and malignant ovarian neoplasms. CI, confidence interval.

parties (20). The 11 QUADAS items were: The spectrum of patients was representative of the patients that will receive the test in practice (QUADAS01); the selection criteria were clearly described (QUADAS02); the time period between the reference standard and index test was short enough to be reasonably sure that the target condition did not change between the two tests (QUADAS03); the whole sample or a

random selection of the sample received verification using a reference standard of diagnosis (QUADAS04); patients received the same reference standard regardless of the index test result (QUADAS05); the reference standard was independent of the index test (QUADAS06); the execution of the index test was described in sufficient detail to permit replication of the test (QUADAS07); the execution of the reference




Figure 4. (A) The diagnostic OR forest plot of included studies on the diagnostic capabilities of diffusion weighted imaging in discriminating benign vs. malignant ovarian neoplasms. (B) The SROC curve of included studies on the diagnostic capabilities of diffusion weighted imaging in discriminating benign vs. malignant ovarian neoplasms. OR, odds ratio; CI, confidence interval; SROC, summary receiver operating characteristic; SENS, sensitivity; SPEC, specificity; AUC, area under the curve.

standard was described in sufficient detail to permit replication (QUADAS08); the same clinical data was available at the time test results were interpreted and at the time the test was applied in practice (QUADAS09); uninterpretable test results were reported (QUADAS10); and withdrawals from the study were explained (QUADAS11).

Statistical analysis. The present meta-analysis was performed using STATA software (version 12.0; Stata Corp, College Station, TX, USA) and Meta-disc software (version 1.4; Meta-DiSc, Madrid, Spain). A random effects model or a fixed effects model was applied to calculate the diagnostic odds ratio (DOR). The multiple parameters of specificity, sensitivity, positive likelihood ratio, negative likelihood ratio and summary receiver operating characteristic (SROC) curve to count the area under the curve (AUC) were used to assess the diagnostic value of DWI in discriminating between benign and malignant ovarian neoplasms. A heterogeneity test, that included tests

of threshold effect and non-threshold effect, was evaluated by applying the Spearman correlation coefficients (21) of the logarithms of sensitivity and 1-specificity. A non-threshold effect occurred with $P > 0.05$ ($P < 0.05$ was considered to indicate that a threshold effect existed). The heterogeneity test for a non-threshold effect was applied with an I^2 test (0%, no heterogeneity; 100%, maximal heterogeneity) (22) to reflect the potential for heterogeneity between studies. A random effects analysis was used for cases in which heterogeneity was observed between studies ($P < 0.05$ or $I^2 > 50\%$); otherwise a fixed effects analysis was applied.

Data combination was conducted by applying a SROC to calculate the AUC (23), with scores that ranged from 0-1 in case of heterogeneity due to the threshold effect (an AUC of 1 was considered to be the maximal diagnostic value). If the heterogeneity was not caused by threshold effect, the targets, including specificity, sensitivity, positive likelihood ratio, negative likelihood ratio and DOR, were combined to produce




Figure 5. (A) Results of the pooled likelihood ratio suggested a limited clinical value of DWI for discriminating between benign and malignant ovarian neoplasms. (B) Fagan's nomogram of the included studies on the diagnostic capabilities of diffusion weighted imaging for discriminating between benign and malignant ovarian neoplasms. LUQ, left upper quadrant; LRP, likelihood ratio positive; LRN, likelihood ratio negative; RUQ, right upper quadrant; LLQ, left lower quadrant; RLQ, right lower quadrant.

a forest plot. The studies were excluded in turn, in order to analyze the effect of single study on the overall results. Fagan's nomogram was applied to estimate the pre-test and post-test probability of three diagnostic criteria, and a bivariate boxplot was used to judge the heterogeneity between studies. The included angle between the regression line in Deeks funnel plot (24) and DOR was applied to assess publication bias.

Results

Included studies. A total of 285 articles were retrieved from the 9 databases by keyword search. Following the exclusion of duplicates (n=89), reviews, letters or meta-analyses (n=42), non-human studies (n=53), and studies not relevant to our research topics (n=56), the remaining studies (n=45) were carefully reviewed. An additional 11 studies were excluded as they were not cohort or case-control studies (n=2), not relevant to DWI (n=4), or not relevant to ovarian neoplasms (n=5). Subsequent to additional reviewing of the remaining 34 trials, the studies that did not supply enough information were eliminated (n=4), and 10 high quality studies were included in the final meta-analysis. These 10 case-control studies, published between 2005 and 2014, provided the required information on the diagnostic value of DWI and ovarian neoplasms (9,11,12,25-31). The meta-analysis included

a total of 1,159 ovarian tumor patients, comprising 559 benign tumor patients and 600 malignant tumor patients. Of the 10 studies, 6 were conducted on Asian patient populations and 4 on Caucasian patient populations. The QUADAS of included studies is shown in Fig. 1, and Table I presents the baseline characteristics of the selected study samples.

Although the majority of the studies were located in the middle region of the bivariate boxplot, two studies were located outside of the boxplot, which indicated heterogeneity between the studies (Fig. 2). The result of the Spearman correlation coefficient of the sensitivity logarithm and 1-specificity logarithm indicated that no threshold effect existed (0.345; P=0.328). The I² values of sensitivity (57.5%) and negative likelihood ratio (54.5%) were >50%, which indicated statistical heterogeneity between studies, and a random effects model was applied. However, the I² values for specificity (23.9%) and positive likelihood ratio (33.1%) were <50%, which suggested that no statistical heterogeneity existed; therefore, a fixed effects model was used.

Quantitative data synthesis. The results of the present meta-analysis demonstrated that DWI has high diagnostic capabilities in discriminating between benign and malignant ovarian neoplasms, based on the outcomes of pooled sensitivity [0.93; 95% confidence interval (CI), 0.91-0.95; Fig. 3A], pooled




Figure 6. (A) Subgroup analysis based on ethnicity of the diagnostic capability of diffusion weighted imaging in discriminating benign vs. malignant ovarian neoplasms. (B) The subgroup analysis based on magnetic resonance imaging apparatus manufacturer (GE Healthcare Life Sciences or Siemens AG) of the diagnostic capability of diffusion weighted imaging in discriminating benign vs. malignant ovarian neoplasms. OR, odds ratio; CI, confidence interval.




Figure 7. (A) Sensitivity analysis chart for the publication bias of the included studies regarding the diagnostic capabilities of diffusion weighted imaging for discriminating between benign and malignant ovarian neoplasms. (B) Deeks for the publication bias of the included studies. CI, confidence interval; ESS, effective sample size.

specificity (0.89; 95% CI, 0.86-0.91; Fig. 3B), pooled positive likelihood ratio (7.58; 95% CI, 6.00-9.56), pooled negative likelihood ratio (0.10; 95% CI, 0.06-0.16), pooled DOR (85.33; 95% CI, 57.15-127.40) (Fig. 4A) and the AUC of the SROC

curve (0.95; Fig. 4B). The results of the pooled likelihood ratio suggested a limited clinical value of DWI for discriminating between benign and malignant ovarian neoplasms (Fig. 5A). The results of Fagan's nomogram indicated that the pre-test

probability ratio (20%) x positive likelihood ratio (8.00) yielded a post-test probability ratio of 66%, while the pre-test probability ratio x negative likelihood ratio (0.08) yielded a post-test probability ratio of 2% (Fig. 5B). Subgroup analyses of ethnicity and MRI types were conducted. The results indicated that the DORs for patients of Asian and Caucasian descent were 86.37 (95% CI, 48.35-154.27) and 84.33 (95% CI, 48.44-146.81), respectively (Fig. 6A), suggesting no statistically significant differences between Asians and Caucasians. Notably, the DORs for GE Healthcare Life Sciences and Siemens AG MRIs were 100.76 (95% CI, 65.28-155.53) and 30.85 (95% CI, 10.40-91.53), respectively; thus, the diagnostic value of GE Healthcare Life Sciences machine appears to be superior to that of the Siemens AG equipment (Fig. 6B).

Sensitivity analysis and publication bias. A sensitivity analysis was applied in order to estimate the stability of the present meta-analysis. The outcomes of the sensitivity analysis revealed that none of the included studies had an influence on the pooled DOR of the DWI for discriminating between benign and malignant ovarian neoplasms (Fig. 7A). The included angle between the regression line and the DOR axis was close to 90° in Deeks, which indicated that no publication bias was present in the current meta-analysis (Fig. 7B).

Discussion

In order to evaluate the diagnostic value of DWI in discriminating between benign and malignant ovarian neoplasms, a systematic meta-analysis was conducted. The outcomes of the present study indicated that DWI has a high diagnostic value with regard to discriminating between benign and malignant ovarian neoplasms. Ovarian neoplasms are the sixth most prevalent cancer and the seventh leading cause of cancer-associated mortality among all types of gynecological malignancy worldwide (1,2). Due to the high frequency of peritoneal and distant metastases and the complicated nature of ovarian neoplasms, the patient survival rate is poor (27,32). Diagnostic imaging is used to determine the suitability of the patient for surgery by assessing the primary tumor or by describing the disease volume and extent (7,8). DWI, as a functional MRI technique, provides non-invasive and high-resolution information on tissue characteristics based on molecular water diffusion properties, and the imaging output is useful to determine tissue microstructure, density, microcirculation and cellular organization (9,11,33). Several studies describe a close association between cell density and ADC values (16,33-35). High ADC values are present in benign lesions, including primary neoplasms that contain low cell densities, while low ADC values are typically observed in malignant neoplasms, due to high cell densities and altered membrane permeability (15,16). Based on the results of the current meta-analysis, a major conclusion is that DWI demonstrates an excellent diagnostic performance for discriminating between benign and malignant ovarian neoplasms.

Subgroup analyses based on ethnicity and MRI devices were performed. The result of the subgroup analysis on ethnicity indicated no statistically significant differences between Asians and Caucasians. However, the subgroup analysis on the MRI machine type demonstrated that a better

diagnostic performance of DWI was achieved with GE Healthcare Life Sciences equipment compared with Siemens AG machines.

The present study was subject to certain limitations. First, the number of studies included was small, and several relevant unpublished studies and abstracts were not included, which may have influenced the conclusion of the current study. A second limitation of the present study was that the sample size was relatively small, leading to a lower confidence in assessing the value of DWI for discriminating between benign and malignant ovarian neoplasms. Third, only studies published in English and Chinese were included, with studies published in other languages excluded, which potentially eliminated important studies that may have influenced the conclusion of the present meta-analysis. Additional studies are required that involve a larger sample size and diverse ethnicities for a more comprehensive evaluation of the diagnostic performance of DWI in ovarian neoplasms.

In conclusion, DWI demonstrates an excellent diagnostic value for discriminating between benign and malignant ovarian neoplasms, and may provide a reliable predictive tool for the surgical outcome in ovarian neoplasms.

References

- Zeng ST, Guo L, Liu SK, Wang DH, Xi J, Huang P, Liu DT, Gao JF, Feng J and Zhang L: Egg consumption is associated with increased risk of ovarian cancer: Evidence from a meta-analysis of observational studies. *Clin Nutr* 34: 635-641, 2015.
- Ahmed N, Abubaker K and Findlay JK: Ovarian cancer stem cells: Molecular concepts and relevance as therapeutic targets. *Mol Aspects Med* 39: 110-125, 2014.
- Lengyel E: Ovarian cancer development and metastasis. *Am J Pathol* 177: 1053-1064, 2010.
- Kipps E, Tan DS and Kaye SB: Meeting the challenge of ascites in ovarian cancer: New avenues for therapy and research. *Nat Rev Cancer* 13: 273-282, 2013.
- Conic I, Dimov I, Tasic-Dimov D, Djordjevic B and Stefanovic V: Ovarian epithelial cancer stem cells. *ScientificWorldJournal* 11: 1243-1269, 2011.
- Bristow RE, Puri I and Chi DS: Cytoreductive surgery for recurrent ovarian cancer: A meta-analysis. *Gynecol Oncol* 112: 265-274, 2009.
- Forstner R, Sala E, Kinkel K and Spencer JA; European Society of Urogenital Radiology: ESUR guidelines: Ovarian cancer staging and follow-up. *Eur Radiol* 20: 2773-2780, 2010.
- Kumar Dhingra V, Kand P and Basu S: Impact of FDG-PET and -PET/CT imaging in the clinical decision-making of ovarian carcinoma: An evidence-based approach. *Women's Health (Lond Engl)* 8: 191-203, 2012.
- Michielsen K, Vergote I, Op de Beeck K, Amant F, Leunen K, Moerman P, Deroose C, Souverijns G, Dymarkowski S, De Keyzer F and Vandecaveye V: Whole-body MRI with diffusion-weighted sequence for staging of patients with suspected ovarian cancer: A clinical feasibility study in comparison to CT and FDG-PET/CT. *Eur Radiol* 24: 889-901, 2014.
- Low RN and Gurney J: Diffusion-weighted MRI (DWI) in the oncology patient: Value of breathhold DWI compared to unenhanced and gadolinium-enhanced MRI. *J Magn Reson Imaging* 25: 848-858, 2007.
- Low RN, Sebrechts CP, Barone RM and Muller W: Diffusion-weighted MRI of peritoneal tumors: Comparison with conventional MRI and surgical and histopathologic findings - a feasibility study. *AJR Am J Roentgenol* 193: 461-470, 2009.
- Espada M, Garcia-Flores JR, Jimenez M, Alvarez-Moreno E, De Haro M, Gonzalez-Cortijo L, Hernandez-Cortes G, Martinez-Vega V and Sainz De La Cuesta R: Diffusion-weighted magnetic resonance imaging evaluation of intra-abdominal sites of implants to predict likelihood of suboptimal cytoreductive surgery in patients with ovarian carcinoma. *Eur Radiol* 23: 2636-2642, 2013.

13. Low RN and Barone RM: Combined diffusion-weighted and gadolinium-enhanced MRI can accurately predict the peritoneal cancer index preoperatively in patients being considered for cytoreductive surgical procedures. *Ann Surg Oncol* 19: 1394-1401, 2012.
14. White NS, McDonald CR, Farid N, Kuperman JM, Kesari S and Dale AM: Improved conspicuity and delineation of high-grade primary and metastatic brain tumors using 'restriction spectrum imaging': Quantitative comparison with high B-value DWI and ADC. *AJNR Am J Neuroradiol* 34: 958-964, 2013.
15. Testa ML, Chojniak R, Sene LS, Damascena AS, Guimarães MD, Szklaruk J and Marchiori E: Is DWI/ADC a useful tool in the characterization of focal hepatic lesions suspected of malignancy? *PLoS One* 9: e101944, 2014.
16. Li SP and Padhani AR: Tumor response assessments with diffusion and perfusion MRI. *J Magn Reson Imaging* 35: 745-763, 2012.
17. Galea N, Cantisani V and Taouli B: Liver lesion detection and characterization: Role of diffusion-weighted imaging. *J Magn Reson Imaging* 37: 1260-1276, 2013.
18. Soussan M, Des Guetz G, Barrau V, Aflalo-Hazan V, Pop G, Mehanna Z, Rust E, Aparicio T, Douard R, Benamouzig R, *et al*: Comparison of FDG-PET/CT and MR with diffusion-weighted imaging for assessing peritoneal carcinomatosis from gastrointestinal malignancy. *Eur Radiol* 22: 1479-1487, 2012.
19. Satoh Y, Ichikawa T, Motosugi U, Kimura K, Sou H, Sano K and Araki T: Diagnosis of peritoneal dissemination: Comparison of 18F-FDG PET/CT, diffusion-weighted MRI, and contrast-enhanced MDCT. *AJR Am J Roentgenol* 196: 447-453, 2011.
20. Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM and Kleijnen J: The development of QUADAS: A tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. *BMC Med Res Methodol* 3: 25, 2003.
21. Pfeiffer KA, McIver KL, Dowda M, Almeida MJ and Pate RR: Validation and calibration of the Actical accelerometer in preschool children. *Med Sci Sports Exerc* 38: 152-157, 2006.
22. Higgins JP and Thompson SG: Quantifying heterogeneity in a meta-analysis. *Stat Med* 21: 1539-1558, 2002.
23. Walter SD: Properties of the summary receiver operating characteristic (SROC) curve for diagnostic test data. *Stat Med* 21: 1237-1256, 2002.
24. Deeks JJ, Macaskill P and Irwig L: The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *J Clin Epidemiol* 58: 882-893, 2005.
25. Kierans AS, Bennett GL, Mussi TC, Babb JS, Rusinek H, Melamed J and Rosenkrantz AB: Characterization of malignancy of adnexal lesions using ADC entropy: Comparison with mean ADC and qualitative DWI assessment. *J Magn Reson Imaging* 37: 164-171, 2013.
26. Zhang P, Cui Y, Li W, Ren G, Chu C and Wu X: Diagnostic accuracy of diffusion-weighted imaging with conventional MR imaging for differentiating complex solid and cystic ovarian tumors at 1.5T. *World J Surg Oncol* 10: 237, 2012.
27. Cui YF, Li WH, Zhu MJ, Wang DB, Zhang P, Zhang ZY and Chu CT: Clinical application and research of diffusion weighted MR imaging in complex ovarian tumors. *Zhongguo Linchuang Yixue Yingxiang Zazhi* 12: 856-859, 2012 (In Chinese).
28. Zhang W, Zou S, Shen DH and Tong YX: The value of MR diffusion-weighted imaging combined with serum CA125 in the qualitative diagnosis of ovarian occupying lesion. *Yixue Yingxiangxue Zazhi* 8: 1348-1353, 2012 (In Chinese).
29. Li WH, Chu C, Cui Y, Zhang P and Zhu M: Diffusion-weighted MRI: A useful technique to discriminate benign versus malignant ovarian surface epithelial tumors with solid and cystic components. *Abdom Imaging* 37: 897-903, 2012.
30. Yuan XC, Wang X, Yao G, Zheng L and Hu Y: The value of 3.0T MRI for the diagnosis of the ovarian tumor. *Shiyong Fangshexue Zazhi* 27: 1695-1698, 2011 (In Chinese).
31. Han DM, Yang XP, Wang HP and Li YX: Comparative study of CT and MRI in the diagnosis of ovarian neoplasms. *Zhongguo Linchuang Yixue Yingxiang Zazhi* 16: 441-443, 2005 (In Chinese).
32. Ferlay J, Parkin DM and Steliarova-Foucher E: Estimates of cancer incidence and mortality in Europe in 2008. *Eur J Cancer* 46: 765-781, 2010.
33. Decker G, Mürtz P, Gieseke J, Träber F, Block W, Sprinkart AM, Leitzen C, Buchstab T, Lütter C, Schüller H, *et al*: Intensity-modulated radiotherapy of the prostate: dynamic ADC monitoring by DWI at 3.0 T. *Radiother Oncol* 113: 115-120, 2014.
34. Ginat DT, Mangla R, Yeane G, Johnson M and Ekholm S: Diffusion-weighted imaging for differentiating benign from malignant skull lesions and correlation with cell density. *AJR Am J Roentgenol* 198: W597-601, 2012.
35. Rechichi G, Galimberti S, Signorelli M, Franzesi CT, Perego P, Valsecchi MG and Sironi S: Endometrial cancer: Correlation of apparent diffusion coefficient with tumor grade, depth of myometrial invasion, and presence of lymph node metastases. *AJR Am J Roentgenol* 197: 256-262, 2011.