

Identification of key genes associated with the effect of osmotic stimuli on intervertebral discs using microarray analysis

GUANGXIAO NI^{1*}, GUOBIN LIU^{2*} and KUNLUN YU³

¹Department of Rehabilitation, The Second Hospital of Hebei Medical University;

²Department of Orthopaedics, The First Hospital of Hebei Medical University, Shijiazhuang, Hebei 050000;

³Department of Orthopaedics, The Third Hospital of Hebei Medical University, Shijiazhuang, Hebei 050051, P.R. China

Received June 7, 2016; Accepted March 14, 2017

DOI: 10.3892/ol.2017.6657

Abstract. The present study aimed to explore the effect of osmotic stimuli on intervertebral discs (IVDs) using microarray analysis. Gene expression dataset GSE1648 was downloaded from the Gene Expression Omnibus database. There were 11 IVD cell samples in this dataset, which included 4 hyperosmotic stimuli samples, 3 hypoosmotic stimuli samples and 4 isosmotic stimuli samples. The differentially expressed genes (DEGs) in hyperosmotic or hypoosmotic IVD cells (designated DEGs-hyper or DEGs-hypo) were identified, compared with isosmotic cells, using the limma package of R software. The Database for Annotation, Visualization and Integrated Discovery was used to perform a Gene Ontology (GO) term enrichment analysis for the DEG sets. Protein-protein interaction (PPI) network and microRNA (miRNA) gene-regulatory network data for the DEG sets were obtained using the Human Protein Reference Database (HPRD) and the TargetScan database, respectively, and these networks were constructed and visualized using Cytoscape software. There was a total of 43 DEGs in DEGs-hyper and 9 in DEGs-hypo. Analysis of DEGs-hyper revealed that 41 GO terms were significantly enriched. In total, 376 pairs and 382 nodes were involved in the PPI network, and 1,314 miRNA-gene pairs and 422 nodes were contained in the miRNA-gene-regulated network. The results of the present study indicated that potential target genes (including *NCOA3*, *SOS1*, *XPO1*, *ZBTB18*, *EFNB2* and *SOBP*) may be involved in the effect of osmotic stimuli on IVD, and the biological processes of apoptosis and cell death may be associated with the effect of high osmolality on IVD disease. The potential targets identified in the present study are more reliable than those identified by previous studies.

Introduction

Intervertebral disc (IVD) disease is a common surgical disease, and IVD degeneration is hypothesized to be the first step in degenerative spinal changes (1). IVD degeneration is typically associated with disc herniation and back pain, which has a marked effect on the sufferer's life and causes severe chronic pain (2). Furthermore, lower back pain may limit the activity of individuals <45 years, which has a marked socio-economic impact (3). The etiology of lower back pain is unclear, but in 40% of cases it is associated with IVD degeneration (4). It is reported that ~40% of individuals <30 years of age, and >90% of individuals >55 years, exhibit moderate-to-severe levels of IVD degeneration (5,6). It is estimated that the costs associated with lumbar disc and lower back disorders exceed \$100 billion/year in the USA (7). Traditionally, inflammation has mostly been observed as detrimental and is associated with disease progression. Although the etiology remains unknown, the osmolality has been identified to be associated with IVD disease (8-10). Katz (7) reported that the osmotic environment exerted an appreciable effect on gene expression and also affected responses to mechanical stimuli. Mavrogonatou and Kletsas (11) indicated that high osmolality was able to activate the G₁ and G₂ cell cycle checkpoints and affect the DNA integrity of nucleus pulposus IVD cells, triggering an enhanced DNA repair response. Although changes in the extracellular osmolality markedly influenced the behavior of IVD cells, their response to this condition has not yet been fully elucidated. In the present study, the associated biological processes and potential biomarkers in the response of IVD to osmotic stimuli were investigated using gene expression analysis.

Materials and methods

Gene expression profile. The gene expression dataset GSE1648 (8) was downloaded from the Gene Expression Omnibus (<http://www.ncbi.nlm.nih.gov/geo/>) database. There was a total of 11 IVD cell samples in GSE1648, including 4 hyperosmotic stimuli samples, 3 hypoosmotic stimuli samples and 4 isosmotic stimuli samples.

Data pre-processing and identification of differentially expressed genes (DEGs). For the expression profile, the

Correspondence to: Dr Kunlun Yu, Department of Orthopaedics, The Third Hospital of Hebei Medical University, 139 Zi-Qiang Road, Shijiazhuang, Hebei 050051, P.R. China
E-mail: yukunlun@126.com

*Contributed equally

Key words: hyper-osmotic stimuli, intervertebral disc degeneration, bioinformatics




Figure 1. Cluster graph of differentially expressed genes in intervertebral disc cells of hyperosmotic stimuli samples compared with the isosmotic stimuli.

original data were converted into a recognizable format using R software version 3.1.1 (from bioconductor.org/packages/release/bioc/html/biomaRt.html), and the Affy package (12) version 1.48.0 (from bioconductor.org/packages/release/bioc/html/affy.html) was used for the background correction and normalization. The DEGs in hyperosmotic (designated DEGs-hyper) or hypoosmotic (DEGs-hypo) IVD cells, compared with isosmotic cells, were identified using the limma package (13) version 3.18.13 (from bioconductor.org/packages/release/bioc/html/limma.html) based on gene expression differences of $P<0.05$ and $|log_2(\text{fold-change})|>0.05$.

Gene Ontology (GO) term enrichment analysis. Functional annotation of DEGs is a necessary and critical step in the analysis of microarray data. The Database for Annotation, Visualization and Integrated Discovery (DAVID, <http://david.abcc.ncifcrf.gov/>) (14) is a tool providing a comprehensive set of functional annotation. Enriched GO terms from DEGs-hyper were identified using DAVID with a threshold of $P<0.05$.

Construction of the protein-protein interaction (PPI) network. The Human Protein Reference Database (HPRD) was used as

a centralized platform to visually depict and integrate information pertaining to domain architecture, post-translational modifications, interaction networks and disease association for each protein in the human proteome (15). PPI network data for DEGs-hyper was obtained from HPRD and visualized using Cytoscape (16) software.

Establishment and analysis of the microRNA (miRNA)-regulated-gene network. The software TargetScan (17) was used to predict biological targets of miRNAs by searching for the presence of conserved 8-mer and 7-mer sites that matched the seed region of each miRNA. TargetScan screened for miRNAs associated with the regulation of DEGs-hyper, and miRNA-gene connections were obtained. An miRNA-regulated-gene network was established and visualized using Cytoscape software, and every node was analyzed according to the number of connections with other nodes (degree). miRNAs for which the degree was ≥ 1 were removed from the analysis.

Results

DEGs. A total of 43 DEGs (34 up- and 9 downregulated) were identified in DEGs-hyper, and the cluster graph is presented in

Table I. Top 10 most significantly enriched GO terms of differentially expressed genes in intervertebral disc cells of hyperosmotic stimuli samples compared with the isosmotic stimuli.

Category	GO ID	GO name	Number of genes	P-value
BP	GO:0042981	Regulation of apoptosis	9	3.48x10 ⁻⁴
BP	GO:0043067	Regulation of programmed cell death	9	3.73x10 ⁻⁴
BP	GO:0010941	Regulation of cell death	9	3.82x10 ⁻⁴
BP	GO:0006913	Nucleocytoplasmic transport	5	4.20x10 ⁻⁴
BP	GO:0051169	Nuclear transport	5	4.41x10 ⁻⁴
BP	GO:0012501	Programmed cell death	7	0.00232
BP	GO:0008219	Cell death	7	0.00519
BP	GO:0030183	B cell differentiation	3	0.00537
BP	GO:0016265	Death	7	0.00537
BP	GO:0051168	Nuclear export	3	0.00828

GO, Gene Ontology; ID, identifier; BP, biological process.




Figure 2. Protein-protein interaction network of differentially expressed genes in intervertebral disc cells of hyperosmotic stimuli samples compared with the isosmotic stimuli.

Table II. Top 20 nodes with a higher degree in the protein-protein interaction network.

Gene	Degree
NCOA3	49
SOS1	45
XPO1	45
RRAS2	27
TP53BP2	23
BCL6	22
NXF1	22
EFNB2	21
SGK1	16
ARNTL	13
LSM6	10
PTPRE	10
VCAM1	10
BDNF	9
GPSM2	7
FNBP4	6
GLUL	5
MALT1	4
MAPK1	4
PPM1D	4

Degree indicates the number of connections of each node with other nodes in the network.

Table III. Top 20 nodes with higher degree in the microRNA-gene-regulated network.

Gene	Degree
ZBTB18	81
EFNB2	73
SOBP	72
NXF1	68
BDNF	66
PTBP3	64
PPM1D	59
BAMBI	58
NCOA3	55
KLHL24	47
SGK1	45
GAREM	42
NUAK1	42
SOS1	42
BCL6	38
XPO1	37
FAM110B	34
TNFAIP8	34
KLHL9	29

Degree indicates the number of connections of each node with other nodes in the network.

Fig. 1. However, only 9 DEGs were obtained in DEGs-hypo, which were not studied further.

GO terms of DEGs-hyper. In total, 41 GO terms were enriched in DEGs-hyper, and the 10 most significantly enriched GO terms (e.g., regulation of apoptosis, regulation of programmed cell death and regulation of cell death) are presented in Table I.

PPI network of DEGs-hyper. The PPI network was constructed and is presented in Fig. 2, and included 376 pairs and 382 nodes. The top 20 nodes (e.g., *NCOA3*, *SOS1* and *XPO1*), as defined by highest degree, are presented in Table II.

miRNA-regulated-gene network of DEGs-hyper. In total, 1,314 miRNA-regulated-gene connections were identified with TargetScan. A miRNA-gene-regulated network was established, including 1,314 connections and 422 nodes (Fig. 3). The top 20 nodes (e.g., *ZBTB18*, *EFNB2* and *SOBP*) in the regulated network, as determined by degree, are presented in Table III.

Discussion

The normal upper limit of water in the human IVD is ~500 mOsm/kg, which is higher than that routinely encountered in the majority of other parts of the body (~300 mOsm/kg) (18). It was previously reported that ~18% of the fluid of IVD was lost and re-imbibed during a diurnal cycle, with consequent changes in osmolality (19). Changes in osmolality are an important component of the physicochemical environment of IVD, as variations in disc loading leads to alterations of disc hydration. A study of IVD cells in a three-dimensional alginate culture system confirmed that the biological response to altered osmolarity is mediated, in part, by changes at the transcriptional level (20). In the present study, 43 DEGs were identified in hyperosmotic cells, whereas 9 DEGs were identified in hypoosmotic cells, compared with isosmotic cells. These results suggest that IVD is more sensitive to hyperosmotic stimuli than to hypoosmotic stimuli, and that the effects of hyperosmotic stimuli may be far greater at the transcriptional level. Hyperosmotic stimuli were previously demonstrated to elicit calcium transience in IVD cells that were modulated by the stability of the actin cytoskeleton (21,22). In the present study, DEGs-hyper were the basis of further research, whereas DEGs-hypo were not further studied.

The enriched GO terms for DEGs-hyper were predominantly associated with the biological processes of apoptosis and cell death, and the regulation of these processes. Three subfamilies were identified: Extracellular-signal-regulated kinase (ERK1/2), p38 mitogen-activated protein kinase (p38 MAPK, p38) and c-Jun N-terminal kinase (JNK1/2). Nucleus pulposus cell apoptosis is one of the changes associated with IVD degeneration (23). p38 and JNK1/2 are serine/threonine protein kinases activated by various stress stimuli, including osmotic shock, toxic compounds and pro-inflammatory cytokines (24). ERK1/2 was previously demonstrated to be activated by high osmolality in nucleus pulposus cells *in vitro* (25). Dong *et al* (26) reported that high osmolality activated p38 MAPK, JNK1/2 and ERK1/2 in rabbit nucleus




Figure 3. microRNA-gene-regulated network of differentially expressed genes in intervertebral disc cells of hyperosmotic stimuli samples compared with the isosmotic stimuli.

pulposus cells. The activated p38 MAPK and JNK1/2 induced cell apoptosis; by contrast, the activation of ERK1/2 promoted cell survival. A recent study (27) indicated that the effects of osmolality on nucleus pulposus cell apoptosis depended on the osmolality level (hypo-, iso- or hyper-) and osmolality mode (constant or cyclic). Furthermore, inhibition of the ERK1/2 pathway promoted nucleus pulposus cell apoptosis in this process (27). Therefore, it was suspected that the biological processes of apoptosis and cell death may be associated with the effect of hyperosmolality on IVD diseases.

The PPI network was analyzed, and nuclear receptor coactivator 3 (NCOA3), SOS Ras/Rac guanine nucleotide exchange factor 1 (SOS1) and exportin 1 (XPO1) were the top three nodes as defined by degree. NCOA3 was identified to be associated with the pathophysiology of osteoarthritis (OA) (28,29). A recent study concluded that NCOA3 was subject to a *cis*-acting expression quantitative trait locus in articular cartilage, which was associated with the OA association signal and associated with the OA-associated allele of the functional single-nucleotide polymorphism rs116855380 (30). Although there is no direct evidence that NCOA3 is associated

with IVD degeneration, IVD degeneration and OA may exhibit similarities in their occurrence and development, as they are joint degeneration diseases. *SOS1* primarily encoded membrane-bound guanine nucleotide-binding proteins in humans, which function in the transduction of signals that control cell growth and differentiation (31).

It was previously reported that *SOS1* participates in the process of apoptosis (32,33). It was identified that the biological processes of apoptosis and cell death may be associated with the effect of high osmolality on IVD diseases. The mutation of *SOS1* serves an important role in the salt-tolerance and osmotic stimuli of *Arabidopsis* and tobacco (34-37). XPO1 is a specific receptor for leucine-rich nuclear export sequences (38) and was identified to mediate the nuclear export of proteins in a range of species (39,40). Ferrigno *et al* (41) indicated that XPO1 mediated the export of high osmolality glycerol response MAPK to the cytoplasm of cells adapted to hyper-osmotic stimuli. Thus, *NCOA3*, *SOS1* and *XPO1* may be genes directly affected by osmotic stimuli in IVD cells, although more trials and clinical validation are required. Similarly, *ZBTB18*, *EFNB2* and *SOBP* were the top 3 nodes in the miRNA-gene-regulated network,

and they may exhibit an intimate association with the effects of osmotic stimuli on IVD.

The dataset GSE1648 was created by Boyd *et al* (8); however, the present study adopted different methods and obtained novel results compared with the Boyd *et al* study (8). First, differential expression analysis was performed for cells in hyperosmotic and hypoosmotic conditions in the present study, whereas Boyd *et al* (8) only conducted differential expression analysis of cells in hyperosmotic conditions. Secondly, the PPI network and miRNA-gene-regulated network were constructed in the present study; however, this was not carried out in the study by Boyd *et al* (8). Finally, potential target genes (*NCOA3*, *SOS1*, *XPO1*, *ZBTB18*, *EFNB2* and *SOBP*) were identified from the PPI network and the miRNA-gene-regulated network, whereas they were not identified in the study by Boyd *et al* (8). These genes exhibited more interactions compared with those identified by Boyd *et al*, and therefore they were considered more reliable.

References

- Boos N, Weissbach S, Rohrbach H, Weiler C, Spratt KF and Nerlich AG: Classification of age-related changes in lumbar intervertebral discs: 2002 Volvo Award in basic science. *Spine (Phila Pa 1976)* 27: 2631-2644, 2002.
- Stewart WF, Ricci JA, Chee E, Morganstein D and Lipton R: Lost productive time and cost due to common pain conditions in the US workforce. *JAMA* 290: 2443-2454, 2003.
- Andersson GB: Epidemiological features of chronic low-back pain. *Lancet* 354: 581-585, 1999.
- Schwarzer AC, Aprill CN, Derby R, Fortin J, Kine G and Bogduk N: The prevalence and clinical features of internal disc disruption in patients with chronic low back pain. *Spine (Phila Pa 1976)* 20: 1878-1883, 1995.
- Stolworthy DK, Bowden AE, Roeder BL, Robinson TF, Holland JG, Christensen SL, Beatty AM, Bridgewater LC, Eggett DL, Wendel JD, *et al*: MRI evaluation of spontaneous intervertebral disc degeneration in the alpaca cervical spine. *J Orthop Res* 33: 1776-1783, 2015.
- Cheung KM, Karppinen J, Chan D, Ho DW, Song YQ, Sham P, Cheah KS, Leong JC and Luk KD: Prevalence and pattern of lumbar magnetic resonance imaging changes in a population study of one thousand forty-three individuals. *Spine (Phila Pa 1976)* 34: 934-940, 2009.
- Katz JN: Lumbar disc disorders and low-back pain: Socioeconomic factors and consequences. *J Bone Joint Surg Am* 88 (Suppl 2): S21-S24, 2006.
- Boyd LM, Richardson WJ, Chen J, Kraus VB, Tewari A and Setton LA: Osmolarity regulates gene expression in intervertebral disc cells determined by gene array and real-time quantitative RT-PCR. *Ann Biomed Eng* 33: 1071-1077, 2005.
- Setton LA and Chen J: Mechanobiology of the intervertebral disc and relevance to disc degeneration. *J Bone Joint Surg Am* 88 (Suppl 2): S52-S57, 2006.
- Ma H, Xie YZ, Zhao J and Ye B: Small molecule-enrichment analysis in response to osmotic stimuli in the intervertebral disc. *Genet Mol Res* 11: 3668-3675, 2012.
- Mavrogenatou E and Kletsas D: High osmolality activates the G1 and G2 cell cycle checkpoints and affects the DNA integrity of nucleus pulposus intervertebral disc cells triggering an enhanced DNA repair response. *DNA Repair (Amst)* 8: 930-943, 2009.
- Gautier L, Cope L, Bolstad BM and Irizarry RA: affy-analysis of Affymetrix GeneChip data at the probe level. *Bioinformatics* 20: 307-315, 2004.
- Diboun I, Wernisch L, Orengo CA and Koltzenburg M: Microarray analysis after RNA amplification can detect pronounced differences in gene expression using limma. *BMC Genomics* 7: 252, 2006.
- Sherman BT, Huang da W, Tan Q, Guo Y, Bour S, Liu D, Stephens R, Baseler MW, Lane HC and Lempicki RA: DAVID Knowledgebase: A gene-centered database integrating heterogeneous gene annotation resources to facilitate high-throughput gene functional analysis. *BMC Bioinformatics* 8: 426, 2007.
- Keshava Prasad TS, Goel R, Kandasamy K, Keerthikumar S, Kumar S, Mathivanan S, Telikicherla D, Raju R, Shaferen B, Venugopal A, *et al*: Human protein reference database-2009 update. *Nucleic Acids Res* 37 (Database issue): D767-D772, 2009.
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B and Ideker T: Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Res* 13: 2498-2504, 2003.
- Lewis BP, Shih IH, Jones-Rhoades MW, Bartel DP and Burge CB: Prediction of mammalian microRNA targets. *Cell* 115: 787-798, 2003.
- Urban JP: The role of the physicochemical environment in determining disc cell behaviour. *Biochem Soc Trans* 30: 858-864, 2002.
- McMillan DW, Garbutt G and Adams MA: Effect of sustained loading on the water content of intervertebral discs: Implications for disc metabolism. *Ann Rheum Dis* 55: 880-887, 1996.
- Chen J, Baer AE, Paik PY, Yan W and Setton LA: Matrix protein gene expression in intervertebral disc cells subjected to altered osmolarity. *Biochem Biophys Res Commun* 293: 932-938, 2002.
- Pritchard S, Erickson GR and Guilak F: Hyperosmotically induced volume change and calcium signaling in intervertebral disk cells: The role of the actin cytoskeleton. *Biophys J* 83: 2502-2510, 2002.
- Pritchard S and Guilak F: The role of F-actin in hypo-osmotically induced cell volume change and calcium signaling in anulus fibrosus cells. *Ann Biomed Eng* 32: 103-111, 2004.
- Freemont AJ: The cellular pathobiology of the degenerate intervertebral disc and discogenic back pain. *Rheumatology (Oxford)* 48: 5-10, 2009.
- Johnson NL, Gardner AM, Diener KM, Lange-Carter CA, Gleavy J, Jarpe MB, Minden A, Karin M, Zon LI and Johnson GL: Signal transduction pathways regulated by mitogen-activated/extracellular response kinase kinase kinase induce cell death. *J Biol Chem* 271: 3229-3237, 1996.
- Hiyama A, Gogate SS, Gajghate S, Mochida J, Shapiro IM and Risbud MV: BMP-2 and TGF-beta stimulate expression of beta1,3-glucuronosyl transferase 1 (GlcAT-1) in nucleus pulposus cells through AP1, TonEBP and Sp1: Role of MAPKs. *J Bone Miner Res* 25: 1179-1190, 2010.
- Dong ZH, Wang DC, Liu TT, Li FH, Liu RL, Wei JW and Zhou CL: The roles of MAPKs in rabbit nucleus pulposus cell apoptosis induced by high osmolality. *Eur Rev Med Pharmacol Sci* 18: 2835-2845, 2014.
- Li P, Gan Y, Wang H, Xu Y, Li S, Song L, Zhang C, Ou Y, Wang L and Zhou Q: Role of the ERK1/2 pathway in osmolarity effects on nucleus pulposus cell apoptosis in a disc perfusion culture. *J Orthop Res* 35: 86-92, 2017.
- Evangelou E, Kerkhof HJ, Styrkarsdottir U, Ntzani EE, Bos SD, Esko T, Evans DS, Metrustry S, Panoutsopoulou K, Ramos YF, *et al*: A meta-analysis of genome-wide association studies identifies novel variants associated with osteoarthritis of the hip. *Ann Rheum Dis* 73: 2130-2136, 2014.
- Tsezou A: Osteoarthritis year in review 2014: Genetics and genomics. *Osteoarthritis Cartilage* 22: 2017-2024, 2014.
- Gee F, Rushton MD, Loughlin J and Reynard LN: Correlation of the osteoarthritis susceptibility variants that map to chromosome 20q13 with an expression quantitative trait locus operating on *NCOA3* and with functional variation at the polymorphism rs116855380. *Arthritis Rheumatol* 67: 2923-2932, 2015.
- Pierre S, Bats AS and Coumoul X: Understanding SOS (Son of Sevenless). *Biochem Pharmacol* 82: 1049-1056, 2011.
- Xiao Z, Li L, Li Y, Zhou W, Cheng J, Liu F, Zheng P, Zhang Y and Che Y: Rasfonin, a novel 2-pyrone derivative, induces ras-mutated Panc-1 pancreatic tumor cell death in nude mice. *Cell Death Dis* 5: e1241, 2014.
- Chang YL, Ho BC, Sher S, Yu SL and Yang PC: miR-146a and miR-370 coordinate enterovirus 71-induced cell apoptosis through targeting SOS1 and GADD45beta. *Cell Microbiol* 17: 802-818, 2015.
- Yue Y, Zhang M, Zhang J, Duan L and Li Z: SOS1 gene overexpression increased salt tolerance in transgenic tobacco by maintaining a higher K(+)/Na(+) ratio. *J Plant Physiol* 169: 255-261, 2012.
- Ariga H, Katori T, Yoshihara R, Hase Y, Nozawa S, Narumi I, Iuchi S, Kobayashi M, Tezuka K, Sakata Y, *et al*: Arabidopsis sos1 mutant in a salt-tolerant accession revealed an importance of salt acclimation ability in plant salt tolerance. *Plant Signal Behav* 8: e24779, 2013.

36. Kim JH, Nguyen NH, Jeong CY, Nguyen NT, Hong SW and Lee H: Loss of the R2R3 MYB, AtMyb73, causes hyper-induction of the SOS1 and SOS3 genes in response to high salinity in Arabidopsis. *J Plant Physiol* 170: 1461-1465, 2013.
37. Katschnig D, Bliek T, Rozema J and Schat H: Constitutive high-level SOS1 expression and absence of HKT1;1 expression in the salt-accumulating halophyte *Salicornia dolichostachya*. *Plant Sci* 234: 144-154, 2015.
38. Stade K, Ford CS, Guthrie C and Weis K: Exportin 1 (Crm1p) is an essential nuclear export factor. *Cell* 90: 1041-1050, 1997.
39. Engel K, Kotlyarov A and Gaestel M: Leptomycin B-sensitive nuclear export of MAPKAP kinase 2 is regulated by phosphorylation. *EMBO J* 17: 3363-3371, 1998.
40. Toone WM, Kuge S, Samuels M, Morgan BA, Toda T and Jones N: Regulation of the fission yeast transcription factor Pap1 by oxidative stress: Requirement for the nuclear export factor Crm1 (Exportin) and the stress-activated MAP kinase Sty1/Spc1. *Genes Dev* 12: 1453-1463, 1998.
41. Ferrigno P, Posas F, Koepp D, Saito H and Silver PA: Regulated nucleo/cytoplasmic exchange of HOG1 MAPK requires the importin beta homologs NMD5 and XPO1. *EMBO J* 17: 5606-5614, 1998.